


## Adenocarcinoma of the Esophagus and GE Junction

1

## Anatomy

Food moves from the throat

- esophagus
- stomach
- small bowel (jejunum)



2

## Cancer Staging

Staging refers to the tests to determine

- How large is the tumor?
- Has there been spread to lymph nodes?
- Has it spread to other parts of the body?

**Treatment options depend upon the cancer stage**

3


## Cancer Staging

- **T** = Tumor - Depth of growth into the wall
- **N** = Nodes - Spread to the lymph nodes
- **M** = Metastasis - Spread to liver, lungs, or bone

4

## Early Stage Cancers


Cancers start on the very inside layer called the mucosa



5

## Locally-advanced Cancers


Over time, cancers can grow into the muscular wall



6

### Lymph Nodes


In some cases, cancer cells can break off from the main tumor and spread to lymph nodes



7

### T Stage

Cancers are categorized based upon the thickness of the tumor, known as the T stage




8

### N Stage

Cancers are categorized by whether there is spread to the nodes.

- **N0** cancers have not spread to the nodes
- **N1** cancers have spread to the nodes.




9

### M Stage

Some cancers spread to other parts of the body

- **M0** cancers have not spread to other parts of the body
- **M1** cancers have spread to lungs, liver, or bone




10

### PET scan

Similar to CT scan  
Tracer shows 'hot spots'


- Cancer
- Inflammation or infection
- Normal organs (heart, kidneys)



11

### Endoscopic Ultrasound


- Similar to upper endoscopy (EGD)
- Ultrasound in scope
- Evaluates T stage



12

### Laparoscopy


- Some stomach cancers can spread inside the abdomen
- Areas of spread can be very small (grain of rice)
- Laparoscopy can detect spread inside the abdomen



13

### Laparoscopy

- General anesthetic
- Several 1/4" incisions
- Telescope examines the abdomen
- Biopsies can be performed.



14


### Treatment Plan

- Superficial (T1)
- Endoscopic Therapy
- Localized (T1b/T2)
- Surgery (esophagectomy)
- Locally-advanced (T3M0)
- Chemo±Radiation → Surgery
- Metastatic (M1)
- Chemotherapy

15

### Locally-advanced cancers

Patients with locally-advanced esophageal cancer often have localized spread of cancer cells in the surrounding area



16

### Locally-advanced cancers

If locally-advanced cancers are treated with surgery alone...



17

### Locally-advanced cancers

If locally-advanced cancers are treated with surgery alone...  
There is a risk that cancer cells can be left behind



18

### Preoperative Therapy

It is helpful to start with therapy *before* surgery that will shrink the cancer.



19

### Preoperative Therapy


It is helpful to start with therapy *before* surgery that will shrink the cancer.



20

### Preoperative Therapy


It is helpful to start with therapy *before* surgery that will shrink the cancer.



21

### Surgery after Preoperative Therapy


When surgery is then performed...



22

### Preoperative Therapy


When surgery is then performed...  
The risk of cancer recurrence is minimized



23

### Preoperative Therapy

When surgery is then performed...  
The risk of cancer recurrence is minimized



24

### Chemotherapy + Radiation CROSS Trial

363 patients with esophageal cancer studied

Patients were treated in two groups:

**Surgery Alone**

vs

**Chemo + Radiation** → Surgery

25

### Chemotherapy + Radiation CROSS Trial

363 patients with esophageal cancer studied

Chemotherapy + radiation together over 6 wks

**Surgery Alone**

vs

**Chemo + Radiation** → Surgery ⇒ Longer Survival

26

### Chemo + Radiation CROSS Trial

Typical schedule for chemotherapy + radiation:

- Chemotherapy once per week for six weeks
- Radiation five days per week for six weeks (28)
- PET scan (or CT) 4 weeks after the end of radiation
- Surgery 8 weeks after the end of radiation

27

### Chemo + Radiation - Side Effects

Kills cancer cells in the esophagus and lymph nodes

Can also irritate the lining of the esophagus.

Swallowing can be difficult the last 2 weeks.

Feeding tube may be needed for hydration/nutrition.

28

### Locally-advanced Adenocarcinoma

“Sandwich” chemotherapy before + after surgery:

Chemo (8 wks) → Surgery → Chemo (8 wks)

Two different drug combinations:

- FLOT (more effective)
- FOLFOX (better tolerated)

29

### “Sandwich” Chemotherapy Drugs

#### FLOT

- 5-FU
- Leucovorin
- Oxaliplatin
- Taxotere

#### FOLFOX

- 5-FU
- Leucovorin
- Oxaliplatin

30

### FLOT Treatment Plan

- FLOT Chemo every 2 weeks x 4 (=8 weeks total)
- Surgery (4-6 weeks later)
- FLOT Chemo every 2 weeks x 4 (=8 weeks total)

31

### Durvalumab Immunotherapy

Cancer cells can turn off immune cells using a protein PD-L1

Durvalumab turns on immune cells by counteracting PD-L1

Durvalumab strengthens the immune system to fight cancer

32

### Matterhorn Trial

The Matterhorn clinical trial compared two types of treatment:

FLOT → Surgery → FLOT

FLOT + Durvalumab → Surgery → FLOT + Durvalumab

Better survival with addition of durvalumab to FLOT

**Treatment with durvalumab depends upon approval from insurance company**

33

### FLOT Chemo ± Durvalumab

#### FLOT Chemotherapy

- FLOT (8 weeks)
- Surgery (4-6 weeks later)
- FLOT (8 weeks)

#### FLOT + Durvalumab

- FLOT + Durvalumab (8 wks)
- Surgery (4-6 weeks later)
- FLOT + Durvalumab (8 wks)
- Durvalumab monthly x10

34

### Tumor Biomarkers - Pathology Tests

Show whether other drugs may be helpful:

- HER-2 → Herceptin can be helpful
- PD-L1 → Immunotherapy can be helpful
- MMR → Immunotherapy can be very helpful

Biomarkers reported in a separate pathology report

Your medical oncologist will review these with you

35

### Adenocarcinoma Treatment Options

#### Chemo + Radiation

- Chemo + Radiation (6wk)
- Surgery

#### Chemotherapy

- Chemo (8wk)
- Surgery
- Chemo (8wk)

36

### Adenocarcinoma Treatment Options

#### Chemo + Radiation

- Longer track record
- Better tolerated
- Port usually placed
- Eating worse → better
- May need feeding tube

#### Chemotherapy

- More effective
- More side effects
- Port always required
- Eating slowly improves
- Less likely to need feeding tube

37

### Chemotherapy Administration

Most chemotherapy is administered by vein.

Several options exist to administer chemotherapy:

- Intravenous catheter in peripheral veins
- Peripheral Intravenous Central Catheter (PICC)
- Central Venous port

38

### Intravenous Catheter in Peripheral Vein ("IV")


- IV catheter placed in vein of hand or arm
- Allows administration of chemo and fluids
- Placed for each dose
- Removed that day
- Not suitable for FLOT chemo



39

### PICC Lines


- Placed in Radiology
- Stay in place during all of treatment
- Needs to be kept clean and dry
- Suitable for FLOT chemotherapy



40

### Central Venous Port


- Implantable device makes chemo easier
- May shower in 24 hrs
- No special care at home
- OK for FLOT chemo
- Allows for blood draws



41

### Central Venous Port


- Implanted under skin
- Neck incision (1/4")
- Incision below the collarbone
- Sutures dissolve
- "Superglue" on incisions



42

### Central Venous Port

When it is time for chemotherapy, a needle is inserted through the skin into the port



43


### Restaging

CT or PET scan will be performed after preoperative therapy

- Surgery performed after restaging
- Timing depends upon recovery from therapy

44

### Restaging after Chemotherapy ± Radiation




45

### Fitness Evaluation prior to Surgery




46

### Surgery vs No Surgery



47


### Surgery Advantages + Disadvantages



48




## Outcomes after Chemotherapy ± Radiation



49

## Active Surveillance



50

## Active Surveillance

Active Surveillance is appropriate when:

- CT/PET shows no signs of spread
- EGD biopsies are negative

Active Surveillance reduces the need for surgery by 50% *but* survival and disease control is probably better with immediate surgery.

51

## Primary Care Practitioner (PCP)

Critical to coordinate care between specialists.

We will update your PCP after each visit

PCP Referral Line (844) 235-6998

52

## My Atrium Patient Portal

- Critical to good communication with your care team
- Available for desktop or laptop or phone
- Sign up at [my.atriumhealth.org](http://my.atriumhealth.org)

53

## Exercise

- Reduces risk of complications from treatment
- Goal is 30min/day of vigorous exercise 6 days/wk
  - Working hard enough that you can't converse
  - Start slowly and build up
  - Every day counts! (Aim for daily activity)

54

### Smoking Cessation

- Smoking makes cancer treatment more difficult
  - Increases risk of complications after surgery
- Options for help with smoking cessation:
  - NC Quit Line 1-800-QUIT-NOW (1-800-784-8669)
  - American Lung Assn  
[www.freedomfromsmoking.org](http://www.freedomfromsmoking.org)
  - Smoking Cessation Counseling (Metro Charlotte)

55


### Protein Needs

- Men: Average 75 grams/day
  - Women: Average 60 grams/day
- Protein Shakes provide protein with minimal sugar



56

### Feeding Tubes



57

### Gastrostomy Tube

#### Feeding Gastrostomy

- Feeding with a syringe several times per day.
- Tube can be hidden underneath clothing
- Tube does not interfere with eating by mouth
- Removed easily in the office when no longer needed

58

### Gastrostomy Tube Methods

PEG: Tube placed by endoscopy  
 Laparoscopic: Tube placed surgically by laparoscopy  
 Preferred method depends upon whether esophagectomy is planned

59


### Gastrostomy Tube

- Outpatient Placement (go home the same day)
- Central venous port can be placed at the same time (if needed)

60

### Jejunostomy Tube

- Nutrition to bypasses the esophagus and stomach
- Placed in small intestine
- Pump administers feedings slowly
- Feeding done at night



61

### Jejunostomy Typical Regimen

- Jejunostomy tube feeds for 16 hours (6pm-10am)
    - Men: 75mL/hour x 16 hours = 5 cartons
    - Women: 60mL/hour x 16 hours = 4 cartons
  - Water 240ml (8oz) via syringe 4x/day
- Hospital nurses will teach use of the feeding tube

62

### Jejunostomy Feeds with Diabetes

Jejunostomy feedings elevate blood sugars

- Insulin may be required along with feeds
- Typical Pattern for tube feeds
- Feeds run via pump from 6pm to 10am
  - Insulin at 6pm (70/30 insulin)
  - Insulin at Midnight (70/30 insulin)
  - No insulin if tube feedings are not run

63

### Jejunostomy Video

A video is available to help become familiar with the feeding jejunostomy



64

### Surgery for Esophageal Cancer


Surgery for esophageal cancer is performed for:

- Superficial Tumors (T1) not removed by endoscopy
- Localized Tumors (T2 N0 M0)
- Locally Advanced (T3 M0) after preop therapy

65

### Goals of Surgery


- Remove tumor from esophagus
- Remove surrounding lymph nodes
- Create a new esophagus



66

### Ivor Lewis (Transthoracic) Esophagectomy


- Removes tumor and lower 1/3 esophagus
- Removes surrounding lymph nodes
- GI tract reconstructed



67

### Reconstruction


New esophagus is created from the stomach in the abdomen by fashioning it into a tube.



68

### Ivor Lewis esophagectomy


The new esophagus is now brought up into the chest. A connection is made between the esophagus and the stomach, called an *anastomosis*.



69

### Minimally-invasive Ivor Lewis


- Small incisions abdomen and chest
- Surgical telescope and instruments
- Smaller incisions → faster recovery and less discomfort



70

### Open Ivor Lewis


Minimally-invasive approach feasible in 95% of cases  
In some cases, an open approach is still necessary.



71


### Total Esophagectomy

For patients with tumors in the upper esophagus, we need to remove more of the esophagus  
We need to remove the whole esophagus, including the portion in the neck



72

### McKeown Esophagectomy



73

### Risks of Esophagectomy

Esophagectomy is a complex operation, with a real risk of complications.


Two significant complications:

- Anastomotic leak
- Pneumonia

74

### Anastomotic Leak

The anastomosis is surgical connection between the esophagus and the stomach.




75

### Anastomotic Leak

If healing doesn't occur:

- Leakage of fluid from the esophagus
- Infection in the space between the lungs
- Requires additional time in the hospital




76

### Anastomotic Leak

If leak occurs:

- Some leaks will seal
- Stent may be required to help healing
- Occasionally additional surgery is required




77

### Anastomotic Leak

Risk of leak depends on:


- Type of operation performed
- Nutritional status of patient
- Experience of the surgeon



78

### Pneumonia

- Occurs in 10-15% of patients after esophagectomy.
- Requires treatment with antibiotics
- Requires a longer hospitalization.



79

### Preventing Pneumonia

Several ways to help prevent pneumonia:



- Deep breathing
- Coughing
- Walking

After surgery, this means:

- Sitting in a chair most of the day
- Walking in the halls as soon as possible

80

### Minimally-invasive Esophagectomy



81

### Risks of Surgery

Risks related to anesthesia

- Heart attack (5%)
- Irregular heart rhythm (15%)
- Pneumonia (10%)
- Blood clots in legs (<5%)
- Pulmonary embolism (2%)

82

### Risks of Surgery

Risks related to Surgery

- Anastomotic leak (5%)
- Stricture at anastomosis (15%)
- Death within 90 days of surgery
  - Low risk patients = 2%
  - Intermediate risk = 10%
  - High risk = 30%

83

### Risks of Surgery

	Age <75	Age >75
Normal Muscle (75%)	2%	10%
Low Muscle (25%)	10%	30%

Risks of Death within 90 Days of Surgery

84

### Day Prior to Surgery

- Clear liquids for 24 hours prior to surgery
- Check with Pre-op nurse regarding medicines day prior to surgery
- No tube feedings the night before surgery

85

### Day of Surgery

- Arrive at 5am – nothing to eat or drink after midnight.
- Medicines OK w/ a sip of water
- sip of black coffee but **no cream**.
- Surgery will be cancelled if you have cream/milk
- Waiting room for family and friends on 5th floor

86

### Epidural Catheter for Pain Control

- Remains in place for 2-5 days
- Dosage can be adjusted as needed
- Can make it more difficult to urinate
- May require foley catheter in bladder
- Foley catheter removed after epidural removed

87

### Intensive Care Unit (ICU) (2-4 days)

- Surgical ICU on 11th floor
- NG tube in nose to drain stomach and esophagus
- Catheter in bladder
- Chest tube right chest
- Abdominal drains (usually 2)
- Feeding jejunostomy (usually stays in 8 wks)

88

### Intensive Care Unit (ICU)

- Bladder catheter removed → check that bladder empties properly
- Chest tube removed (day 2-4) → follow-up x-ray
- Fluid emptied from drains every few hours
- Start tube feedings by feeding
- Feeding jejunostomy (stays in 8 weeks)

89

### Ward - 6Tower

- Jejunostomy feeds started
- Up in a chair most of the day
- Walking in the halls
  - Start with assistance
  - Improves lung function
  - Prevents loss of muscle strength

90

### Jejunostomy Feeds


Jejunostomy tube feeds

- Start continuous (24 hours)
  - Convert to night-time only (16 hours)
- Water administered through feeding tube
- Usually 8oz 4 times/day
  - Important to prevent dehydration

91

### Jejunostomy Tube

- Nutrition to bypasses the esophagus and stomach
- Placed in small intestine
- Pump administers feedings slowly
- Feeding done at night



92

### Jejunostomy Typical Regimen

- Jejunostomy tube feeds for 16 hours (6pm-10am)
    - Men: 75mL/hour x 16 hours = 5 cartons
    - Women: 60mL/hour x 16 hours = 4 cartons
  - Water 240ml (8oz) via syringe 4x/day
- Hospital nurses will teach use of the feeding tube

93

### Jejunostomy Feeds with Diabetes

- Jejunostomy feedings elevate blood sugars
- Insulin may be required along with feeds
- Typical Pattern for tube feeds
- Feeds run via pump from 6pm to 10am
  - Insulin at 6pm (70/30 insulin)
  - Insulin at Midnight (70/30 insulin)
  - No insulin if tube feedings are not run

94

### Jejunostomy Video

A video is available to help become familiar with the feeding jejunostomy



95

### Activity after Surgery

- Up in chair most of the day
- Walking with help from nurse/Physical Therapist
- Goals:
  - Improve lung function
  - Prevent muscle loss

96



### Nasogastric (NG) Tube

Tube passed through nose into stomach

- Drains fluid from stomach
- Prevents vomiting

Upper GI X-ray on 2nd or 3rd day after surgery

- If stomach empties well → NG tube removed
- Otherwise, X-ray repeated 2-3 days later

97

### Swallowing Evaluation

Once NG tube has been removed:

Modified barium swallow in radiology

- Drink a white chalky liquid (barium)
- Look for proper swallowing function
- 70% of patients ⇒ liquids started by mouth

98

### Oral Intake at Home

Most are taking protein shakes when they go home

Protein shakes are started after tolerating water

- 2 oz per hour to start
- 4 oz per hour if 2oz are tolerated well

99

### Discharge

Goal: ready to leave day #6/7 after surgery

- Night-time tube feedings (6pm to 10am)
- Nutrition by mouth (70% of patients)
  - 1 oz of water per hour by mouth OR
  - Protein shakes 4oz every 2 hours
- Water through tube 8oz four times per day
- Home care nursing (feeding tube teaching)
- Home infusion (tube feeding supplies)

100

### Nutrition after Surgery

At discharge home:

- Protein shakes 4oz every 2 hrs
- Tube feeds 4-5 cans at night (6pm-10am)

10-12 Days: Increase protein shakes

- Tube feeds 3-4 cans at night

Three weeks: Post-esophagectomy Diet

8-12 weeks: Remove feeding tube (in office)

101

### Transition from Tube Feeds → Eating

Dietitian will calculate daily protein goal

- Typically 60-75 grams protein/day
- Each carton of tube feeding has 15 grams
  - 75 grams protein = 5 cartons/night
- More intake by mouth → tube feeds reduced
- Spread out protein during the day (20gm/meal)
- Three meals + 2-3 high-protein snacks

102

### Post-esophagectomy Diet

- Soft Consistency
- High Protein
- Avoid sugary liquids (can cause 'dumping')
- Avoid raw vegetables (and salads)
- Eating
  - Small, frequent meals
  - Sit up for 30-45 minutes after eating
  - Avoid eating within 2 hours of bedtime

103

### Medicines at Home - Pain

Acetaminophen (Tylenol) 1000mg 4x/day

Gabapentin 300mg 3 times/day

Oxycodone

- As needed in addition to Tylenol/gabapentin
- Will begin reducing dose at first postop visit
- Can usually discontinue by 4 weeks
- **NO DRIVING WHILE ON OXYCODONE**

104

### Non-steroidals Anti Inflammatory (NSAID)

Non-steroidal anti-inflammatories (Celebrex)

- 200 mg every 12 hours starting at 2 weeks

**NO GOODY POWDERS OR BCs**

- (Can cause permanent scarring at the surgery site)

105

### Acid Blockers = Proton Pump Inhibitors

Examples include omeprazole and pantoprazole

- Will stay on for at 1-2 years to prevent acid reflux
- Important in preventing scarring at anastomosis (new connection between esophagus and stomach)
- To administer through feeding tube, open capsule and resuspend beads in 60mL (2oz) of water

106

### Medicines at Home

Reglan – Helps stomach empty

- Will plan to stop after six weeks
- 0.1% risk of tardive dyskinesia (nervous tic)

Remeron – Helps improve appetite

- Can cause vivid dreams
- Used for several weeks after surgery
- Will stop within first three months of surgery

107

### Metoprolol = Beta Blockers

- Slows heart rate and lowers blood pressure

- Used to prevent rapid heart rate

- Patients not taking a beta blocker prior to surgery  
→ wean after surgery

- Patients taking a beta blocker prior to surgery →  
return to prior dose and drug after surgery

108

### Sleeping at Home

Reflux can occur the first few weeks/months after surgery

This improves over the first few months

A wedge pillow can be helpful for sleep



109

### Postoperative Visit at 7-10 Days

Check surgical site

- Remove staples (if needed)

Adjust medicines as needed

- Insulin (for diabetic patients on insulin)

- Reduce beta blocker medicines

Advance diet

Reduce tube feeds

110

### After surgery

Wean off medicines added after surgery

- Pain medicines
- Beta-blockers
- Reglan and Remeron

Continue acid blockers for at least 1 year

111

### Jejunostomy Removal

Jejunostomy tube is removed in the office once you can take in enough nutrients by mouth

Removal usually around 8 weeks after surgery

May take 30 minutes and some local anesthetic to loosen up the tube for removal.

112

### Nutritional Monitoring after Surgery

You may have difficulty absorbing some nutrients:

- Iron
- Vitamin B12
- Vitamin D

113

### Nutritional Monitoring after Surgery

About 3 months after the jejunostomy tube is removed, we will check blood levels:

- Iron (ferritin)
- Vitamin B12
- Vitamin D

114

### Nutritional Replacements after Surgery

Vitamin or iron replacements can be ordered by:

- Primary Care Provider (PCP)
- Medical Oncologist
- Surgeon

If levels are low

- Replacement
- Repeat testing in 3-6 months

115

### Team Members - Physicians

Primary Care Provider

Gastroenterologist

Medical Oncologist (chemotherapy)

Radiation Oncologist (radiation)

Surgeons

- Jonathan Salo MD
- Jeffrey Hagen MD
- Michael Roach MD

116

### Team Members - Support Staff

Dietitian - Liz Koch

Nurses

- Matthew Carpenter RN
- Brandon Galloway LPN

Navigator - Laura Swift

117

### Team Members - Physicians

Primary Care Provider

Gastroenterologist

Medical Oncologist (chemotherapy)

Radiation Oncologist (radiation)

Surgeons

- Jonathan Salo MD
- Jeffrey Hagen MD
- Michael Roach MD

118

### Team Members - Support Staff

Dietitian - Liz Koch

Nurses

- Matthew Carpenter RN
- Brandon Galloway LPN

Navigator - Laura Swift

119