

Locally-Advanced Esophageal Adenocarcinoma

Intro L 1

I'm Dr Jonathan Salo, a GI Cancer Surgeon at the in Charlotte, North Carolina If you're seeing this video, changes are you or someone close to you has had an encounter with esophageal cancer. These videos are designed to educate you about cancer and its treatment and help you and you cancer care team make the right decisions for you.

Of course, there is no substitute for the expert opinions of your cancer care team.

Locally-advanced Esophageal Cancer L 4. (2)

The topic of this video is locally-advanced adenocarcinoma of the esophagus and gastroesophageal junction.

Esophagus or Gastroesophageal junction 2. (3)

We will consider cancers of the esophagus and gastro-esophageal junction together, as the treatment is similar

Treatment is guided by the Stage of cancer (4)

The treatment recommended will depend upon the stage of the cancer. If you haven't already, this may be a good time to view the video on Diagnosis and Staging.

Staging (5)

A PET or CT scan is used for staging. If the scans show that the tumor is localized, without signs of spread beyond the nearby lymph nodes, it is considered M0.

On the other hand, if the scans show signs of spread, this is considered M1.

Metastatic Disease = M1 (6)

Patients with metastatic disease are best treated with chemotherapy and are generally not treated with surgery

Locally Advanced Esophageal Cancer 4 (7)

We will focus in this video on Locally-advanced Esophageal cancer

Locally-advanced Esophageal Cancer T2/3 or N+. (8)

Locally advanced esophageal cancer has not spread to other organs in the body. Locally-advanced cancers are not only M0, but they are either T2 or T3 or node-positive.

If this terminology is not familiar to you, please refer to our video on Diagnosis and Staging. There is a link above and in the description below.

Locally-advanced: M0 (9)

Locally-advanced cancers are M0, which means there is no evidence of metastasis, or spread to other parts of the body.

Adenocarcinoma vs Squamous Cell. (10)

To make things a bit more complicated, there are two types of esophageal cancer: Adenocarcinoma and Squamous Cell Carcinoma. The treatment of these cancers is somewhat similar, but different enough that we have prepared a video to specifically address squamous cell carcinoma.

Types of Esophageal Cancer (11)

This video will focus on locally-advanced *adenocarcinoma*

Why not just Cut it out? (12)

I am occasionally asked by patients why we don't just remove esophageal cancer with surgery as the first treatment.

The answer is a bit complex.

(13)

When a patient is diagnosed with cancer of the esophagus or gastroesophageal junction, it can look like the cancer is confined to the esophagus.

<esophageal cancer with 'fingers'> (14)

what we see from inside the esophagus is just the tip of the iceberg.

There is frequently spread of cancer cells into the nearby tissue or lymph nodes

(15)

If surgery is performed as the only therapy for locally-advanced cancer,

(16)

there is a risk that cancer cells can be left behind

Locally Advanced (17)

Preoperative therapy is administered before surgery And is designed to shrink the tumor

Locally Advanced (18)

and kill cancer cells in the surrounding area

This therapy typically involves chemotherapy

Locally Advanced (19)

and may involve radiation

Locally Advanced (20)

The overall goal is shrinking the tumor and killing nearby cancer cells, to make the surgery more effective

Surgery (21)

After preoperative therapy has shrunk the tumor and any other cancer cells, surgery can be performed

Post Surgery (22)

Preoperative therapy combined with surgery offers the best chance to remove all of the cancer, without leaving any behind

Post Surgery and Reconstruction (23)

...without leaving any cancer cells behind

Preoperative Therapy (24). 4

- Shrinks the tumor
- Make surgery more effective
- Leads to better cancer control than surgery alone

Neoadjuvant (25)

You hear preoperative therapy referred to as “neoadjuvant therapy”

Preop Therapy (26)

For adenocarcinoma of the esophagus, there are two different approaches to preoperative therapy.

One uses chemotherapy and radiation together, followed by surgery

Preop Therapy (27)

The other uses chemotherapy before and after surgery, without radiation therapy

Chemotherapy + Radiation -> Surgery (28)

The chemotherapy and radiation strategy uses chemotherapy and radiation together

Chemotherapy is given intravenously once per week

Radiation is given five days per week

Both are administered together for 5 1/2 weeks

This is also known as the "CROSS" treatment regimen

Chemotherapy + Radiation -> Surgery (29)

The chemotherapy is a low dose, designed to make the radiation more effective

The therapy is generally well tolerated

Hair loss is uncommon

Low blood counts are also uncommon, and occur in about 7% of cases

Esophagitis (30)

The radiation can, however, cause inflammation in the esophagus called esophagitis.

This can make eating more uncomfortable during the last two weeks of radiation

The good news is that this is temporarily and usually resolves within 2 or 3 weeks after the radiation is completed.

Esophagitis (31)

What this may mean, however, is placement of a temporary feeding tube may be needed.

Esophagitis (32)

If a feeding tube is necessary, it can usually be removed by 3 weeks after the end of radiation.

Chemo + Radiation -> Surgery (33) 2

To summarize, This approach to preoperative therapy uses six weeks of chemotherapy and radiation together.

Surgery is typically performed 6-8 weeks after the end of radiation, as soon as patients have recovered.

(4) Adenocarcinoma Preoperative Therapy (34). 4

The second and somewhat newer approach to preoperative therapy for esophageal adenocarcinoma is treatment with chemotherapy, followed by surgery, followed by additional chemotherapy

Chemotherapy is given every other week for 4 doses,

Followed by surgery,

Followed by chemotherapy every other week for another 4 doses

Chemo -> Surgery -> Chemo (35)

Chemotherapy is administered intravenously every other week for 4 doses over 8 weeks

Surgery is then performed 4-6 weeks later

Chemotherapy is again administered intravenously every other week for 4 more doses over 8 weeks

Chemotherapy Options (36)

There are two commonly used “recipes” for chemotherapy:

FLOT consists of four drugs

FOLFOX consists of three drugs

Chemotherapy Options (37)

As you might expect, FLOT is more effective against cancer, while FOLFOX has fewer side effects

Durvalumab Immunotherapy. (38)

Durvalumab is a drug which activates the immune system to fight cancer

The combination of FLOT and Durvalumab appears to be more effective against FLOT alone

Durvalumab is typically given along with FLOT chemotherapy, followed by monthly doses

Chemo -> Surgery -> Chemo (39) 2

To summarize, the chemotherapy strategy calls for 4 doses of chemotherapy before surgery, and an additional 4 doses of chemotherapy afterwards.

(2) Preoperative Therapy (40). 4

If we compare Chemotherapy + Radiation (or CROSS) with chemotherapy (FLOT)..

We see that Chemotherapy + Radiation has fewer side effects, but FLOT chemotherapy is more effective

A central venous port is required for FLOT chemotherapy

Due to radiation esophagitis, a feeding tube is a little more likely to be needed with chemotherapy and radiation

ChemoRT vs Chemo (41)

Your medical oncologist will guide you in the decision regarding chemotherapy + radiation vs chemotherapy alone.

In general, FLOT is recommended for younger and healthier patients

While chemotherapy and radiation is recommended for patients who may have difficulty tolerating FLOT

Preparing for Preoperative Therapy (42)

XXX

Preparing for Preoperative Therapy-2. (43)

XXX

Wrap up (44) 1

We hope you have found this video helpful. Here are some other topics for which videos have been posted or are planned.

Feel free to leave a comment or a question, or if you have suggestions for future videos.

If you or a family member have had an encounter with esophageal cancer, I would love to hear about your experience, so please take a minute to leave a comment below.

We're constantly creating new videos, so please subscribe to be notified of new videos when we post them.

The radiation can lead to irritation of the lining of the esophagus called radiation esophagitis. One can think of this as a sunburn on the inside of the esophagus. The result is that the therapy can make eating temporarily worse before it gets better. Usually the last two weeks of radiation therapy are the worst. By two weeks after radiation is completed, the tumor has shrunk enough that eating is usually better than before therapy.

For this reason, a temporary feeding tube is frequently necessary for patients getting chemotherapy and radiation

Preoperative Therapy (31)

On the other hand, FLOT chemotherapy, administered both before and after surgery, has been shown to be more effective than chemotherapy and radiation.

FLOT chemotherapy, however, has more side effects than chemotherapy and radiation.

As the chemotherapy begins to work, patients who have trouble eating before therapy find gradual improvement in their eating. As a result, a feeding tube might be less likely used with FLOT chemotherapy.

Preoperative Therapy (32)

So we have two different approaches for therapy prior to surgery, and each has favorable and unfavorable features

Preoperative Therapy (33)

In general, we think that FLOT may be more effective, but CROSS is better tolerated

Preoperative Therapy (34)

A medical oncologist, who specializes in chemotherapy treatment, will make a recommendation for preoperative therapy based upon a patient's age, overall health, and any medical problems such as heart disease or kidney disease.

Additional Topics (35)

The effectiveness of this approach was proven scientifically in the CROSS clinical trial

This was a scientific study published in 2010

363 patients with locally-advanced esophageal cancer were divided into two treatment groups:

One group was treated with surgery alone

The other group was treated with chemotherapy and radiation first, followed by surgery

CROSS Clinical Trial (25)

The results of the trial were quite dramatic: There was better control of the cancer in the group that received chemotherapy and radiation prior to surgery. This group had longer survival than the group treated with surgery alone.

CROSS Clinical Trial (26)

For the past 15 years, chemotherapy and radiation, followed by surgery, was established as the most effective therapy. This is known as trimodality therapy, because it uses a combination of three different therapies.

Matterhorn Trial

The Matterhorn clinical trial compared two types of treatment:

FLOT → Surgery → FLOT

FLOT + Durvalumab → Surgery → FLOT + Durvalumab

Better survival with addition of durvalumab to FLOT

Treatment with durvalumab depends upon approval from insurance company

FLOT Chemo \pm Durvalumab

FLOT Chemotherapy

- FLOT (8 weeks)
- Surgery (4-6 weeks later)
- FLOT (8 weeks)

FLOT + Durvalumab

- FLOT + Durvalumab (8 wks)
- Surgery (4-6 weeks later)
- FLOT + Durvalumab (8 wks)
- Durvalumab monthly x10