GI Stromal Tumors of the Stomach

GI Stromal Tumors

- · Arise from the wall of the stomach
- Grow slowly over time

2

- Lymph nodes rarely involved
- Not conventional stomach cancer

GIST vs Adenocarcinoma

GIST

1

Adenocarcinoma

- Starts from wall of stomach
- Slow growing
- Rarely spreads to lymph nodes
- Conventional "stomach cancer"
- Starts from lining of stomach
- Can spread to lymph nodes
- More aggressive

3

Benign or Malignant?

GIST tumors have a range of behavior:

- Small tumors tend to behave in a benign manner but can grow over time
- Larger tumors tend to behave in a malignant (cancerous) manner

GIST Treatment

- · Initial treatment usually surgery
- Gleevec pills after surgery for patients at high risk of recurrence
 - 1 to 3 years depending upon risk
- Large tumors treated with Gleevec before surgery

Patial Gastrectomy

GI Stromal Tumor can come in a variety of sizes

6



5

Partial Gastrectomy

- Tumor removed from
- Stomach wall closed
- · Lymph nodes not removed

7



Partial Gastrectomy

Risks of Partial Gastrectomy


- Leakage from closure of
- Bleeding requiring return to surgery
- Delayed stomach emptying



Distal Gastrectomy

Subtotal Gastrectomy

- Removes bottom half of the stomach
- Rarely required for GI **Stromal Tumors**



10


Distal Gastrectomy







• Removes bottom 2/3 of stomach

• Rarely required for GI **Stromal Tumors**





11















Risks of Surgery

- Leak where bowel is joined together (anastomosis)
- · Bleeding requiring reoperation
- Delayed stomach function
- Infection in the abdomen

Laparoscopy

- Some stomach cancers can spread inside the abdomen
- Areas of spread can be very small (grain of rice)
- Laparoscopy can detect spread inside the abdomen

20




19

Laparoscopy

A laparoscopy is performed under a general anesthetic.

- Several incisions 1/4" long
- A telescope is inserted to look inside the abdominal cavity.



Primary Care Practitioner (PCP)

A PCP is critical to coordinate care between specialists.

We will update your PCP after each visit
Call our referral line at (844) 235-6998 if you need a

21 22

My Atrium Patient Portal

- Critical to good communication with your care team
- Available for desktop or laptop or phone
- · Sign up at my.atriumhealth.org

Exercise

- Reduces risk of complications from treatment
- Goal is 30min/day of vigorous exercise 6 days/week
 - Working hard enough that you can't converse
 - Start slowly and build up
 - Every day counts! (Aim for some activity every day)

Smoking Cessation

- Smoking makes cancer treatment more difficult
 - Increases risk of complications after surgery
- Options for help with smoking cessation:
 - NC Quit Line 1-800-QUIT-NOW (1-800-784-8669)
 - American Lung Assn www.freddomfromsmoking.org
 - Smoking Cessation Counseling (Metro Charlotte)