


Adenocarcinoma of the Esophagus and GE Junction

1

Anatomy

Food moves from the throat
 → esophagus
 → stomach
 → small bowel (jejunum)



2

Cancer Staging

Staging refers to the tests to determine

- How large is the tumor?
- Has there been spread to lymph nodes?
- Has it spread to other parts of the body?

Treatment options depend upon the cancer stage

3


Cancer Staging

- **T** = Tumor - Depth of growth into the wall
- **N** = Nodes - Spread to the lymph nodes
- **M** = Metastasis - Spread to liver, lungs, or bone

4

Early Stage Cancers


Cancers start on the very inside layer called the mucosa



5

Locally-advanced Cancers


Over time, cancers can grow into the muscular wall



6

Lymph Nodes


In some cases, cancer cells can break off from the main tumor and spread to lymph nodes



7

T Stage

Cancers are categorized based upon the thickness of the tumor, known as the T stage




8

N Stage

Cancers are categorized by whether there is spread to the nodes.

- **N0** cancers have not spread to the nodes
- **N1** cancers have spread to the nodes.




9

M Stage

Some cancers spread to other parts of the body

- **M0** cancers have not spread to other parts of the body
- **M1** cancers have spread to lungs, liver, or bone




10

PET scan

PET scan is similar to CT scan


- Tracer shows 'hot spots'
 - Cancer
 - Inflammation or infection
 - Normal organs (heart)



11

Endoscopic Ultrasound


- Similar to upper endoscopy (EGD)
- Ultrasound in scope
- Evaluates T stage



12

Laparoscopy


- Some stomach cancers can spread inside the abdomen
- Areas of spread can be very small (grain of rice)
- Laparoscopy can detect spread inside the abdomen



13

Laparoscopy

- General anesthetic
- Several 1/4" incisions 1/4"
- Telescope examines the abdomen
- Biopsies can be performed.



14

Treatment Plan

Superficial (T1) \Rightarrow Endoscopic Therapy

Localized (T1b/T2) \Rightarrow Surgery


Locally-advanced (T3) \Rightarrow Chemo \pm Radiation
 \rightarrow Surgery

Metastatic (M1) \Rightarrow Chemotherapy

15

Locally-advanced cancers

Patients with locally-advanced esophageal cancer often have localized spread of cancer cells in the surrounding area



16

Locally-advanced cancers

If locally-advanced cancers are treated with surgery alone...



17

Locally-advanced cancers

If locally-advanced cancers are treated with surgery alone...
There is a risk that cancer cells can be left behind



18

Preoperative Therapy

It is helpful to start with therapy *before* surgery that will shrink the cancer.



19

Preoperative Therapy


It is helpful to start with therapy *before* surgery that will shrink the cancer.



20

Preoperative Therapy


It is helpful to start with therapy *before* surgery that will shrink the cancer.



21

Surgery after Preoperative Therapy


When surgery is then performed...



22

Preoperative Therapy

When surgery is then performed...
The risk of cancer recurrence is minimized



23

Chemotherapy + Radiation CROSS Trial

363 patients with esophageal cancer studied
Patients were treated in two groups:

Surgery Alone

vs

Chemotherapy + Radiation → Surgery

24

Chemotherapy + Radiation CROSS Trial

363 patients with esophageal cancer studied
Chemotherapy + radiation together over 6 wks

Surgery Alone

vs

Chemotherapy + Radiation → Surgery ⇒ Longer Survival

25

Chemotherapy + Radiation CROSS Trial

Typical schedule for chemotherapy + radiation:

- Chemotherapy once per week for six weeks
- Radiation five days per week for six weeks (28)
- PET scan (or CT) 4 weeks after the end of radiation
- Surgery 8 weeks after the end of radiation

26

Chemotherapy + Radiation - Side Effects

Kills cancer cells in the esophagus and lymph nodes
Can also irritate the lining of the esophagus.
Swallowing can be difficult the last 2 weeks.
Feeding tube may be needed for hydration/nutrition.

27

Locally-advanced Adenocarcinoma

“Sandwich” chemotherapy before + after surgery:

Chemo (8 wks) → Surgery → Chemo (8 wks)

Two different drug combinations:

- FLOT (more effective)
- FOLFOX (better tolerated)

28

“Sandwich” Chemotherapy Drugs

FLOT

- 5-FU
- Leucovorin
- Oxaliplatin
- Taxotere

FOLFOX

- 5-FU
- Leucovorin
- Oxaliplatin

29

Adenocarcinoma Treatment Options

Chemo + Radiation

- Chemo + Radiation (6wk)
- Surgery

Chemotherapy

- Chemo (8wk)
- Surgery
- Chemo (8wk)

30

Adenocarcinoma Treatment Options

Chemo + Radiation

- Longer track record
- Better tolerated
- Port usually placed
- Eating worse → better
- May need feeding tube

Chemotherapy

- More effective
- More side effects
- Port always required
- Eating slowly improves
- Less likely to need feeding tube

31

Chemotherapy Administration

Most chemotherapy is administered by vein.

Several options exist to administer chemotherapy:

- Intravenous catheter in peripheral veins
- Peripheral Intravenous Central Catheter (PICC)
- Central Venous port

32

Intravenous Catheter in Peripheral Vein ("IV")


- IV catheter placed in vein of hand or arm
- Allows administration of chemo and fluids
- Placed for each dose
- Removed that day
- Not suitable for FLOT chemo



33

PICC Lines


- Placed in Radiology
- Stay in place during all of treatment
- Needs to be kept clean and dry
- Suitable for FLOT chemotherapy



34

Central Venous Port


- Implantable device makes chemo easier
- May shower in 24 hrs
- No special care at home
- OK for FLOT chemo
- Allows for blood draws



35

Central Venous Port


- Implanted under the skin
- Neck incision (1/4")
- Incision below the collarbone
- Sutures dissolve
- "Superglue" on incisions



36

Central Venous Port

When it is time for chemotherapy, a needle is inserted through the skin into the port



37

Restaging

CT or PET scan will be performed after preoperative therapy

- Surgery performed after restaging
- Timing depends upon recovery from therapy

38

Primary Care Practitioner (PCP)

Critical to coordinate care between specialists.
We will update your PCP after each visit
PCP Referral Line (844) 235-6998

39

My Atrium Patient Portal

- Critical to good communication with your care team
- Available for desktop or laptop or phone
- Sign up at my.atriumhealth.org

40

Exercise

- Reduces risk of complications from treatment
- Goal is 30min/day of vigorous exercise 6 days/week
 - Working hard enough that you can't converse
 - Start slowly and build up
 - Every day counts! (Aim for daily activity)

41

Smoking Cessation

- Smoking makes cancer treatment more difficult
 - Increases risk of complications after surgery
- Options for help with smoking cessation:
 - NC Quit Line 1-800-QUIT-NOW (1-800-784-8669)
 - American Lung Assn
www.freedomfromsmoking.org
 - Smoking Cessation Counseling (Metro Charlotte)

42


Protein Needs

- Men: Average 75 grams/day
 - Women: Average 60 grams/day
- Protein Shakes provide protein with minimal sugar



43

Feeding Tubes



44

Gastrostomy Tube

Feeding Gastrostomy

- Feeding with a syringe several times per day.
- Tube can be hidden underneath clothing
- Tube does not interfere with eating by mouth
- Removed easily in the office when no longer needed

45

Gastrostomy Tube Methods

A gastrostomy tube can be placed either by endoscopy, which is called a PEG tube

A gastrostomy tube can also be placed by laparoscopy, which is usually preferred if surgery on the esophagus is planned in the future.

Your surgeon will help you decide which kind of tube is best for you. This is especially important if you will need esophageal surgery in the future, as the stomach is frequently used to make a new esophagus

46


Gastrostomy Tube

- Outpatient Placement (go home the same day)
- Central venous port can be placed at the same time (if needed)

47

Jejunostomy Tube

- Nutrition to bypasses the esophagus and stomach
- Placed in small intestine
- Pump administers feedings slowly
- Feeding usually done at night



48

Jejunostomy Typical Regimen

- Jejunostomy tube feeds for 16 hours (6pm-10am)
 - Men: 75mL/hour x 16 hours = 5 cartons
 - Women: 60mL/hour x 16 hours = 4 cartons
 - Water 240ml (8oz) via syringe 4x/day
- Hospital nurses will teach use of the feeding tube

49

Jejunostomy Feeds with Diabetes

Jejunostomy feedings elevate blood sugars

- Insulin may be required along with feeds

Typical Pattern for tube feeds

- Feeds run via pump from 6pm to 10am
- Insulin at 6pm (70/30 insulin)
- Insulin at Midnight (70/30 insulin)
- No insulin if tube feedings are not run

50

Jejunostomy Video

A video is available to help become familiar with the feeding jejunostomy



51

Surgery for Esophageal Cancer


Surgery for esophageal cancer is performed for:

- Superficial Tumors (T1) not completely removed by endoscopy
- Localized Tumors (T2N0)
- Locally Advanced (T3) after preoperative therapy.

52

Goals of Surgery


- Remove tumor from esophagus
- Remove surrounding lymph nodes
- Create a new esophagus



53

Ivor Lewis (Transthoracic) Esophagectomy


- Removes tumor
- Removes lower 1/3 of esophagus
- Removes surrounding lymph nodes
- Reconstruction of GI tract



54

Reconstruction


A new esophagus is created from the stomach in the abdomen by fashioning it into a tube.



55

Ivor Lewis esophagectomy


The new esophagus is now brought up into the chest. A new connection is made between the esophagus and the stomach, called an *anastomosis*.



56

Minimally-invasive Ivor Lewis


- Small incisions abdomen and chest
- Surgical telescope and instruments
- Smaller incisions → faster recovery and less discomfort



57

Open Ivor Lewis


We use the minimally-invasive approach in 95% of cases
In some cases, an open approach is still necessary.



58


Total Esophagectomy

For patients with tumors in the upper esophagus, we need to remove more of the esophagus
We need to remove the whole esophagus, including the portion in the neck



59


McKeown Esophagectomy



60


Colon Interposition

If the stomach is not suitable to make a new esophagus, the colon can be used to replace the esophagus



61

Colon Interposition



62

Risks of Surgery


An esophagectomy is a substantial operation, and in some cases there can be postoperative complications. We're going to talk about two of these complications and what you can do to reduce your risk of complications:

- Anastomotic leak
- Pneumonia

63

Anastomotic Leak

The anastomosis is surgical connection between the esophagus and the stomach.




64

Anastomotic Leak

If anastomosis doesn't heal:

- Leakage of fluid from the esophagus
- Infection in the space between the lungs
- Requires additional time in the hospital




65

Anastomotic Leak

If an anastomotic leak occurs:

- Some leaks will seal on their own
- A stent may be required to help healing
- Occasionally additional surgery is required




66

Anastomotic Leak

Risk of a leak depends upon:

- Type of operation performed
- Overall nutritional status of patient
- Experience of the surgeon




67

Pneumonia

Can occur in 10-15% of patients after esophagectomy.

Requires treatment with antibiotics and frequently requires a longer hospitalization.



68

Preventing Pneumonia

Several ways to help prevent pneumonia:



- Deep breathing
- Coughing
- Walking

After surgery, this means:

- Sitting in a chair most of the day
- Walking in the halls as soon as possible

69

Minimally-invasive Esophagectomy



70

Risks of Surgery

Risks related to anesthesia

- Heart attack (5%)
- Irregular heart rhythm (15%)
- Pneumonia (10%)
- Blood clots in legs (<5%)
- Pulmonary embolism (2%)

71

Risks of Surgery

Risks related to Surgery

- Anastomotic leak (5%)
- Stricture at anastomosis (15%)
- Death within 90 days of surgery
 - Low risk patients = 2%
 - Intermediate risk = 10%
 - High risk = 30%

72

Risks of Surgery

	Age <75	Age >75
Normal Muscle (75%)	2%	10%
Low Muscle (25%)	10%	30%

Risks of Death within 90 Days of Surgery

73

Day Prior to Surgery

- Clear liquids for 24 hours prior to surgery
- Check with Pre-op nurse regarding medicines day prior to surgery
- No tube feedings the night before surgery

74

Day of Surgery

- Arrive at 5am – nothing to eat or drink after midnight.
- Medicines w/ a sip of water (or black coffee) but **no cream**.
- Surgery will be cancelled if you have cream/milk
- Waiting room for family and friends on 5th floor
- Post-operative care in STICU (11th floor)

75

Epidural Catheter for Pain Control

- Remains in place for 2-5 days
- Dosage can be adjusted as needed
- Can make it more difficult to urinate
- May require foley catheter in bladder
- Foley catheter removed after epidural removed

76

ICU Stay (2-4 days)

- NG tube in nose to drain stomach and esophagus
- Catheter in bladder
- Chest tube right chest
- Abdominal drains (usually 2)
- Feeding jejunostomy (usually stays in 8 wks)

77

ICU

- Bladder catheter removed → check that bladder empties properly
- Chest tube removed (day 2-4) → follow-up chest x-ray
- Fluid emptied from drains every few hours
- Start tube feedings by feeding
- Feeding jejunostomy (stays in 8 weeks)

78

Ward - 6Tower

- Jejunostomy feeds started
- Up in a chair most of the day
- Walking in the halls
 - Start with assistance
 - Improves lung function
 - Prevents loss of muscle strength

79

Jejunostomy Feeds


Jejunostomy tube feeds

- Start continuous (24 hours)
 - Convert to night-time only (16 hours)
- Water administered through feeding tube
- Usually 8oz 4 times/day
 - Important to prevent dehydration

80

Jejunostomy Tube

- Nutrition to bypasses the esophagus and stomach
- Placed in small intestine
- Pump administers feedings slowly
- Feeding usually done at night



81

Jejunostomy Typical Regimen

- Jejunostomy tube feeds for 16 hours (6pm-10am)
 - Men: 75mL/hour x 16 hours = 5 cartons
 - Women: 60mL/hour x 16 hours = 4 cartons
 - Water 240ml (8oz) via syringe 4x/day
- Hospital nurses will teach use of the feeding tube

82

Jejunostomy Feeds with Diabetes

Jejunostomy feedings elevate blood sugars

- Insulin may be required along with feeds
- Typical Pattern for tube feeds
- Feeds run via pump from 6pm to 10am
 - Insulin at 6pm (70/30 insulin)
 - Insulin at Midnight (70/30 insulin)
 - No insulin if tube feedings are not run

83

Jejunostomy Video

A video is available to help become familiar with the feeding jejunostomy



84

Activity

- Up in chair most of the day
- Walking with help from nurse/Physical Therapist
- Goals:
 - Improve lung function
 - Prevent muscle loss

85

Nasogastric (NG) Tube

Tube passed through nose into stomach

- Drains fluid from stomach
 - Prevents vomiting
- Upper GI X-ray on 2nd or 3rd day after surgery
- If stomach empties well → NG tube removed
 - Otherwise, X-ray repeated 2-3 days later

86

Swallowing Evaluation

Once NG tube has been removed:

Modified barium swallow in radiology

- Drink a white chalky liquid (barium)
- Look for proper swallowing function
- 70% of patients ⇒ liquids started by mouth

87

Protein Shakes

Most are taking protein shakes when they go home

Protein shakes are started after tolerating water

- 2 oz per hour to start
- 4 oz per hour if 2oz are tolerated well

88

Discharge

Goal: ready to leave day #6/7 after surgery

- Night-time tube feedings (6pm to 10am)
- Nutrition by mouth (70% of patients)
 - 1 oz of water per hour by mouth OR
 - Protein shakes 4oz every 2 hours
- Water through tube 8oz four times per day
- Home care nursing (feeding tube teaching)
- Home infusion (tube feeding supplies)

89

Nutrition after Surgery

At discharge home:

- Protein shakes 4oz every 2 hrs
 - Tube feeds 4-5 cans at night (6pm-10am)
- 10-12 Days: Increase protein shakes
- Tube feeds 3-4 cans at night
- Three weeks: Post-esophagectomy Diet
- 8-12 weeks: Remove feeding tube (in office)

90

Transition from Tube Feeds → Eating

Dietitian will calculate daily protein goal

- Typically 60-75 grams protein/day
- Each carton of tube feeding has 15 grams
 - 75 grams protein = 5 cartons/night
- As intake by mouth increases, tube feeds are reduced

Spread out protein during the day (20gm/meal)

- Three meals + 2-3 high-protein snacks

91

Post-esophagectomy Diet

- Soft Consistency
- High Protein
- Avoid sugary liquids (can cause 'dumping')
- Avoid raw vegetables (and salads)
- Eating
 - Small, frequent meals
 - Sit up for 30-45 minutes after eating
 - Avoid eating within 2 hours of bedtime

92

Medicines at Home - Pain

Acetaminophen (Tylenol) 1000mg 4x/day

Gabapentin 300mg 3 times/day

Oxycodone

- As needed in addition to Tylenol/gabapentin
- Will begin reducing dose at first postop visit
- Can usually discontinue by 4 weeks
- NO DRIVING WHILE ON OXYCODONE

93

Non-steroidals Anti Inflammatory (NSAID)

Non-steroidal anti-inflammatories (Celebrex)

- 200 mg every 12 hours starting at 2 weeks

NO GOODY POWDERS OR BCs

- (Can cause permanent scarring at the surgery site)

94

Acid Blockers = Proton Pump Inhibitors

Examples include omeprazole and pantoprazole

- Will stay on for at 1-2 years to prevent acid reflux
- Important in preventing scarring at anastomosis (new connection between esophagus and stomach)
- To administer through feeding tube, open capsule and resuspend beads in 60mL (2oz) of water

95

Medicines at Home

Reglan – Helps stomach empty

- Will plan to stop after six weeks
- 0.1% risk of tardive dyskinesia (nervous tic)

Remeron – Helps improve appetite

- Can cause vivid dreams
- Used for several weeks after surgery
- Will stop within first three months of surgery

96

Metoprolol = Beta Blockers

- Slows heart rate and lowers blood pressure
- Used to prevent rapid heart rate
- Patients not taking a beta blocker prior to surgery → wean after surgery
- Patients taking a beta blocker prior to surgery → return to prior dose and drug after surgery

97

Sleeping

Reflux can occur the first few weeks/months after surgery
This improves over the first few months
A wedge pillow can be helpful for sleep



98

Postoperative Visit

Check surgical site

- Remove staples (if needed)

Adjust medicines as needed

- Insulin (for diabetic patients on insulin)
- Reduce beta blocker medicines

Advance diet

Reduce tube feeds

99

After surgery

Wean off medicines added after surgery

- Pain medicines
- Beta-blockers
- Reglan and Remeron

Continue acid blockers for at least 1 year

100

Jejunostomy Removal

Jejunostomy tube is removed in the office once you can take in enough nutrients by mouth

Removal usually around 8 weeks after surgery

May take 30 minutes and some local anesthetic to loosen up the tube for removal.

101

Nutritional Monitoring after Surgery

You may have difficulty absorbing some nutrients:

- Iron
- Vitamin B12
- Vitamin D

102

Nutritional Monitoring after Surgery

About 3 months after the jejunostomy tube is removed, we will check blood levels:

- Iron (ferritin)
- Vitamin B12
- Vitamin D

103

Nutritional Replacements after Surgery

Vitamin or iron replacements can be ordered by:

- Primary Care Provider (PCP)
- Medical Oncologist
- Surgeon

If levels are low

- Replacement
- Repeat testing in 3-6 months

104

Team Members - Physicians

Primary Care Provider

Gastroenterologist

Medical Oncologist (chemotherapy)

Radiation Oncologist (radiation)

Surgeons

- Jonathan Salo
- Jeffrey Hagen
- Michael Roach

105

Team Members - Support Staff

Dietitian - Liz Koch

Nurses - Brandon Galloway & Kit Sluder & Rebecca Wicks

Schedulers - Stacey Singleton & Tony Bethea

Navigator - Laura Swift

106