

Gastrostomy

I'm Dr Jonathan Salo, I'm a GI Cancer Surgeon in Charlotte, North Carolina.

Cancers in the GI tract can cause difficulty with eating, which can limit your body's ability to get the nutrition you need in order to keep your body healthy during cancer treatment.

Normally, food passes from the mouth into the esophagus, and then into the stomach. The stomach serves as a reservoir for food, to allow you to eat a big Thanksgiving meal. The stomach starts digestion, and then after the meal slowly allows small portions of food to pass into the small intestines, where most of the digestion occurs. Food passes from the stomach into the duodenum, then into the jejunum, which is the first portion of the small bowel.

A feeding tube can be used to administer

- formula, which is similar to Boost or Ensure.
- Water
- Medicines

There are two types of feeding tubes:

Gastrostomy tubes

- Placed into the stomach.
- Allow feeding with a syringe

Jejunostomy tubes

- Placed into the small intestine
- Require a pump for feeding

Gastrostomy is also referred to as a "G-Tube"

A jejunostomy is also referred to as a "J-Tube"

This video will focus on the gastrostomy tube

Our friend Brad has a gastrostomy tube, and we'll demonstrate how to administer water or formula.

The end of the gastrostomy tube has a special fitting with a cap.

To administer water or formula,

Open the cap

Attach the syringe to the fitting by twisting it on.

Empty the syringe

Remove the syringe

Close the cap.

This technique can be used for either water or formula or medicines

Another approach is to remove the plunger from the syringe, and use it as a funnel

The cap is opened,

The syringe attached to the fitting with a twisting motion

The water is simply poured into the syringe body and gravity will take it into the stomach.

When finished, simply remove the syringe and close the cap.

Formula can also be administered using this method.

Whatever method you use, it is extremely important to flush the gastrostomy tube with water after each use to avoid clogging the tube.

Some patients will need to take their medicines through the feeding tube.

Some medicines are available as a liquid, which makes administration easier.

Please check with your care team to see if any of your prescriptions can be converted from a pill to a liquid form.

In order to administer pills through a feeding tube, a pill crusher is indispensable.

Simply open the pill crusher by unscrewing the cap

Place the pill inside the container

Then screw the crusher cap down tightly against the tablet.

You will want to rotate the cap back and forth to crush the tablet and make a uniform powder.

When you remove the cap you will have a uniform powder which can be resuspended in water.

There are some medications which should not be crushed and cannot be given through a feeding tube.

Most enteric-coated medicines, or extended-release or time-release capsules should not be crushed

One common example is tamsulosin, also known as Flomax. This cannot be crushed as it will clog the feeding tube.

It is extremely important to flush the tube with at least 30mL (or at least one ounce) of water after each medicine to avoid clogging the tube.

Again, ask your care team for help in managing administration of medicines through the feeding tube.

Your dietitian will calculate how much formula you need. Typically this is 4 to 6 cartons of formula per day

You will also need to administer water through your tube, typically 8oz four times per day, or 32 oz per day.

Your dietitian will calculate how many cartons of tube feeds that you need, but typically it will consist of 4 to 6 cartons of formula per day.

You will also need water in addition to the tube feeding formula. Typically this is 8 ounces of water 4 times per day, for a total of 32 ounces of water per day.