

A user–friendly tool to evaluate the effectiveness of no–take marine reserves

Marine reserve evaluation

Juan Carlos Villaseñor-Derbez^{1¶*}, Caio Faro^{1¶}, Melaina Wright^{1¶}, Jael Martínez^{1¶}, Sean Fitzgerald^{1&}, Stuart Fulton^{2&}, Maria del Mar Mancha-Cisneros^{3&}, Gavin McDonald^{1,4,5&}, Fiorenza Micheli^{6&}, Alvin Suárez^{2&}, Jorge Torre^{2&}, Christopher Costello^{1,4,5¶}

¹ Bren School of Environmental Science and Management, University of California Santa Barbara, Santa Barbara, California, United States

² Comunidad y Biodiversidad A.C., Calle Isla del Peruano, Guaymas, Sonora, México

³ School of Life Sciences, Arizona State University, Tempe, Arizona, United States

⁴ Sustainable Fisheries Group, University of California Santa Barbara, Santa Barbara, California, United States

⁵ Marine Science Institute, University of California Santa Barbara, Santa Barbara, California, United States

⁶ Hopkins Marine Station and Center for Ocean Solutions, Stanford University, Pacific Grove, CA 93950, USA

*Corresponding author

Email: jvillasenor@bren.ucsb.edu (JCVD)

¶ These authors contributed equally to this work.

& These authors also contributed equally to this work.

Abstract

Marine reserves are implemented to achieve a variety of objectives, but are seldom rigorously evaluated to determine whether those objectives are met. In the rare cases when evaluations do take place, they typically focus on ecological indicators and ignore other relevant objectives such as socioeconomics and governance. And regardless of the objectives, the diversity of locations, monitoring protocols, and analysis approaches hinder the ability to compare results across case studies. Moreover, analysis and evaluation of reserves is generally conducted by outside researchers, not the reserve managers or users, plausibly thereby hindering effective local management and rapid response to change. We present a framework and tool, called “MAREA”, to overcome these challenges. Its purpose is to evaluate the extent to which any given reserve has achieved its stated objectives. MAREA provides specific guidance on data collection and formatting, and then conducts rigorous causal inference analysis based on data input by the user, providing real-time outputs about the effectiveness of the reserve. MAREA’s ease of use, standardization of state-of-the-art inference methods, and ability to analyze marine reserve effectiveness across ecological, socioeconomic, and governance objectives could dramatically further our understanding and support of effective marine reserve management.

Introduction

Unsustainable fishing practices threaten biodiversity, conservation, economic and social outcomes [1,2]. Marine Protected Areas (MPAs; and marine reserves, in which all extractive efforts are prohibited) are frequently proposed to aid in the recovery of fish and invertebrate stocks [3–6] by limiting or restricting fishing effort and gears.

~~Available empirical evidence on marine reserve effectiveness is mixed 7. Some studies have~~
~~shown~~ Empirical evidence shows that MPAs increase biomass [4,87], enhance resilience to

climatic impacts [9,108,9], and preserve genetic diversity [1110]. Compared to ~~partially~~
~~protected MPAs~~ MPAs that grant partial protection, marine reserves have higher levels of
biomass, density, richness, and larger organisms [3,12–1411–13]. ~~These~~ However, these effects
are often measured as biological changes ~~in the area through time~~ within the reserves through
time, and many lack a control site for comparison [1514]. This approach does not account for
other factors (*e.g.* system-level changes in productivity ; [15]) for which one must control [16]
in order to causally attribute a biological change to the reserve. ~~While some~~ Other studies
have used ~~control sites, these analyses do not estimate the effect of the reserve, and often use~~
a control–impact comparison approach that uses control sites but does not address temporal
variability [4,87,17–19]. A smaller fraction of studies have used a before–after–control–impact
(*i.e.* BACI) design comparing reserves to control sites before and after implementation
[4,20,21], which allows the use of causal inference techniques that estimate the effect of the
reserve.

But even when proper causal inference can be drawn, results are often ~~idiosyncratic~~ different
across reserves. Effects of reserves on ecological and economic outcomes are highly heteroge-
neous, and often depend on the specific ecological, economic, and social context. ~~The purpose~~
~~of this paper is to describe a user-friendly tool, called “MAREA”, to rigorously systematize~~
~~the evaluation of marine reserve effectiveness. The tool is in the form of an open-source~~
~~application that uses state-of-the-art methods from program evaluation to compare a reserve~~
~~to control sites along a number of ecological, economic, and governance dimensions.~~

~~The challenge of how to standardize~~ Standardization of marine reserve evaluation is not a
~~new one. The recent~~ new. The IUCN framework “How is your MPA doing?” [22,23] provides
a comprehensive list of biological, socioeconomic, and governance indicators, and insights
into how these ~~indicators~~ may be measured or collected. But this framework stops short
of analysis, ~~so and~~ provides a user with little guidance about establishing causal inference
about the reserve. Recent work ~~by Mascia et al.~~ [24] integrates these three dimensions via

the Social Ecological Systems Framework [25,26] and suggests the use of causal inference techniques to provide a measure of the effect of conservation interventions. However, ~~these two novel approaches do not~~ neither of these approaches provide a user-friendly tool that ~~enables~~ ensures replicability and scalability of the analysis, particularly when used by the fishers and decision makers themselves.

An increasingly popular way to make science accessible, reproducible, scalable, and replicable is through Open Science and the development of open-access tools [27]. The Ocean Health Index [28,29], for example, successfully standardized a way to measure the health and benefits of the oceans. This approach has been implemented at a global scale, but also at country-level [30], and regionally [31,32]. Open access tools are not limited to conservation, and have also been developed to evaluate fishery performance [33,34], design territorial use rights for fisheries [35], and improve decision making in the hydro power industry [36].

~~This paper presents a framework and~~ The purpose of this paper is to describe a user-friendly tool ~~to evaluate~~ , called “MAREA”, to rigorously systematize the evaluation of marine reserve effectiveness ~~, which incorporates the biological, socioeconomic~~ in terms of fisheries and marine conservation. The tool is in the form of an open-source application that uses state-of-the-art methods from program evaluation to compare a reserve to control sites along a number of ecological, economic, and governance dimensions ~~of any fishery~~. We first provide a list of commonly stated management objectives and match them to appropriate indicators. We then develop a simple approach to analyzing these indicators building on causal inference techniques [20], which help us understand the effect of management interventions [24,37]. To implement the analytical approach ~~in a user-friendly format~~, we introduce the Marine Reserve Evaluation Application (MAREA), an open source, web-based tool that automates the framework described in this paper and enables its broader use. Finally, we present a case study on the evaluation of a marine reserve established by the fishers of Isla Natividad (Mexico) in 2006, to demonstrate the potential of MAREA.

Materials and methods

Here, we describe the proposed framework to evaluate the effectiveness of marine reserves (Fig. 1). We explain how management objectives were identified and matched to appropriate indicators that allow the evaluation of the reserves, and provide brief guidelines on data collection. Alongside, methodologies to analyze these indicators are presented. We then describe the development of MAREA and explain how this ~~user-friendly~~ open-access tool can be used by fishermen, managers, and other stakeholders with little scientific background. Finally, we provide guidelines on how to interpret and use the results and output generated by MAREA to inform management.

Fig 1. Workflow to evaluate the effectiveness of marine reserves.

Marine Reserve objectives and indicators

Throughout this study, we will refer to the stated goals for which a marine reserve was designed as “objectives.” This work was motivated by the case of Mexico, where 39 reserves have been implemented over the past five years to achieve objectives such as increasing productivity in nearby waters or ~~recover~~ recovery of overexploited species; most of these reserves have never been formally evaluated for effectiveness at meeting those objectives. Thus, our focus was on identifying common objectives of marine reserves in Mexico. However, a literature review and discussions with marine reserve researchers ~~worldwide~~ suggest that the objectives driving Mexican marine reserve implementation are similar to those in the rest of the world. Thus, we group these objectives into seven major categories, which ~~can~~ may be applied to marine reserves worldwide. ~~The list of objectives includes stated objectives~~ Any given reserve may have been implemented to meet one or more of these. The list includes objectives stated in legislation [38,39] and official documents such as the Technical Justification Studies (*Estudios Técnicos Justificativos*), agreements, and decrees associated to these areas:

1. Avoid overexploitation
2. Conserve species under a special protection regime
3. Maintain biological processes (reproduction, recruitment, growth, feeding)
4. Improve fishery production in adjacent waters
5. Preserve biological diversity and the ecosystem
6. Recover overexploited species
7. Recover species of economic interest

Based on these seven objectives, we determined a set of associated indicators to evaluate reserve effectiveness. These indicators are specific variables on which data could be collected, and analyzed, to ultimately determine whether the corresponding objective was causally being achieved by the marine reserve. The list of indicators was compiled through a review of scientific literature in which we identified indicators that were used to measure similar objectives[3–5,7,11,13,14,17–20,22,23,40–43]. A first filter eliminated indicators for which baseline data do not typically exist in Mexico. The preliminary list of indicators was reviewed at a workshop with participation of members from Mexican fishery management agencies and non-government organizations. Later, these were presented to fishers from the Ensenada Fishing Cooperative (*S.C.P.P. Ensenada*), in El Rosario, Baja California, who provided input. Our final list of indicators includes those identified in review works [4,40,43].

Indicators are divided into three main categories: biological, socioeconomic, and governance (Table 1). The nine biological indicators focus on fish and invertebrate communities that are evaluated using underwater ecological surveys performed inside and outside the reserve (see Data and Analysis section for specific sampling design and methodologies). Five socioeconomic indicators reflect the performance of the fishery in terms of landings, income from landings, and availability of alternative livelihoods. Fifteen governance indicators describe the governance structures under which the community operates (*e.g.*, access rights to the fishery, number of fishers, legal recognition of the reserve). Some indicators are quantitative and require

a numerical entry (*e.g.* Fish biomass) while others are ~~more descriptive qualitative and~~
~~rely on a descriptive entry~~ (*e.g.* Reasoning for reserve location). Many of them specifically
 measure an outcome of the reserve, though some are designed to further the understanding
 of the mechanisms driving a reserve's performance. In that sense, most biological and
 socioeconomic indicators are outcome variables. On the other hand, governance indicators
 are viewed as possible explanatory variables of reserve performance. Whenever an indicator
 is applied to "Target species", it means that the indicator can be used for all species (*e.g.*
 Fish Biomass) and/or for individual species that are either the conservation target of the
 reserve or are of particular economic or ecological interest (*e.g.* Grouper Biomass). Finally,
indicators B3 and B4 are different in that B3 look only at the density of organisms above
size at first maturity (related to reproductive potential), while B4 measures the density of all
fish or of a target species. Each indicator targets different plausible desired outcomes, like
increased reproductive potential (*i.e.* B3; [44]) or having more fish -regardless of their size-
to attract tourism (*i.e.* B4). Table 1 presents the proposed indicators, and Table 2 shows how
 objectives are matched with biological and socioeconomic indicators. Governance indicators
are excluded from Table 2, as we suggest that all of them should be used in an evaluation,
regardless of the objectives.

Table 1: List of indicators to evaluate the effectiveness of no-take marine reserves.

Code	Indicator	Data type	Unit
Biological			
B1	Shannon diversity index	Continuous	
B2	Species richness	Discrete	Number of species/transect
B3	Density of mature organisms	Continuous	Percent
B4	Density*	Continuous	Organisms/transect
B5	Natural Disturbance	Descriptive	
B6	Mean Trophic Level	Continuous	
B7	Biomass*	Continuous	kg/transect
Socioeconomic			
S1	Total landings*	Continuous	kg
S2	Income from total landings*	Continuous	\$
S3	Alternative economic opportunities	Ordinal	
Governance			
G1	Access to the fishery	Categorical	
G2	Number of fishers	Discrete	
G3	Legal recognition of reserve	Binary	
G4	Reserve type	Descriptive	
G5	Illegal harvesting	Ordinal	
G6	Management plan	Binary	
G7	Reserve enforcement	Descriptive	
G8	Size of reserve	Discrete	
G9	Reasoning for reserve location	Descriptive	
G10	Membership to fisher organizations	Binary	
G11	Type of fisheries organizations	Categorical	
G12	Representation	Ordinal	
G13	Internal Regulation	Binary	
G14	Perceived Effectiveness	Categorical	
G15	Social Impact of Reserve	Categorical	

* The indicator is also applied to target species

Data and analyses

In many coastal marine reserves of Mexico, biological data are often collected via underwater visual censuses as part of a reserve’s monitoring program. Scientific divers ~~(which are often local fishermen with guidance from Civil Society Organizations; CSOs)~~ record fish and invertebrate richness and abundances, as well as fish total length along belt transects. Ecological surveys are typically performed annually in each reserve and corresponding control site(s), before and after the implementation of the reserve, providing a sampling design that can be used to draw causal inference. Control sites are areas where habitat is similar to that of the reserve, but with presence of fishing activity; in principle these are areas that are otherwise observationally identical to the reserve site, but where, for presumably random reasons, a reserve was not implemented. While transect dimensions (*i.e.* length and width) and sampling methods might vary from study to study, the general idea remains the same: richness, abundances, and sizes of organisms are recorded in a study-specific standardized way. For this reason, MAREA does not assume specific transect dimensions, and pertinent indicators are calculated per transect (Table 1). [More information on data collection and formatting is provided in a guidebook \[45\], which is available in English and Spanish in MAREA’s interface.](#)

This sampling design for biological data allows us to use causal inference techniques [20, 41–46] to evaluate the effect of the reserve on biological indicators. The hypothesis that the indicators will respond to implementation of the reserve is tested by analyzing spatial and temporal changes in each numeric biological indicator (all but B5) using generalized linear models [20]. To account for variations in the environment and survey conditions, covariates that are gathered during the underwater ecological surveys are included in the difference-in-differences model with form:

$$I_{i,t,z} = \beta_0 + \sum_{t=2}^T \gamma_t Y_t + \beta_1 Z_{i,z} + \beta_2 P_{i,t} \times Z_{i,z} + \beta_3 T_{i,t,z} + \beta_4 V_{i,t,z} + \beta_5 D_{i,t,z} + \epsilon_{i,t,z} \quad (1)$$

Table 2: Management objectives and respective performance indicators.

Objective	B1	B2	B3	B4	B4*	B5	B6	B7	B7*	S1	S1*	S2	S2*	S3
Avoid overexploitation			x		x	x			x	x	x	x	x	x
Conserve species under a special protection			x		x	x			x	x		x		x
Maintain biological process				x	x	x		x	x	x	x	x	x	x
Improve fishery production in nearby waters			x	x	x	x	x	x	x	x	x	x	x	x
Preserve biological diversity and the ecosystem			x		x	x			x		x		x	x
Recover overexploited species	x	x		x		x	x	x						x
Recover species of economic interest	x	x		x		x	x	x						x

* The indicator is applied to target species

Objective	B1	B2	B3	B4	B4*	B5	B6	B7	B7*	S1	S1*	S2	S2*	S3
Avoid overexploitation			x	x	x	x	x	x	x	x	x	x	x	x
Conserve species under a special protection			x		x	x			x	x		x		x
Maintain biological process	x	x		x		x	x	x						x
Improve fishery production in nearby waters				x	x	x		x	x	x	x	x	x	x
Preserve biological diversity and the ecosystem	x	x		x		x	x	x						x
Recover overexploited species			x		x	x			x		x		x	x
Recover species of economic interest			x		x	x			x		x		x	x

* The indicator is applied to target species

188 In this model, i , t , and z are indices for transect, time, and zone (control or reserve site),
189 respectively. This model allows us to estimate the change in an indicator (I) based on
190 the year (Y), a dummy variable that indicates treatment (Z ; *i.e.* control or reserve), an
191 interaction between a dummy variable that indicates ~~pre-or-post-implementation~~ before or
192 after implementation (P) and treatment (Z), and covariates such as bottom temperature
193 (T ; in °C), horizontal visibility during the survey (V ; in m), and depth at which survey was
194 performed (D ; in m). ϵ represents the error term associated to the regression. Here, years
195 are modeled as factors, using the first year as the reference level. This does not impose a
196 linear structure in the way an indicator changes through time (*i.e.* the change in biomass
197 between 2006 and 2007 does not have to be the same as the change between 2015 and 2016).
198 The treatment and implementation variables, modeled as dummy variables, are coded as
199 Control = 0 and Reserve = 1; and ~~Pre-implementation~~ Before implementation = 0 and
200 ~~Post-implementation~~ After implementation = 1, respectively.

201 Socioeconomic data are often collected by fishers, natural resource management agencies, or
202 CSOs by recording landings, income, and sometimes prices for each species. To control for
203 inflation, income is adjusted with the country's consumer price index [4247]:

$$I_t = RI \times \frac{CPI_t}{CPI_T} \quad (2)$$

204 Where I_t represents the adjusted income for year t as the product between the reported
205 income for that year and the ratio between the consumer price index (CPI) in that year to
206 the most recent year's (T) CPI. Since no control sites are typically available for this data
207 type, numeric socioeconomic indicators (S1 and S2) are evaluated with a simplified version of
208 eq. 1:

$$I_t = \beta_0 + \beta_1 P_t + \epsilon_t \quad (3)$$

While this model does not formally allow for causal inference, we can still measure changes in mean landings and income before and after the implementation of the reserve and provide valuable input. For both models (eq. 1 and eq. 3), we estimate the model ~~coefficients~~ coefficient with ordinary least squares, and calculate heteroskedastic-robust standard errors [48].

While biological and some economic data are regularly collected, governance data are typically not available nor systematically collected by the community or other organizations. Therefore, we created a survey specifically designed to collect information needed for the proposed indicators (B5, S3, and G1–G15). The survey is included as supplementary material in English (S1 Appendix) and Spanish (S2 Appendix). To analyze governance information, we developed a framework based on a literature review of common governance structures and their relation to effectiveness in managing fisheries or marine reserves (S3 Table). This approach has been proven to successfully evaluate governance structures [4349]. Unlike with biological and socioeconomic objectives (see eqs 1 and 3), MAREA does not quantitatively analyze governance information. Rather, it is presented along with the biological and socioeconomic indicators to provide managers and users with a more complete description of the reserve.

Marine Reserve Evaluation App (MAREA)

We developed MAREAI in R version 3.4.2 and R Studio 1.1.383 [4450] using the Shiny package [4551], to build an interactive web application hosted on an open server; the MAREA app can be accessed at turfeffect.shinyapps.io/marea. While the original version was developed in Spanish because it was aimed for Mexico and other Latin-American countries, all of its content can be translated by a translation widget available within the app.

MAREA is designed as a 6-step process, divided in tabs that appear upon launching the app. The first tab introduces the app and summarizes the evaluation process. Then, the user selects management objectives, which MAREA automatically matches to appropriate

indicators, based on Table 2. Users can also ~~manually modify selected~~ select and un-select indicators based on their interests and data availability by “clicking” on the check-boxes in MAREA. The user can then load data on one or more reserves, using standard *.csv text files; sample datasets are provided within MAREA. Once data have been loaded, MAREA identifies all reserves in the data, and lets the user select the reserve to be evaluated. At this point, the user can also specify the year of implementation of the reserve, reserve dimensions, and indicate target species that are of particular management interest. MAREA provides the user with a section to confirm that all the decisions made leading up to that point are correct. Once the user has confirmed all input data, objectives, and other information, MAREA performs the formal program evaluation analyses discussed above. For a typical data set, the automated analysis step takes less than one second. Finally, the user is taken to the results tab where all results are presented in a simple format. The user can also download a more comprehensive technical report produced in *.pdf format.

The first output is a color-coded scorecard intended to provide a general overview of the effectiveness of the reserve. The scorecard provides a global score for the reserve, a general score for each category of indicators, and an individual score for each indicator. The global and category-level scores are determined by the percentage of positive indicators, overall and for each category, respectively. For numeric biological indicators (all but B5), the color is defined by the sign of the interaction term coefficient (β_2) in eq. 1. For socioeconomic indicators, colors are assigned based on the direction of the slope (β_1) in eq. 3. Red, yellow, and green are used for $\beta_i < 0$, $\beta_i = 0$, and $\beta_i > 0$, respectively. The intensity of the color is defined by the significance of the coefficient, testing the null hypothesis of no change (*i.e.* $H_0 : \beta_i = 0$) with a Student’s t-test. Cutoff values are $p < 0.05$ and $p < 0.1$. Thus, even in a case where $\beta_i > 0$, if the coefficient is not significant by standard measures (*i.e.* $p > 0.1$), the indicator will be assigned a yellow color. A legend (Fig. 2) is provided within the scorecard to aid in the interpretation of these results. Governance indicators are represented simply by red or green. The color is defined based on what literature shows to be a negative (red) or

positive (green) factor for a reserve (S3 Table). For example, if the perceived degree of illegal fishing is high, this indicator will be assigned a red color. However, due to the nature of some governance indicators, which require the user to provide a narrative, only some indicators are presented in the scorecard (although all are included in the technical report).

Fig 2. Legend used to interpret the scorecard produced by MAREA. Colors indicate direction of change (red = negative; green = positive), and color intensity is given by the statistical significance.

The second output from MAREA is a technical report intended to communicate information and statistical results in a more comprehensive and technical way. This report also includes a scorecard as a summary of the results, but provides more information for each indicator. For all numeric biological indicators, the report includes a graph of the value of the indicator in the reserve and control sites through time. It also provides a regression table that summarizes the value of all coefficients in the regression and their respective robust standard errors. The summary table also provides information on model fit (R^2) and significance of the regression.

The scorecard is produced with functions from the Shinydashboard package [4652]. The technical report is produced by a parameterized Rmarkdown document [4753] processed by the knitr package [4854]. Another feature of MAREA is that the user can choose to share the data. Once the technical report is downloaded, the information on the reserve, its management objectives, and all uploaded data are saved into a central repository. These data can be accessed at any time by any person interested in acquiring them at github.com/turfeffect/MAREAdata.

Case study

While MAREA is a general tool that can be easily employed to evaluate the effectiveness of any marine reserve with the required input data, we illustrate its use here by applying it to

one marine reserve near Isla Natividad, in Baja California Sur, Mexico. Isla Natividad is located 8 Km off the Pacific Coast of the Baja California Peninsula (Fig. 3), where fishers operate under a fishing cooperative (*S.C.P.P. Buzos y Pescadores de la Baja California*) that promotes co-management of marine resources [49,50,55,56]. Additionally, fishers have Territorial Use Rights for Fisheries (TURFs) that provide them with exclusive access rights to exploit the benthic marine resources within a given perimeter [50,56].

Fig 3. General location of Isla Natividad (left) and map of the island (right). The marine reserve polygon is indicated in red, and the approximate location of control sites is indicated by blue squares (B = Babencho, D = La Dulce). [Shapefiles for Mexican coastline and the US were obtained from the Instituto Nacional de Estadística, Geografía e Informática of Mexico \(INEGI: \[www.inegi.org.mx/geo/contenidos/geoestadistica/m_geoestadistico.aspx\]\(http://www.inegi.org.mx/geo/contenidos/geoestadistica/m_geoestadistico.aspx\)\) and the tmap R package \[57\], respectively.](#)

In 2006, the Isla Natividad community implemented two community-based marine reserves within their TURF [9,51,52,40,58] [after establishing a baseline for the soon-to-be reserves and control sites](#). Evidence suggest that these reserves have been effective at enhancing resilience to climate variations [98] and preserving genetic diversity of high value commercial species such as abalone [41,10]. These ecological benefits have been translated into economic benefits, enhancing population persistence and bolstering abalone fisheries [53,42]. For the purpose of this evaluation, we focused on the “La Plana / Las Cuevas” marine reserve, located at the southern end of the island (Fig. 3) and its corresponding control site “La Dulce / Babencho”.

The objective of this reserve was to recover species of economic interest —which were overexploited— and to enhance fishery production in nearby waters. Fishers were also interested in preserving biological diversity and the ecosystem. Thus, objectives 4—7 were selected. Using Table 2 to match these objectives with appropriate management indicators, we selected all biological, socioeconomic, and governance indicators included as options in

the framework.

Local fishers (who were trained in scientific diving by the CSO Comunidad y Biodiversidad, A.C. (COBI; www.cobi.org), ReefCheck California, and Stanford University) and personnel from these institutions performed SCUBA dives to record fish and invertebrate richness and abundances, as well as fish total length. They recorded information along 30 m transects, with a sampling window of 2 m x 2 m following a standardized ReefCheck protocol [5459]. Ecological surveys were performed yearly in each reserve and corresponding control site(s), before and after the implementation of the reserve, providing the requisite time series data inside the reserve and for a suitable control site. Annual surveys (2006–2016) were carried out in late July – early August, performing a total of 242 and 245 transects in the reserve site for fish and invertebrate surveys, respectively. Similar sampling effort was applied to the control site, with 221 fish and 222 invertebrate transects. Between 12 and 27 transects were performed in each site every year.

Socioeconomic data were obtained from the National Commission for Aquaculture and Fisheries (*Comisión Nacional de Acuacultura y Pesca*; CONAPESCA). The data contains species-level information on monthly landings and income from nine species from 2000 to 2014. Data on landings and income were aggregated by year and species, and adjusted by the Consumer Price Index [4247]. From the nine species available, we selected as objective species those that contributed the most (88.27%) income from 2000 to 2014: lobster (*Panulirus interruptus*; 71.76%), red sea urchin (*Mesocentrotus franciscanus*; 9.33%), snail (*Megastrea undosa*; 3.93%), and sea cucumber (*Parastichopus parvimensis*; 3.23%). Abalone species (*Haliotis fulgens*; 4.52% and *Haliotis corrugata*; 6.16%) were excluded because the cooperative implemented an informal closure of these fisheries in 2010 to allow the population recover. Eliminating all fishing pressure on abalones means that the control site receives (for this species) the same treatment as the reserve.

We constructed the governance data based on local knowledge of the area and the community.

Results from illustrative example

In this section we show the results of the application of MAREA to the La Plana/Las Cuevas marine reserve in Isla Natividad, Mexico. These results are intended to highlight the relevance and utility of the MAREA framework and app, which automate the analysis and make it replicable. While we highlight some of the general observed trends, we focus on the utility of the tool rather than on the specific effectiveness of this case study marine reserve.

The scorecard (Fig. 4) shows that this reserve achieves a general score of 64%, suggesting that 64% of all indicators are positive. All category-level scores were also high, with values of 67%, 60%, and 71% positive indicators for biological, socioeconomic and governance, respectively.

Fig 4. Scorecard produced by MAREA for the “La Plana / Las Cuevas” marine reserve in Isla Natividad, Mexico.

Among the biological indicators, the greatest effect of the reserve was observed for snail and sea cucumber densities, with values of $\beta_2 = 97.17$ ($p < 0.05$) and $\beta_2 = 2.31$ ($p < 0.05$), respectively. Fish indicators showed no significant change ($p > 0.1$), with negative trends for Shannon’s diversity index and fish species richness and positive trends for density, biomass, and mean trophic level. Changes through time for these indicators are presented in Figure 5, and a summary of β_2 coefficients is provided in Table 3.

Table 3: Summary of average treatment effect of the reserve on biological indicators.

Indicator	Estimate (SD)	t-score
Shannon fish	-0.22 (0.16)	-1.3969
Richness fish	-0.61 (0.43)	-1.4073
Density fish	0.74 (6.15)	0.1205
Trophic fish	0.00 (0.01)	0.1399
Biomass fish	0.22 (1.47)	0.1476
Shannon invert	-0.67 (0.22)**	-3.0481
Richness invert	-2.71 (0.81)**	-3.3519
Density invert	91.21 (47.11)*	1.9362
Lobster	7.66 (8.93)	0.8583
Urchin	2.15 (1.23)*	1.7425
Snail	97.17 (42.90)**	2.2652
Cucumber	2.31 (1.17)**	1.9782

* Asterisks indicate significance level, with (*) indicating $p < 0.1$ and (**) $p < 0.05$.

Fig. 5 Plots for values of each biological indicator (y-axis) through time (x-axis).

Red and blue correspond to the reserve and control sites, respectively. Black lines indicate yearly mean values, and ribbons indicate ± 1 standard error. Dots are horizontally jittered to aid visualization. This figure contains information for fish Shannon’s diversity index (a), fish species richness (b), fish density (c), fish trophic level (d), fish biomass (e), invertebrate Shannon’s diversity index (f), invertebrate species richness (g), invertebrate density (h), lobster density (i), urchin density (j), snail density (k), and sea cucumber density (l).

One of the main objectives of this reserve was to increase landings. Results of the socioeconomic indicators show that total landings were, on average, 64.20 metric tonnes higher ($p > 0.1$) after the implementation of the reserves, though this cannot necessarily be interpreted as causal, because it relies entirely on a before–after comparison. Total income was \$10,344.85 ($p < 0.05$) thousands of Mexican Pesos (K MXP) higher after the implementation of the reserves. On average, lobster and sea cucumber landings increased, while urchin and snail landings and income decreased. Figure 6 presents the changes in these indicators through time, and Table 4 summarizes these results.

Table 4: Summary of differences in socioeconomic indicators before and after the implementation of the reserve.

Indicator	Estimate (SD)	t-score
Landings	64.20 (90.07)	0.7127
Income	10344.85 (3982.20)**	2.5978
Lobster landings	7.37 (13.95)	0.5281
Urchin landings	-30.00 (9.49)**	-3.1620
Snail landings	-69.53 (33.82)*	-2.0561
Cucumber landings	9.34 (6.72)	1.3906
Lobster income	14372.85 (3634.64)**	3.9544
Urchin income	-5800.46 (1867.50)**	-3.1060
Snail income	-404.85 (187.07)**	-2.1641
Cucumber income	131.49 (185.66)	0.7082

* Asterisks indicate significance level, with (*) indicating $p < 0.1$ and (**) $p < 0.05$.

Fig. 6 Plots for values of each socioeconomic indicator (y-axis) through time (x-axis). Red and blue correspond to before and after the implementation of the reserve, respectively. This figure contains information for total landings (a), total income (b), lobster landings (c), urchin landings (d), snail landings (e), sea cucumber landings (f), lobster income (g), urchin income (h), snail income (i), and sea cucumber income (j).

Recall that the governance objectives are evaluated based on the institutions present, not on a specific quantitative linkage between governance and biological or economic outcomes. Data for this reserve suggest that the community is strongly organized, which is a likely driver of the successes reported above [5560]. The first point of success is the existence of a fishing cooperative that is also affiliated with a regional federation of cooperatives. These polycentric governance structures allow various levels of organization that have been shown to foster communication and cooperation [43,5049,56]; federations also provide bargain power with governments [43,5649,61]. Access to fishing resources is managed through a TURF, permits, and fishing quotas (for some species). McCay [4955] suggests that the TURF promotes a sense of stewardship of their resources and incentivizes sustainable management. Together, these structures enabled a participative, bottom-up process during the reserve design phase;

385 opinions of all fishing members —and often non-fishing community members— were included.
386 Participation of community members in reserve surveillance and yearly monitoring indicate
387 commitment and interest, and allow informal communication of results to un-involved
388 community members. Furthermore, the reserve is partially isolated from poaching activity
389 and fishers have internal regulations pertaining to the reserves. The low level of illegal fishing
390 by members of the community and outsiders both inside and outside the reserve represents
391 another indication of effectiveness. Governance indicators are summarized in Table 5.

Table 5: Summary of governance indicators.

Indicator	Description
Access to the fishery	Permits, Territorial Use Rights for Fisheries, Quotas (for some fisheries)
Number of fishers	Stable
Legal recognition of reserve	Not recognized
Reserve type	Community-based Marine Reserve
Illegal harvesting	Due to its relative isolations, neither the reserve or TURF suffer from significant illegal harvesting
Management plan	The reserve does not have a management plan, but written rules exist within the cooperative
Reserve enforcement	Fishers have two land stations equipped with radars and patrol boats 24/7 to patrol the reserves.
Size of reserve	The reserve is big enough to protect the targeted sessile or not highly mobile invertebrates (lobster, urchin, snail, cucumber, and abalone)
Reasoning for reserve location	The reserves were put in place in zones that, according to local knowledge, were once very productive. Habitat heterogeneity and ease of monitoring, surveillance and enforcement were also considered.
Membership to fisher organizations	The fishers are part of fisher organizations.
Type of fisheries organizations	The fishers are part of a cooperative (S.C.P.P. Buzos y Pescadores de la Baja California) and are affiliated to a federation (FEDECOOP).
Representation	Reserves were designed by fishers in a bottom-up approach, incorporating expertise from academics and CSO members. This was a highly inclusive and participatory process.
Internal Regulation	Fishers have stringent internal regulations to control fishing effort throughout their TURF, assigning different fishing zones and gears to different teams. Rules pertaining the marine reserves also exist.
Perceived Effectiveness	The fishers have a positive perception about the effectiveness of their reserve, often stating that they have seen significant economic benefits.
Social Impact of Reserve	The reserves have had a significant positive social impact. Fishers are proud to be an example of successgul marine conservation, allowing them to have increased social capital.

Discussion

We have developed and presented ~~a user-friendly, an~~ automated approach for evaluating the effectiveness of marine reserves in Mexico, and perhaps around the world. Here we highlight MAREA’s utility for evidence—based management, and comment on a few of its shortcomings. The findings from Isla Natividad are used purely to validate the relevance of MAREA rather than to discuss particularities of the marine reserve effectiveness, which has been described before [9,11,538,10,42]. We use examples from the case study to build on the utility of MAREA and discuss ways in which results can be interpreted to inform management.

The causal inference techniques used by MAREA have been suggested [37,~~41~~46] and used [20] before in other ad hoc studies. This approach reduces ambiguity in the interpretation of results. For example, invertebrate density decreased through time inside and outside of the reserve (Fig. 5h). In this case, a before–after evaluation of the reserve (*i.e.* ignoring the control site) would have incorrectly concluded that the reserve failed to protect invertebrates. On the other hand, a control–impact approach (*i.e.* compare reserve vs. control site only in 2016) would have identified higher densities inside the reserve, concluding that the reserve increases invertebrate density. However, by executing a formal difference–in–differences approach for causal inference, MAREA identifies the changes through time and across sites, and estimates the effect of the reserve on density at $\beta_2 = 91.21$ ($p < 0.05$). This approach reveals that invertebrate densities decrease in both sites through time, but the decrease is faster for the control site, thus yielding a positive value for β_2 .

The approach used by MAREA to estimate the effect of the reserve on biological indicators requires cautious interpretation of the results. The value of the β_2 coefficient represents the difference between the temporal trends of the reserve and control sites [20]. As exemplified by the case of invertebrate densities, a positive value (*i.e.* $\beta_2 > 0$) does not necessarily indicate an increase in the indicator through time, but rather a positive difference with respect to the

temporal trend of the control site. The inverse occurs for negative values of β_2 .

MAREA provides in-depth analysis and a convenient snapshot overview of the effect of the reserve, allowing users to rapidly identify trends. However, users must interpret multiple indicators at a time to better understand the results. For example, with additional knowledge of local environmental variability (*i.e.* indicator B5: Natural Disturbance), we can better understand the trends in invertebrate densities. As reported before [98], hypoxic conditions that have occurred in Isla Natividad can cause decreases in invertebrate densities, and reserves buffer the negative effect. While MAREA automates the analysis and makes results replicable, proper interpretation will still depend on the user. Results produced by MAREA can only aid in management and decision making when results have been correctly interpreted.

Socioeconomic and governance indicators typically lack a control site, which impede us from using the causal inference techniques employed to measure biological changes [24]. However, we can still extract useful information from them. Again, by combining results from multiple indicators, MAREA can provide insights into the effect of the reserve. For example, lobster and sea cucumber have shown increases in densities, landings, and income. We cannot conclude that landings and income from these species have increased due to the reserve, but we can at least conclude that landings have not decreased. While further information on market behavior of each fishery is needed, these results provide insights into the state of the reserve and its associated fisheries.

As for the governance information, it is difficult to establish causal links between the state of the reserve and the governance structures present in the community. However, providing a single platform (*i.e.* scorecard) or document (*i.e.* technical report) where biological, socioeconomic, and governance information is comprehensively included can aid in management. By using MAREA, this information will be reported across reserves in a standardized way, and can help managers identify overarching patterns across sites.

By making results straightforward to interpret, MAREA may also assist in communication

with a broader stakeholder community. While stakeholder involvement in the design and implementation phases of marine reserves is important, that may not be sufficient for ensuring long-run buy-in or success. The scorecard is easily understandable by experts and non-experts, and can be used as an effective tool for communicating the results of annual evaluations. Additionally, the technical report can serve as a tool for managers and scientists to rapidly produce and communicate information at a more technical level.

We recognize that the seven objectives and 29 indicators used by MAREA might not fully describe a reserve. ~~However, they~~ In order to ensure the applicability of the tool to reserves in other countries, further testing in other regions should take place. However, the proposed objectives and indicators provide a starting point to perform the evaluation, to which managers and users can add other indicators (*e.g.* larval dispersal or connectivity) that are relevant to their reserve. Furthermore, MAREA's value is that it provides a free, simple, and replicable way to perform rigorous impact analysis. The tool can easily be used by fishers, CSO members, and managers in government agencies, providing transparency of the analysis and results. In addition, it can empower and enable local managers and fishers to respond to local change and adapt by allowing direct and easy access to the information.

An evident limitation of MAREA is its dependence on data obtained through a BACI design, and the amount of samples needed to estimate coefficients in Eqn. 1. It is not uncommon for control sites or baselines that describe the biological state of the area before the management intervention to be absent. Properly designing marine reserves by identifying control sites and establishing a baseline before the implementation of the reserve is enough to overcome this issue; reserves for which there is no control site and baseline can not be evaluated with MAREA. On the second case, typical underwater surveys require that at least 12 - 16 transects are performed on each site (*i.e.* reserve and control) each year. This provides with at least 48 samples (12 samples per site, per year), enough to avoid overfitting Eqn 1. On either case, these problems can be easily avoided by accounting for data requirements in the

eventual evaluation, an important phase during the design process.

To the best of our knowledge, MAREA is the first tool designed to evaluate marine reserves. Previous work [22,24] addressed MPA evaluation and provided the foundation for our contribution. However, these did not intended to create user-friendly tools to aid in the evaluation. As with other conservation management tools, development of tools that automatize complex calculations can have an important impact in management [62]. The use of open data science enables the creation of open-access tools that can address technical gaps and improve management [27].

The effectiveness of marine reserves continues to be a matter of debate [7,12,40,11,43,63]. With current targets set to increase ocean protection, it is important that we understand the effects of our interventions [37] so we can better inform management [41,46]. It is therefore important that academics, managers, fishers, and CSOs have access to open access tools like MAREA. This is particularly relevant for Mexico and other Latin American countries, where management agencies are often understaffed and underfunded [57,64], or where materials are often not available in their language. In this context, MAREA provides a simple and replicable way to align management objectives with performance indicators. The proposed methodologies, especially the way in which biological indicators are evaluated, provide valuable information for managers. We acknowledge there is room for improvement in the way in which socioeconomic and governance data are analyzed. Despite this, providing a unifying platform where all indicators can be analyzed and comprehensively presented represents a valuable step towards effective evidence—based management [41,46].

The first release of MAREA is now available, and it will continue to be developed and maintained to keep up to date with the literature. This process will incorporate new features, and enhance current ones, aiming to improve user experience and expand the scope of the analysis. Other modifications may also include addition of more objectives and indicators to ensure applicability in other regions, full translation into other languages

to avoid any ambiguities introduced via the automatic translation, or reporting effects over time in percentages to aid interpretation. Yet, we believe that this first release represents a major step towards effective, replicable evaluation and management of marine reserves.

Acknowledgements

We thank Olivier Deschenes and Andrew Plantinga, who provided valuable input to design the model that evaluates the biological indicators. Special thanks to the fishers from Isla Natividad, who gathered the data used in this study, and the fishers from El Rosario, who helped us validate our survey and framework, and to Arturo Hernández and Alfonso Romero who provided help with the logistics. Finally, we thank the Editor for a professional handling, and two anonymous reviewers for their suggestions, which significantly improved the quality of this paper.

References

1. Pauly D, Watson R, Alder J. Global trends in world fisheries: Impacts on marine ecosystems and food security. *Philos Trans R Soc Lond, B, Biol Sci.* 2005;360: 5–12. doi:10.1098/rstb.2004.1574
2. Halpern BS, Walbridge S, Selkoe KA, Kappel CV, Micheli F, D’Agrosa C, et al. A global map of human impact on marine ecosystems. *Science.* 2008;319: 948–952. doi:10.1126/science.1149345
3. Lester S, Halpern B. Biological responses in marine no-take reserves versus partially protected areas. *Mar Ecol Prog Ser.* 2008;367: 49–56. doi:10.3354/meps07599
4. Lester S, Halpern B, Grorud-Colvert K, Lubchenco J, Ruttenberg B, Gaines S, et al. Biological effects within no-take marine reserves: A global synthesis. *Mar Ecol Prog Ser.* 2009;384: 33–46. doi:10.3354/meps08029
5. Sala E, Costello C, De Bourbon Parme J, Fiorese M, Heal G, Kelleher K, et al. Fish banks: An economic model to scale marine conservation. *Marine Policy.* 2016;73: 154–161. doi:10.1016/j.marpol.2016.07.032
6. Hastings A, Gaines SD, Costello C. Marine reserves solve an important bycatch problem in fisheries. *Proc Natl Acad Sci U S A.* 2017;doi: [114: 8927–8934](#). doi:10.1073/pnas.1705169114
- ~~7. Padleton L, Aghmadian G, Browman H, Thurstand R, Kaplan D, Bartolino V. Debating the effectiveness of marine protected areas. *ICES Journal of Marine Science.* 2017; doi: [8: 7](#).~~
- Aburto-Oropeza O, Erisman B, Galland GR, Mascareñas-Osorio I, Sala E, Ezcurra E. Large recovery of fish biomass in a no-take marine reserve. *PLoS ONE.* 2011;6: e23601. doi:10.1371/journal.pone.0023601
- ~~9.~~ [8.](#) Micheli F, Saenz-Arroyo A, Greenley A, Vazquez L, Espinoza Montes JA, Rossetto M, et al. Evidence that marine reserves enhance resilience to climatic impacts. *PLoS ONE.*

2012;7: e40832. doi:10.1371/journal.pone.0040832

~~10.~~9. Roberts CM, O’Leary BC, McCauley DJ, Cury PM, Duarte CM, Lubchenco J, et al.
Marine reserves can mitigate and promote adaptation to climate change. *Proc Natl Acad Sci*
U S A. 2017;114: 6167–6175. doi:10.1073/pnas.1701262114

~~11.~~10. Munguía-Vega A, Sáenz-Arroyo A, Greenley AP, Espinoza-Montes JA, Palumbi SR,
Rossetto M, et al. Marine reserves help preserve genetic diversity after impacts derived from
climate variability: Lessons from the pink abalone in baja california. *Global Ecology and*
Conservation. 2015;4: 264–276. doi:10.1016/j.gecco.2015.07.005

~~12.~~11. Edgar GJ, Stuart-Smith RD, Willis TJ, Kininmonth S, Baker SC, Banks S, et
al. Global conservation outcomes depend on marine protected areas with five key features.
Nature. 2014;506: 216–220. doi:10.1038/nature13022

~~13.~~12. Giakoumi S, Scianna C, Plass-Johnson J, Micheli F, Grorud-Colvert K, Thiriet P,
et al. Ecological effects of full and partial protection in the crowded mediterranean sea: A
regional meta-analysis. *Sci Rep*. 2017;7: 8940. doi:10.1038/s41598-017-08850-w

~~14.~~13. Sala E, Giakoumi S. No-take marine reserves are the most effective protected areas
in the ocean. *ICES Journal of Marine Science*. 2017; doi:10.1093/icesjms/fsx059

~~15.~~14. Betti F, Bavestrello G, Bo M, Asnaghi V, Chiantore M, Bava S, et al. Over 10 years
of variation in mediterranean reef benthic communities. *Marine Ecology*. 2017;38: e12439.
doi:10.1111/maec.12439

15. Szuwalski CS, Burgess MG, Costello C, Gaines SD. High fishery catches through trophic
cascades in china. *Proc Natl Acad Sci U S A*. 2017;114: 717–721. doi:10.1073/pnas.1612722114

16. Davies TK, Mees CC, Milner-Gulland EJ. Use of a counterfactual approach to evaluate
the effect of area closures on fishing location in a tropical tuna fishery. *PLoS ONE*. 2017;12:

554 e0174758. doi:10.1371/journal.pone.0174758

555 17. Guidetti P, Baiata P, Ballesteros E, Di Franco A, Hereu B, Macpherson E, et al. Large-
556 scale assessment of mediterranean marine protected areas effects on fish assemblages. PLoS
557 ONE. 2014;9: e91841. doi:10.1371/journal.pone.0091841

558 18. Friedlander AM, Golbuu Y, Ballesteros E, Caselle JE, Gouezo M, Olsudong D, et al.
559 Size, age, and habitat determine effectiveness of palau’s marine protected areas. PLoS ONE.
560 2017;12: e0174787. doi:10.1371/journal.pone.0174787

561 19. Rodriguez AG, Fanning LM. Assessing marine protected areas effectiveness: A case study
562 with the tobago cays marine park. OJMS. 2017;07: 379–408. doi:10.4236/ojms.2017.73027

563 20. Moland E, Olsen EM, Knutsen H, Garrigou P, Espeland SH, Kleiven AR, et al.
564 Lobster and cod benefit from small-scale northern marine protected areas: Inference
565 from an empirical before-after control-impact study. Proc Biol Sci. 2013;280: 20122679.
566 doi:10.1098/rspb.2012.2679

567 21. Soykan CU, Lewison RL. Using community-level metrics to monitor the effects of marine
568 protected areas on biodiversity. Conserv Biol. 2015;29: 775–783. doi:10.1111/cobi.12445

569 22. Pomeroy RS, Watson LM, Parks JE, Cid GA. How is your mpa doing? A methodology
570 for evaluating the management effectiveness of marine protected areas. Ocean Coast Manag.
571 2005;48: 485–502. doi:10.1016/j.ocecoaman.2005.05.004

572 23. Pomeroy RS, Parks JE, Watson LM. How is your mpa doing ? A guidebook of natural and
573 social indicators for evaluating marine protected areas management effectiveness [Internet].
574 IUCN; 2004. doi:10.2305/IUCN.CH.2004.PAPS.1.en

575 24. Mascia MB, Fox HE, Glew L, Ahmadi GN, Agrawal A, Barnes M, et al. A novel
576 framework for analyzing conservation impacts: Evaluation, theory, and marine protected

577 areas. *Ann N Y Acad Sci.* 2017;1399: 93–115. doi:10.1111/nyas.13428

578 25. Ostrom E. A general framework for analyzing sustainability of social-ecological systems.
579 *Science.* 2009;325: 419–422. doi:10.1126/science.1172133

580 26. Basurto X, Gelcich S, Ostrom E. The social–ecological system framework as a knowledge
581 classificatory system for benthic small-scale fisheries. *Global Environmental Change.* 2013;23:
582 1366–1380. doi:10.1016/j.gloenvcha.2013.08.001

583 27. Lowndes JSS, Best BD, Scarborough C, Afflerbach JC, Frazier MR, O’Hara CC, et al.
584 Our path to better science in less time using open data science tools. *Nat ecol evol.* 2017;1:
585 0160. doi:10.1038/s41559-017-0160

586 28. Halpern BS, Longo C, Hardy D, McLeod KL, Samhouri JF, Katona SK, et al. An
587 index to assess the health and benefits of the global ocean. *Nature.* 2012;488: 615–620.
588 doi:10.1038/nature11397

589 29. Halpern BS, Frazier M, Afflerbach J, O’Hara C, Katona S, Stewart Lowndes JS, et al.
590 Drivers and implications of change in global ocean health over the past five years. *PLoS*
591 *ONE.* 2017;12: e0178267. doi:10.1371/journal.pone.0178267

592 30. Selig ER, Frazier M, O’Leary JK, Jupiter SD, Halpern BS, Longo C, et al. Measuring
593 indicators of ocean health for an island nation: The ocean health index for fiji. *Ecosystem*
594 *Services.* 2015;16: 403–412. doi:10.1016/j.ecoser.2014.11.007

595 31. Halpern BS, Longo C, Scarborough C, Hardy D, Best BD, Doney SC, et al. Assessing
596 the health of the u.S. west coast with a regional-scale application of the ocean health index.
597 *PLoS ONE.* 2014;9: e98995. doi:10.1371/journal.pone.0098995

598 32. Elfes CT, Longo C, Halpern BS, Hardy D, Scarborough C, Best BD, et al. A regional-scale
599 ocean health index for brazil. *PLoS ONE.* 2014;9: e92589. doi:10.1371/journal.pone.0092589

600 33. Anderson JL, Anderson CM, Chu J, Meredith J, Asche F, Sylvia G, et al. The fishery

- performance indicators: A management tool for triple bottom line outcomes. PLoS ONE. 2015;10: e0122809. doi:10.1371/journal.pone.0122809
34. Dowling N, Wilson J, Rudd M, Babcock E, Caillaux M, Cope J, et al. FishPath: A decision support system for assessing and managing data- and capacity- limited fisheries. In: Quinn II T, Armstrong J, Baker M, Heifetz J, Witherell D, editors. Assessing and managing data-limited fish stocks. Alaska Sea Grant, University of Alaska Fairbanks; 2016. doi:10.4027/amdlfs.2016.03
35. Oyanedel R, Macy Humberstone J, Shattenkirk K, Rodriguez Van-Dyck S, Joye Moyer K, Poon S, et al. A decision support tool for designing turf-reserves. BMS. 2017;93: 155–172. doi:10.5343/bms.2015.1095
36. Vilela T, Reid J. Improving hydropower choices via an online and open access tool. PLoS ONE. 2017;12: e0179393. doi:10.1371/journal.pone.0179393
37. Burgess MG, Clemence M, McDermott GR, Costello C, Gaines SD. Five rules for pragmatic blue growth. Marine Policy. 2016; doi:10.1016/j.marpol.2016.12.005
38. NOM-049-SAG/PESC. NORMA oficial mexicana nom-049-sag/pesc-2014, que determina el procedimiento para establecer zonas de refugio para los recursos pesqueros en aguas de jurisdicción federal de los estados unidos mexicanos. DOF. 2014;
39. LGEEPA. Ley general del equilibrio ecológico y la protección al ambiente. DOF. 2017; Available: http://www.diputados.gob.mx/LeyesBiblio/pdf/148/_240117.pdf
40. [Lester S, McDonald G, Clemence M, Dougherty D, Szuwalski C. Impacts of turfs and marine reserves on fisheries and conservation goals: Theory, empirical evidence, and modeling. BMS. 2017;93: 173–198. doi:10.5343/bms.2015.1083](#)
41. [Chirico AAD, McClanahan TR, Eklöf JS. Community- and government-managed marine protected areas increase fish size, biomass and potential value. PLoS ONE. 2017;12: e0182342. doi:10.1371/journal.pone.0182342](#)

- [42. Rossetto M, Micheli F, Saenz-Arroyo A, Montes JAE, De Leo GA, Rochet M-J. No-take marine reserves can enhance population persistence and support the fishery of abalone. *Can J Fish Aquat Sci.* 2015;72: 1503–1517. doi:10.1139/cjfas-2013-0623](#)
- [43. Woodcock P, O’Leary BC, Kaiser MJ, Pullin AS. Your evidence or mine? Systematic evaluation of reviews of marine protected area effectiveness. *Fish Fish.* 2017;18: 668–681. doi:10.1111/faf.12196](#)
- [44. Carter AB, Davies CR, Emslie MJ, Mapstone BD, Russ GR, Tobin AJ, et al. Reproductive benefits of no-take marine reserves vary with region for an exploited coral reef fish. *Sci Rep.* 2017;7: 9693. doi:10.1038/s41598-017-10180-w](#)
- [45. Villaseñor-Derbez JC, Faro C, Wright M, Martínez J. A guide to evaluate the effectiveness of no-take marine reserves in mexico \[Internet\]. 2017. Available: \[https://www.researchgate.net/publication/317840581/_A/_guide/_to/_evaluate/_the/_effectiveness/_of/_no-take/_marine/_reserves/_in/_Mexico\]\(https://www.researchgate.net/publication/317840581/_A/_guide/_to/_evaluate/_the/_effectiveness/_of/_no-take/_marine/_reserves/_in/_Mexico\)](#)
- ~~41.~~ [46. Ferraro PJ, Pattanayak SK. Money for nothing? A call for empirical evaluation of biodiversity conservation investments. *PLoS Biol.* 2006;4: e105. doi:10.1371/journal.pbio.0040105](#)
- ~~42.~~ [47. OECD. Prices - inflation \(cpi\) - oecd data \[Internet\]. 2017. Available: <https://data.oecd.org/price/inflation-cpi.htm>](#)
- [48. Zeileis A. Econometric computing with hc and hac covariance matrix estimators. *Journal of Statistical Software.* 2004;11: 1–17.](#)
- ~~43.~~ [49. Espinosa-Romero MJ, Rodriguez LF, Weaver AH, Villanueva-Aznar C, Torre J. The changing role of ngos in mexican small-scale fisheries: From environmental conservation to multi-scale governance. *Marine Policy.* 2014;50: 290–299. doi:10.1016/j.marpol.2014.07.005](#)
- ~~44.~~ [50. R Core Team. R: A language and environment for statistical computing \[Internet\]. Vienna, Austria: R Foundation for Statistical Computing; 2017. Available: <https://www>.](#)

651 R-project.org/

652 ~~45.~~ 51. Chang W, Cheng J, Allaire J, Xie Y, McPherson J. Shiny: Web application framework
653 for r [Internet]. 2017. Available: <https://CRAN.R-project.org/package=shiny>

654 ~~46.~~ 52. Chang W, Borges Ribeiro B. Shinydashboard: Create dashboards with 'shiny'
655 [Internet]. 2017. Available: <https://CRAN.R-project.org/package=shinydashboard>

656 ~~47.~~ 53. Allaire J, Cheng J, Xie Y, McPherson J, Chang W, Allen J, et al. Rmarkdown:
657 Dynamic documents for r [Internet]. 2017. Available: [https://CRAN.R-project.org/package=](https://CRAN.R-project.org/package=rmarkdown)
658 rmarkdown

659 ~~48.~~ 54. Xie Y. Knitr: A general-purpose package for dynamic report generation in r [Internet].
660 2017. Available: <http://yihui.name/knitr/>

661 ~~49.~~ 55. McCay B. Territorial use rights in fisheries of the northern pacific coast of mexico.
662 BMS. 2017;93: 69–81. doi:10.5343/bms.2015.1091

663 ~~50.~~ 56. McCay BJ, Micheli F, Ponce-Díaz G, Murray G, Shester G, Ramirez-Sanchez S, et
664 al. Cooperatives, concessions, and co-management on the pacific coast of mexico. Marine
665 Policy. 2014;44: 49–59. doi:10.1016/j.marpol.2013.08.001

666 57. Tennekes M. Tmap: Thematic maps [Internet]. 2017. Available: [https://CRAN.](https://CRAN.R-project.org/package=tmap)
667 R-project.org/package=tmap

668 ~~51.~~ 58. Afflerbach JC, Lester SE, Dougherty DT, Poon SE. A global survey of -reserves, terri-
669 torial use rights for fisheries coupled with marine reserves. Global Ecology and Conservation.
670 2014;2: 97–106. doi:10.1016/j.gecco.2014.08.001

671 ~~52.~~ ~~Lester S, McDonald G, Clemence M, Dougherty D, Szuwalski C. Impacts of turfs~~
672 ~~and marine reserves on fisheries and conservation goals: Theory, empirical evidence, and~~
673 ~~modeling. BMS. 2017;93: 173–198. doi:~~

674 ~~53.~~ ~~Rossetto M, Micheli F, Saenz-Arroyo A, Montes JAE, De Leo GA, Rochet M-J. No-take~~

~~marine reserves can enhance population persistence and support the fishery of abalone. Can
J Fish Aquat Sci. 2015;72: 1503–1517. doi:~~

~~54.~~ 59. Suman CS, Saenz-Arroyo A, Dawson C, Luna MC. Manual de instruccion de reef
check california: Guia de instruccion para el monitoreo del bosque de sargazo en la peninsula
de baja california. Pacific Palisades, CA, USA: Reef Check Foundation; 2010.

~~55.~~ 60. Gutiérrez NL, Hilborn R, Defeo O. Leadership, social capital and incentives promote
successful fisheries. Nature. 2011;470: 386–389. doi:10.1038/nature09689

~~56.~~ 61. Finkbeiner EM, Basurto X. Re-defining co-management to facilitate small-scale
fisheries reform: An illustration from northwest mexico. Marine Policy. 2015;51: 433–441.
doi:10.1016/j.marpol.2014.10.010

62. Ball IR, Possingham HP, Watts ME. Marxan and relatives: Software for spatial
conservation prioritization. Spatial conservation prioritization quantitative methods &
computational tools. United Kingdom: Oxford University Press; 2009. pp. 185–195.
Available: <https://espace.library.uq.edu.au/view/UQ:200259>

63. Padleton L, Aghmadian G, Browman H, Thurstand R, Kaplan D, Bartolino V. Debating
the effectiveness of marine protected areas. ICES Journal of Marine Science. 2017;
doi:10.1093/icesjms/fsx154

~~57.~~ 64. Lundquist CJ, Granek EF. Strategies for successful marine conservation: Inte-
grating socioeconomic, political, and scientific factors. Conserv Biol. 2005;19: 1771–1778.
doi:10.1111/j.1523-1739.2005.00279.x

Supporting information

**S1 Appendix. Survey to collect governance information from fishing communi-
ties.** English version

698 **S2 Appendix. Survey to collect governance information from fishing communi-**
699 **ties. Spanish version**

700 **S3 Table. Assigned values and reasoning of socioeconomic and governance indi-**
701 **cators used to color-code the scorecard in MAREA**