

Regulatory network of gene expression in the drought and cold stress responses

Kazuo Shinozaki^{*}, Kazuko Yamaguchi-Shinozaki[†] and Motoaki Seki[‡]

Molecular and genomic studies have shown that several genes with various functions are induced by drought and cold stresses, and that various transcription factors are involved in the regulation of stress-inducible genes. The products of stress-inducible genes function not only in stress tolerance but also in stress response. Genetic studies have identified many factors that modify the regulation of stress responses. Recent progress has been made in analyzing the complex cascades of gene expression in drought and cold stress responses, especially in identifying specificity and crosstalk in stress signaling.

Addresses

^{*}[‡]Laboratory of Plant Molecular Biology, RIKEN Tsukuba Institute, 3-1-1 Koyadai, Tsukuba 305-0074, Japan

[‡]Plant Mutation Exploration Team, Plant Functional Genomics Research Group, RIKEN Genomic Sciences Center, 1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama, Kanagawa 230-0045, Japan

[†]Biological Resources Division, Japan International Research Center for Agricultural Sciences, Ministry of Agriculture, Forestry, and Fisheries, 2-1 Ohwashi, Tsukuba, Ibaraki 305-0851, Japan

^{*}CREST, Japan Science and Technology Corporation (JST), Japan

*e-mail: shinozaki@rtc.riken.go.jp

Correspondence: Kazuo Shinozaki

NCED 9-*cis*-epoxycarotenoid dioxygenase

RD RESPONSIVE TO DEHYDRATION

SnRK SNF1-related protein kinase

Introduction

Environmental stresses, such as drought, high salinity and low temperature, have adverse effects on plant growth and seed production. Plants respond and adapt to these stresses through various biochemical and physiological processes, thereby acquiring stress tolerance. Many genes respond to drought, salt and/or cold stress at the transcriptional level, and the products of these genes function in the stress response and tolerance [1–3]. Transcriptome analyses using microarray technology [4–6] have identified several genes that are induced by abiotic stresses, and these genes have been classified into two major groups [1–3]. One group encodes products that directly protect plant cells against stresses, whereas the products of the other group regulate gene expression and signal transduction in abiotic stress responses. Molecular and genomic analyses have shown that several different transcriptional regulatory systems are involved in stress-responsive gene induction. Several different sets of *cis*- and *trans*-acting factors are known to be involved in stress-responsive transcription. Some of them are controlled by abscisic acid (ABA) but others are not, indicating the involvement of both ABA-dependent and -independent regulatory systems for stress-responsive gene expression [1–3,7]. Many genes are induced by both drought and cold stress, suggesting the existence of crosstalk between the drought and cold-stress signaling pathways. The use of mutants isolated from transgenic *Arabidopsis* that contain promoter::luciferase (LUC) constructs provides a powerful method for the analysis of stress signaling pathways [7,8]. In this short review, we highlight recent progress in understanding the regulation of gene expression in response to drought and/or cold stress, and in revealing complex gene networks for specificity and crosstalk in abiotic-stress-responsive gene expression.

Transcriptome analysis of stress-inducible gene expression using microarray technology

Microarray technology using cDNAs or oligonucleotides has become a powerful and useful tool for analyzing the gene expression profiles of plants that are exposed to abiotic stresses, such as drought, cold and high salinity, or to ABA treatment [4,5,9,10^{••},11[•],12,13[•]–15[•]]. Potential *cis*-acting DNA elements have been analyzed by comparing their expression profiles with those of the promoter

Current Opinion in Plant Biology 2003, **6**:410–417

This review comes from a themed issue on
Cell signalling and gene regulation
Edited by Kazuo Shinozaki and Elizabeth Dennis

1369-5266/\$ – see front matter
© 2003 Elsevier Ltd. All rights reserved.

DOI 10.1016/S1369-5266(03)00092-X

Abbreviations

ABA	abscisic acid
aba	ABA-deficient
ABF	ABRE-binding factor
abi	ABA-insensitive
ABRE	ABA-responsive element
AREB	ABRE-binding protein
bHLH	basic helix–loop–helix
bZIP	basic-domain leucine zipper
CBF	C-REPEAT-BINDING FACTOR
CRT	C-repeat
DRE	dehydration-responsive element
DREB	DRE-binding protein
ERD1	EARLY RESPONSIVE TO DEHYDRATION1
ERF	ethylene-responsive element binding factor
fry1	fry1
hos	high expression of osmotically responsive genes
ICE1	INDUCER OF CBF EXPRESSION1
los	low expression of osmotically responsive genes
LT1	LOW-TEMPERATURE-INDUCED
LUC	luciferase

sequences of stress-inducible genes [9,10^{••},11[•],13[•]]. Microarray technology is also useful in identifying target genes for stress-related transcription factors, opening a way to analyze gene networks in abiotic stress responses. There are two major forms of microarray technology: the cDNA-spotted glass microarray and the oligonucleotides-based microarray, which is used by GeneChip [4–6]. Both kinds of microarray have been used to analyze gene-expression profiles under conditions of drought, cold and high salinity stress. First, cDNA microarrays were used to identify drought-, cold- and high-salinity-inducible genes [9,10^{••},11[•],12]. Recently, 299 drought-inducible genes, 54 cold-inducible genes, 213 high-salinity-stress-inducible genes and 245 ABA-inducible genes were identified using a cDNA microarray containing around 7000 independent *Arabidopsis* full-length cDNA groups [10^{••},11[•]]. These cDNAs were collected from stress-treated plants as well as normally grown plants. Thousands of stress-inducible genes were identified using a GeneChip array containing oligonucleotides for about 8000 independent *Arabidopsis* genes. The identification of these stress-inducible genes was based on the annotation of these genes on partial genomics sequences [15[•]]. Many differences in the lists of stress-inducible genes were found between cDNA microarray and GeneChip analyses. This is mainly due to the different sets of genes that are arrayed in the two systems (only 1919 genes are represented in both systems) and different conditions of plant growth and stress treatments (K Maruyama *et al.*, unpublished data).

More than half of the drought-inducible genes are also induced by high salinity and/or ABA treatments, indicating the existence of significant crosstalk among the drought, high-salinity and ABA responses [10^{••},11[•]]. By contrast, only 10% of the drought-inducible genes were also induced by cold stress. Many transcription-factor genes were found among the stress-inducible genes, suggesting that various transcriptional regulatory mechanisms function in the drought-, cold- or high-salinity-stress signal transduction pathways [10^{••},11[•],13[•]–15[•]]. These stress-inducible transcription factors include members of the DRE-binding protein (DREB) family, the ethylene-responsive element binding factor (ERF) family, the zinc-finger family, the WRKY family, the MYB family, the basic helix–loop–helix (bHLH) family, the basic-domain leucine zipper (bZIP) family, the NAC family, and the homeodomain transcription factor family. These transcription factors could regulate various stress-inducible genes cooperatively or separately, and may constitute gene networks. Functional analysis of these stress-inducible transcription factors should provide more information on the complex regulatory gene networks that are involved in responses to drought, cold and high-salinity stresses.

Transcriptome analysis has also revealed many genes that respond to rehydration after drought stress, indicating

their involvement in the process of recovery from abiotic stress [16^{••}]. The products of these genes are thought to function not only in recovery from stress but also in cell growth and elongation. The expression and function of the rehydration-inducible proline dehydrogenase gene (*ProDH*) has been precisely analyzed. This gene is involved in the degradation of the proline that accumulates during dehydration [17]. Promoter analysis of the *ProDH* gene revealed an important *cis*-acting element, ACTCAT, that is involved in rehydration-inducible gene expression [18[•]]. Many rehydration-inducible gene promoters contain the ACTCAT motif. There are probably many other genes that are induced during the deacclimation process after cold stress. Similar molecular and genomic analyses will help to reveal total plant responses not only during abiotic stress but also during recovery from stress.


Systems that regulate gene expression in response to drought and cold stress: identification of *cis*-acting elements and their DNA-binding proteins in ABA-independent pathways

The *cis*-acting elements of some genes that have a typical stress-inducible expression profile and the transcription factors that affect the expression of these genes have been analyzed precisely [1–3]. Promoter analyses of drought- and/or cold-inducible genes have provided at least four independent regulatory systems for gene expression [2].

The promoter of a drought-, high-salinity- and cold-inducible gene, *RESPONSIVE TO DEHYDRATION29A* (*RD29A*)/*COLD-REGULATED78* (*COR78*)/*LOW-TEMPERATURE-INDUCED78* (*LTI78*), has been found to contain two major *cis*-acting elements, the ABA-responsive element (ABRE) and the dehydration-responsive element (DRE)/C-repeat (CRT), that are involved in stress-inducible gene expression [19]. ABRE and DRE/CRT are *cis*-acting elements that function in ABA-dependent and ABA-independent gene expression in response to abiotic stress, respectively (Figure 1). Transcription factors belonging to the ERF/APETALA2 (AP2) family that bind to DRE/CRT have also been isolated and termed *C-REPEAT-BINDING FACTOR* (*CBF*)/*DREB1* and *DREB2* [20,21]. The conserved DNA-binding motif of *DREB1A/CBF3* and *DREB2* is A/GCCGAC [22]. The *CBF/DREB1* genes are quickly and transiently induced by cold stress, and their products activate the expression of target stress-inducible genes. The *DREB2* genes are induced by dehydration, leading to the expression of various genes that are involved in drought-stress tolerance [21]. The inducible expression of *CBF/DREB1* also activates target genes that are involved in acquired stress tolerance.

Overexpression of *CBF/DREB1* in transgenics increased tolerance of freezing, drought and high salt concentrations

Figure 1



Regulatory network of gene expression in response to drought and cold stresses: specificity and crosstalk of gene networks. *Cis*-acting elements that are involved in stress-responsive transcription are shown in boxes. Transcription factors that control stress-inducible gene expression are shown in circles or ovals. Small shaded circles indicate the modification of transcription factors in response to stress signals for their activation, such as phosphorylation. The upper part of the figure shows transcription cascades that are involved in rapid and emergency responses to drought and cold stress, such as those involving ICE, DREB2 or NCED. Lower parts of the figure show transcription cascades that are involved in slow and adaptive processes in stress responses, such as those involving AREB/ABF, MYB, MYC and CBF/DREB1. A two-component histidine kinase 1 (ATHK1) is thought to function as an osmosensor, and phospholipase C (PLC) is thought to function upstream of the DREB2 system [2]. HOS1 functions as a negative regulator of ICE and CBF/DREB1 [49]. FRY1 functions as a negative regulator of drought, cold and ABA responses [46]. SnRK2- and SnRK3-like protein kinases are involved in ABA signaling [40–42]. ABI1/ABI2 functions as a negative regulator for ABA signaling. A dashed arrow indicates that CBF4/DREB1D are involved in gene expression in response to drought stress [25]. The open double-headed arrow suggests crosstalk between DREB2 and AREB/ABF that is based on DRE/CRT's acting as a coupling element for ABRE [55]. ERA1, ENHANCED RESPONSE TO ABA1; MYBR, MYB recognition site; MYCR, MYC recognition site.

[21,23,24], suggesting that the CBF/DREB1 proteins function without modification in the development of cold-stress tolerance [21]. Many *CBF/DREB1* target genes have been identified using both cDNA and GeneChip microarrays [9,14]; K Maruyama *et al.*, unpublished data). Most of these target genes contain DRE- or DRE-related CCGAC core motif sequences in their promoter regions. By contrast, overexpression of *DREB2* in transgenics does not improve stress tolerance, suggesting that DREB2 proteins require posttranslational activation [21]. However, one of the *CBF/DREB1* genes, *CBF4/DREB1D*, is induced by osmotic stress, suggesting the existence of crosstalk between the CBF/DREB1 and the DREB2 pathways [22,25]. The DREB2 protein is expressed under normal growth conditions and is activated in the early stage of the osmotic stress response through posttranslational modification. The dehydration-inducible CBF/DREB1 may function in next stage of the osmotic stress response. Rice homologs for *CBF/DREB1* and *DREB2*, four *OsDREB1s* and one *OsDREB2* have been identified in the rice genome sequence and function in stress-inducible gene expression [26]. Overexpression of *OsDREB1A* in *Arabidopsis* revealed that this gene has a

similar function to that of its *Arabidopsis* homolog in stress-responsive gene expression and stress tolerance. This indicates that similar transcription factors function in dicotyledons and monocotyledons.

Recently, the *INDUCER OF CBF EXPRESSION1 (ICE1)* gene was identified through the map-based cloning of the *Arabidopsis ice1* mutation [27], which affected the expression of the CBF3/DREB1A promoter::LUC transgene [28]. *ICE* encodes a MYC-type bHLH transcription factor that regulates the expression of *CBF3/DREB1A* but not of other *CBF/DREB1* genes. Overexpression of *ICE1* in transgenics resulted in improved freezing tolerance, supporting an important role for *ICE1* in the cold-stress response (Figure 1). Molecular analysis of the *DREB1C/CBF2* promoter has identified a *cis*-acting element that is involved in cold-inducible gene expression and its DNA-binding protein. The DNA-binding protein has been cloned and shown to be a MYC-type bHLH transcription factor that is different from *ICE1* (Y Imura *et al.*, unpublished data). These results suggest the redundant involvement of MYC-type bHLH transcription factors in the upregulation of *CBF/DREB1* genes. A cold

signal is necessary for the activation of the ICE proteins but the mechanism of this signal remains to be solved.

Several drought-inducible genes do not respond to either cold or ABA treatment, suggesting the existence of an ABA-independent pathway in the dehydration stress response. These genes include *EARLY RESPONSIVE TO DEHYDRATION1* (*ERD1*), which encodes a Clp protease regulatory subunit, ClpD [29]. The *ERD1* gene is not only induced by dehydration but also upregulated during natural senescence and dark-induced senescence [30[•]]. Promoter analysis of the *ERD1* gene in transgenic plant indicates that the *ERD1* promoter contains *cis*-acting element(s) that are involved not only in ABA-independent stress-responsive gene expression but also in senescence-activated gene expression. Analysis of the *ERD1* promoter also identified two novel *cis*-acting elements that are involved in induction by dehydration stress and in dark-induced senescence [30[•]]. Cloning of the DNA-binding proteins for these *cis*-elements will reveal this novel regulatory system (Figure 1).

Cis-acting elements and transcription factors involved in ABA-dependent gene expression

ABA is synthesized *de novo* mainly in response to drought and high-salinity stress but not in response to cold stress. Many stress-inducible genes are regulated by the endogenous ABA that accumulates during drought and high-salinity stress [1,2]. Recently, genes that are involved in ABA biosynthesis have been identified through genetic and genomics analyses [7[•],31]. Several of these genes are induced by drought and high salinity but not by cold stress. This indicates important roles for ABA in drought-stress responses. The genes encoding 9-*cis*-epoxycarotenoid dioxygenase (NCED), a key enzyme of ABA biosynthesis, are strongly induced by drought stress [32–34]. However, regulatory factors that control the drought-inducible expression of the *NCED* gene have not been elucidated (Figure 1).

ABRE is a major *cis*-acting element in ABA-responsive gene expression (Figure 1). Two ABRE motifs are important in the ABA-responsive expression of the *Arabidopsis* gene *RD29B* [35]. The bZIP transcription factors ABRE-binding protein (AREB)/ABRE-binding factor (ABF) can bind to ABRE and activate ABA-dependent gene expression [35,36]. The AREB/ABF proteins have reduced activity in ABA-deficient *aba2* mutants and in ABA-insensitive *abi1* mutants, and enhanced activity in the ABA-hypersensitive *era1* mutant. Hence, activation of the AREB/ABF proteins has been shown to require an ABA-mediated signal [35], which is probably ABA-dependent phosphorylation. Overexpression of *ABF3* or *AREB2/ABF4* caused ABA hypersensitivity, reduced transpiration rate and enhanced drought tolerance of the transgenics [37[•]].

The induction of the drought-inducible gene *RD22* is mediated by ABA, and this gene requires protein biosynthesis for its ABA-dependent expression [2,38]. A MYC transcription factor, *RD22BP1* (*AtMYC2*), and a MYB transcription factor, *AtMYB2*, have been shown to bind *cis*-elements in the *RD22* promoter and cooperatively activate *RD22* ([38]; Figure 1). These MYC and MYB proteins are synthesized after the accumulation of endogenous ABA, indicating that their role is in a late stage of the stress responses. Microarray analysis of *MYC/MYB*-overexpressing transgenics revealed target genes for MYC/MYB, such as the alcohol dehydrogenase gene and ABA- or jasmonic-acid (JA)-inducible genes [39^{••}]. Overexpression of both *AtMYC2* and *AtMYB2* not only caused an ABA-hypersensitive phenotype but also improved the osmotic-stress tolerance of the transgenic plants [39^{••}].

Recently, the type-2 SNF1-related protein kinase (SnRK2) that is activated by ABA (OPEN STOMATA1 [OST1]/SnRK2E) has been shown to function upstream of ABA-responsive expression of *RD22* and *RD29B* and of stomatal closure in an ABA signal transduction pathway [40,41]. The *ost1/srk2e* mutant has a wilted phenotype. The SnRK3-like protein kinase, PKS3 (a SALT OVERLY SENSITIVE2 [SOS2]-like protein kinase), functions in general ABA signaling, and interacts with a SOS3-like calcium-binding protein (SCaBP5) and ABI2, a 2C-type protein phosphatase [42]. These protein kinases may be involved in the activation of transcription factors in ABA-responsive gene expression.

In addition to the transcription factors that bind to *cis*-acting elements of stress-inducible genes, many genes for transcription factors of various types are induced by drought and/or cold stress. As described above, microarray analyses have also identified many stress-inducible transcription-factor genes. These transcription factors probably function in stress-inducible gene expression, although their target genes have not yet been identified. Among them, *Arabidopsis* ATHB6, which contains the homeodomain, functions downstream of ABI1 as a negative regulator in the ABA signal transduction pathway [43[•]]. The soybean transcription factor SCOF-1, a zinc-finger protein, can activate stress-inducible genes and increase freezing tolerance in transgenic plants. SCOF-1 interacts with the GBF-type bZIP protein SGBF-1, which binds to the G-box and ABRE [44].

Genetic analysis of drought and cold-stress signaling pathways

A unique mutant screening system using transgenic *Arabidopsis* plants with a firefly *LUC* reporter gene under the control of the *RD29A* promoter has been developed to screen mutants that have defects in their abiotic-stress signal transduction pathways [45]. Using this system, many *Arabidopsis* mutants have been isolated that have altered induction of stress-responsive genes by drought,

high-salinity, cold and ABA. These mutants exhibited altered expression of the *RD29A::LUC* gene at a constitutive (*cos*), high (*hos*), or low (*los*) level in response to various abiotic-stress or ABA treatments [45]. The mutated genes are thought to function not only upstream of *RD29A* induction in signal transduction pathways but also in posttranscriptional regulation of the activation of the DREB1/CBF, DREB2 and/or AREB/ABF transcription factors.

The occurrence of mutations with differential responses to drought, cold, high-salinity and/or ABA treatments predicts crosstalk among the drought, salinity, cold and ABA signal transduction pathways [31]. The loci of the mutations, such as *fiery1* (*fry1*) [46], *fry2* [47,48], *hos1* [49], *los1* [50], *los2* [51], *los5/aba3* [52], *los6/aba1* [53] and *sad1* [54], have been cloned. The roles of their gene products have been described precisely in recent reviews by Zhu and his colleagues [7,8]. Among them, three gene products are thought to function directly in transcription. The *hos1* mutation enhances the expression of stress-inducible genes and functions as a negative regulator of a cold-specific signal transduction pathway [49]. HOS1 encodes a novel protein that contains a RING-finger motif, which may function in the degradation of CBF/DREB1 and ICE proteins (Figure 1). It is located in the cytoplasm but moves to the nucleus upon cold treatment, and is thought to control the expression of *CBF/DREB1*. LOS2 is involved in cold-responsive gene expression and encodes a bi-functional enolase [51]. This enzyme is thought to function as a transcription factor that controls STZ/ZAT10, a zinc-finger transcriptional repressor, and to repress *RD29A* expression. FRY2/CPL1 also functions as a transcriptional repressor and contains a region that is partially homologous to the catalytic domain of RNA polymerase II carboxy-terminal domain phosphatases, which are found in humans and yeast [47,48]. FRY2 also contains two double-stranded RNA-binding domains, which may function in mRNA processing. The mutant screening system using transgenic *Arabidopsis* plants containing stress-inducible promoter::LUC provides a powerful tool not only for the analysis of signal transduction pathways upstream of gene expression but also for studies of the posttranscriptional regulation of gene expression. Recently, ICE1 has been identified using the *CBF3/DREB1A* promoter::LUC system as described above [28••].

Complexity of stress-responsive gene networks: specificity and crosstalk

Four transcriptional regulatory systems have been reported, two of them are ABA-independent whereas two are ABA-dependent. Crosstalk between these regulatory systems has, however, been suggested by genetic and molecular analyses. Genomic analyses of stress-inducible genes using microarrays have recently revealed crosstalk in stress-responsive gene expression [10••,11•,15•]. Most

drought-inducible genes are also induced by high-salinity stress, and many drought-inducible genes are also induced by ABA. However, only 10% of drought-inducible genes are also cold-inducible. In fact, ABA is involved in osmotic-stress-responsive gene expression but does not play an important role in the low-temperature stress response.

Crosstalk occurs at the interaction between different *cis*-acting elements. Many drought- and cold-inducible genes contain both DRE/CRT and ABRE motifs in their promoters. These *cis*-acting elements are thought to function independently. However, precise analysis of these *cis*-acting elements in *RD29A* gene expression revealed that DRE/CRT functions cooperatively with ABRE as a coupling element in ABA-responsive gene expression in response to drought stress [55••]. This indicates that interactions between different transcription machineries function to provide crosstalk between different stress signaling pathways. Recently, an osmotic-stress inducible *CBF4/DREB1D* gene has been identified [25•]. Genes of the *CBF/DREB1* family are mainly induced by cold stress, but the drought-inducible gene *CBF4* functions to provide crosstalk between DREB2 and CBF/DREB1 regulatory systems. The drought-inducible expression of *CBF4* is controlled by ABA-dependent pathways, suggesting that CBF4 may function in the slow response to drought that relies on the accumulation of ABA. A maize DRE-binding protein, DBF1, has been shown to function as a transcriptional activator of the *rab17* promoter by ABA [56]. This also suggests the existence in some plants of an ABA-dependent pathway for the regulation of stress-inducible genes that involves DRE/CRT. Cold-induced gene expression through DRE/CRT is greatly enhanced by a light signal mediated by phytochrome B, which indicates crosstalk between abiotic-stress and light-signaling pathways [57•].

The *fry1* mutation results in elevated levels of inositol 1,4,5-triphosphate (Ins[1,4,5]P₃) in both control and ABA-treated plants. It also causes the super-induction of ABA- and stress-inducible genes in response to ABA, salt and dehydration, and gives rise to plants that are less tolerant to freezing, drought and salt stress [46]. The *FRY1* gene encodes an inositol polyphosphate 1-phosphatase that is involved in Ins(1,4,5)P₃ signaling. *FRY1* functions as a negative regulator of ABA and stress signaling, and provides genetic evidence that phosphoinositols have an important role in ABA and stress-related signal transduction in plants (Figure 1). Biochemical analysis has shown that phospholipase C functions upstream of *RD29A* expression in osmotic stress signaling [58], which may be negatively regulated by FRY1.

Conclusions and perspectives

Molecular and genomic analyses have revealed many genes that are induced by abiotic stress and their functions in stress-responsive gene expression and stress

tolerance. The many stress-inducible genes include those that encode signaling molecules, such as enzymes involved in phospholipid metabolism and various protein kinases including mitogen-activated protein (MAP) kinases, calcium-dependent protein kinases (CDPKs), receptor-like kinases and histidine kinases [2,7,8]. Transgenic plants and T-DNA or transposon-tagged mutants have been analyzed to reveal the function of stress-responsive loci in *Arabidopsis*. The reverse genetics approach will be even more important in extending our understanding of regulatory factors in stress signaling. Genetic analysis of regulatory factors using the stress-inducible promoter::LUC system is promising, and can be used to reveal not only upstream stress signaling pathways but also posttranscriptional regulatory systems that modify abiotic stress responses. In the future, a combination of molecular, genomic and genetic analyses will be used to elucidate the complex systems that regulate the responses of gene expression to abiotic stresses.

Acknowledgements

This work was supported in part by a grant for Genome Research from RIKEN, by the Special Coordination Fund of the Science and Technology Agency and by a Grant-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan (MECSST) to KS. KS and KY-S are also recipients of a grant from the Program for the Promotion of Basic Research Activities for Innovative Biosciences. The work was also supported in part by a Grant-in-Aid for Scientific Research on Priority Areas (C) 'Genome Science' from MECSST to MS.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Bray E, Bailey-Serres J, Weretilnyk E: **Responses to abiotic stresses**. In *Biochemistry and Molecular Biology of Plants*. Edited by Buchanan B, Gruissem W, Jones R Rockville: American Society of Plant Biologists; 2000:1158-1203.
2. Shinozaki K, Yamaguchi-Shinozaki K: **Molecular responses to dehydration and low temperature: differences and cross-talk between two stress signaling pathways**. *Curr Opin Plant Biol* 2000, **3**:217-223.
3. Thomashow MF: **Plant cold acclimation: freezing tolerance genes and regulatory mechanisms**. *Annu Rev Plant Physiol Plant Mol Biol* 1999, **50**:571-599.
4. Seki M, Narusaka M, Yamaguchi-Shinozaki K, Carninci P, Kawai J, Hayashizaki Y, Shinozaki K: **Arabidopsis encyclopedia using full-length cDNAs and its application**. *Plant Physiol Biochem* 2001, **39**:211-220.
5. Bohnert HJ, Ayoubi P, Borchert C, Bresson RA, Burnap RL, Cushman JC, Cushman MA, Deyholos M, Fischer R, Galbraith DW *et al.*: **A genomics approach towards salt stress tolerance**. *Plant Physiol Biochem* 2001, **39**:295-311.
6. Zhu T, Budworth P, Han B, Brown D, Chang HS, Zou G, Wang X: **Toward elucidating the global expression patterns of developing Arabidopsis: parallel analysis of 8300 genes by a high-density oligonucleotide probe array**. *Plant Physiol Biochem* 2001, **39**:221-242.
7. Xiong L, Schumaker KS, Zhu JK: **Cell signaling during cold, drought, and salt stress**. *Plant Cell* 2002 (Suppl), S165-S183.
This is an excellent review article on signal transduction and gene expression in plants that are exposed to abiotic stress.
8. Zhu JK: **Salt and drought stress signal transduction in plants**. *Annu Rev Plant Biol* 2002, **53**:247-273.
9. Seki M, Narusaka M, Abe H, Kasuga M, Yamaguchi-Shinozaki K, Carninci P, Hayashizaki Y, Shinozaki K: **Monitoring the expression pattern of 1300 Arabidopsis genes under drought and cold stresses using a full-length cDNA microarray**. *Plant Cell* 2001, **13**:61-72.
10. Seki M, Narusaka M, Ishida J, Nanjo T, Fujita M, Oono Y, Kamiya A, Nakajima M, Enju A, Sakurai T *et al.*: **Monitoring the expression profiles of 7000 Arabidopsis genes under drought, cold, and high-salinity stresses using a full-length cDNA microarray**. *Plant J* 2002, **31**:279-292.
The authors used 7000 *Arabidopsis* full-length cDNA microarrays to analyze drought-, cold-, and high-salinity stress-inducible genes. The results indicate the existence of greater crosstalk between drought and high-salinity stress signaling processes than between cold and high-salinity stress signaling processes.
11. Seki M, Ishida J, Narusaka M, Fujita M, Nanjo T, Umezawa T, Kamiya A, Nakajima M, Enju A, Sakurai T *et al.*: **Monitoring the expression pattern of ca. 7000 Arabidopsis genes under ABA treatments using a full-length cDNA microarray**. *Funct Integr Genom* 2002, **2**:282-291.
ABA-inducible genes were analyzed using 7000 *Arabidopsis* full-length cDNA microarrays. The existence of great crosstalk between ABA and osmotic stress signaling processes was indicated.
12. Kawasaki S, Borchert C, Deyholos M, Wang H, Brazille S, Kawai K, Galbraith D, Bohnert H: **Gene expression profiles during the initial phase of salt stress in rice**. *Plant Cell* 2001, **13**:889-905.
13. Chen W, Provart NJ, Glazebrook J, Katagiri F, Chang HS, Eulgem T, Mauch F, Luan S, Zou G, Whitham SA *et al.*: **Expression profile matrix of Arabidopsis transcription factor genes suggests their putative functions in response to environmental stresses**. *Plant Cell* 2002, **14**:559-574.
The expression profiles of 402 *Arabidopsis* transcription factors are reported to respond to various stresses and hormone treatments. This work reveals overlap in the expression of several transcription factors between different stress signaling pathways.
14. Fowler S, Thomashow MF: **Arabidopsis transcriptome profiling indicates that multiple regulatory pathways are activated during cold acclimation in addition to the CBF cold response pathway**. *Plant Cell* 2002, **14**:1675-1690.
The authors used the 8000 GeneChip array to analyze the expression profiles of cold-inducible genes, thereby revealing multiple regulatory pathways in cold-inducible gene expression. The CBF/DREB1 target genes were also analyzed to identify novel stress-inducible genes including transcription factors.
15. Krebs JA, Wu Y, Chang HS, Zhu T, Wang X, Harper J: **Transcriptome changes for Arabidopsis in response to salt, osmotic, and cold stress**. *Plant Physiol* 2002, **130**:2129-2141.
The authors used the 8000 GeneChip array to describe the expression profiles of high-salinity-, osmotic-stress and cold-inducible genes. They discuss crosstalk in the induction of gene expression by different abiotic stresses.
16. Oono Y, Seki M, Nanjo T, Narusaka M, Fujita M, Satoh R, Satou MK, Sakurai T, Ishida J, Akiyama K *et al.*: **Monitoring expression profiles of Arabidopsis gene expression during rehydration process after dehydration using ca. 7000 full-length cDNA microarray**. *Plant J* 2003, **34**:868-887.
This is the first report of a study in which cDNA microarrays were used to examine gene expression profiles during the rehydration and recovery process following drought stress.
17. Nakashima K, Satoh R, Kiyosue T, Yamaguchi-Shinozaki K, Shinozaki K: **A gene encoding proline dehydrogenase is not only induced by proline and hypoosmolarity, but is also developmentally regulated in the reproductive organs of Arabidopsis**. *Plant Physiol* 1998, **118**:1233-1241.
18. Satoh R, Nakashima K, Seki M, Shinozaki K, Yamaguchi-Shinozaki K: **ACTCAT, a novel cis-acting-element for proline- and hypoosmolarity-responsive expression of the ProDH gene encoding proline dehydrogenase in Arabidopsis**. *Plant Physiol* 2002, **130**:709-719.
The authors identify a novel cis-acting element that is involved in hypoosmolarity, proline treatment and rehydration. This element plays an important role in gene expression during recovery from drought stress.
19. Yamaguchi-Shinozaki K, Shinozaki K: **A novel cis-acting element in an Arabidopsis gene is involved in responsiveness to**

- drought, low temperature, or high-salt stress. *Plant Cell* 1994, 6:251-264.
20. Stockinger EJ, Gilmour SJ, Thomashow MF: ***Arabidopsis thaliana* CBF1 encodes an AP2 domain-containing transcription activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit.** *Proc Natl Acad Sci USA* 1997, 94:1035-1040.
 21. Liu X, Kasuga M, Sakuma Y, Abe H, Miura S, Yamaguchi-Shinozaki K, Shinozaki K: **Two transcription factors, DREB1 and DREB2, with an ERF/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low-temperature-responsive gene expression, respectively, in *Arabidopsis*.** *Plant Cell* 1998, 10:1391-1406.
 22. Sakuma Y, Liu Q, Dubouzet JG, Abe H, Shinozaki K, Yamaguchi-Shinozaki K: **DNA-binding specificity of the ERF/AP2 domain of *Arabidopsis* DREBs, transcription factors involved in dehydration- and cold-inducible gene expression.** *Biochem Biophys Res Com* 2002, 290:998-1009.
 23. Jaglo-Ottosen KR, Gilmour SJ, Zarka DG, Schabenberger O, Thomashow MF: ***Arabidopsis* CBF1 overexpression induces *cor* genes and enhances freezing tolerance.** *Science* 1998, 280:104-106.
 24. Kasuga M, Liu Q, Miura S, Yamaguchi-Shinozaki K, Shinozaki K: **Improving plant drought, salt, and freezing tolerance by gene transfer of a single stress-inducible transcription factor.** *Nat Biotechnol* 1999, 17:287-291.
 25. Haerke V, Cook D, Riechmann JL, Ombra P, Thomashow MF, Zang JZ: **Transcription factor CBF4 is a regulator of drought adaptation in *Arabidopsis*.** *Plant Physiol* 2002, 130:639-648.
- Drought-inducible CBF4 functions in drought-stress responsive gene expression, which suggests crosstalk between drought- and cold-inducible transcription.
26. Dubouzet J, Sakuma Y, Kasuga M, Dubouzet E, Miura S, Seki M, Shinozaki K, Yamaguchi-Shinozaki K: ***OsDREB* genes in rice, *Oryza sativa* L., encode transcription factors that function in drought-, high salt- and cold responsive gene expression.** *Plant J* 2003, 33:751-763.
- Rice has homologs of the *Arabidopsis* transcription factors DREB1 and DREB2, indicating that drought- and cold-inducible gene expression has a similar transcription system in monocotyledons and dicotyledons.
27. Gilmour SJ, Zarka DG, Stockinger EJ, Salazar MP, Houghton JM, Thomashow MF: **Low temperature regulation of *Arabidopsis* CBF family of AP2 transcriptional activators as an early step in cold-induced *COR* gene expression.** *Plant J* 1998, 16:433-442.
 28. Chinnusamy V, Ohta M, Kanrar S, Lee B, Hong X, Agarwal A, Zhu JK: **ICE1: a regulator of cold-induced transcriptome and freezing tolerance in *Arabidopsis*.** *Gene Dev* 2003, 17:1043-1054.
- The authors use map-based cloning of a mutant that was isolated using CBF3/DREB1A promoter::LUC transgenic *Arabidopsis* to identify ICE1 as a key transcription factor that regulates the expression of CBF/DREB1 in response to cold stress. ICE1 is an important transcription factor early in the cold-signaling pathway.
29. Nakashima K, Kiyosue T, Shinozaki K, Yamaguchi-Shinozaki K: **A nuclear gene encoding a chloroplast-targeted Clp protease regulatory subunit homolog is not only induced by water stress but also developmentally upregulated during senescence in *Arabidopsis thaliana*.** *Plant J* 1997, 12:851-861.
 30. Simpson S, Nakashima K, Narusaka Y, Seki M, Shinozaki K, Yamaguchi-Shinozaki K: **Two different novel cis-acting elements of *erd1*, a *clpA* homologous *Arabidopsis* gene function in induction by dehydration stress and dark-induced senescence.** *Plant J* 2003, 33:259-270.
- The authors determine novel cis-acting elements that are involved in drought-stress-responsive gene expression and senescence. These cis elements are involved in a novel pathway in drought-responsive gene expression.
31. Seo M, Koshida T: **Complex regulation of ABA biosynthesis.** *Trends Plant Sci* 2002, 7:41-48.
 32. Quin X, Zeevaart JAD: **The 9-cis-epoxycarotenoid cleavage reaction is the key regulatory step of abscisic acid biosynthesis in water-stressed bean.** *Proc Natl Acad Sci USA* 1999, 96:15354-15361.
 33. Thompson AJ, Jackson AC, Symonds RC, Mulholland BJ, Dadwell AR: **Ectopic expression of tomato 9-cis-epoxycarotenoid dioxygenase gene causes overproduction of abscisic acid.** *Plant J* 2000, 23:363-374.
 34. Iuchi S, Kobayashi M, Taji T, Naramoto M, Seki M, Kato T, Tabata S, Kakubari Y, Yamaguchi-Shinozaki K, Shinozaki K: **Regulation of drought tolerance by gene manipulation of 9-cis-epoxycarotenoid dioxygenase, a key enzyme in abscisic acid biosynthesis in *Arabidopsis*.** *Plant J* 2001, 27:325-333.
 35. Uno Y, Furihata T, Abe H, Yoshida R, Shinozaki K, Yamaguchi-Shinozaki K: ***Arabidopsis* basic leucine zipper transcription factors involved in an abscisic acid-dependent signal transduction pathway under drought and high-salinity conditions.** *Proc Natl Acad Sci USA* 2000, 97:11632-11637.
 36. Choi H, Hong JH, Ha J, Kang JY, Kim SY: **ABFs, a family of ABA-responsive element binding factors.** *J Biol Chem* 2000, 275:1723-1730.
 37. Kang JY, Choi H, Im MY, Kim SY: ***Arabidopsis* basic leucine zipper proteins that mediate stress-responsive abscisic acid signaling.** *Plant Cell* 2002, 14:343-357.
- Overexpression of ABP caused ABA hypersensitivity and improved stress tolerance, indicating important roles for bZIP transcription factors in ABA signal transduction pathways.
38. Abe H, Yamaguchi-Shinozaki K, Urao T, Iwasaki T, Hosokawa D, Shinozaki K: **Role of *Arabidopsis* MYC and MYB homologs in drought- and abscisic acid-regulated gene expression.** *Plant Cell* 1997, 9:1859-1868.
 39. Abe H, Urao T, Ito T, Seki M, Shinozaki K, Yamaguchi-Shinozaki K: ***Arabidopsis* AtMYC2 (bHLH) and AtMYB2 (MYB) function as transcriptional activators in abscisic acid signaling.** *Plant Cell* 2003, 15:63-78.
- ABA-inducible MYC and MYB transcription factors function in drought-stress-responsive gene expression. The target genes of these transcription factors are determined and suggest that these transcription factors function in crosstalk between various stress responses.
40. Mustilli AC, Merlot S, Vavasseur A, Fenzi F, Giraudat J: ***Arabidopsis* OST1 protein kinase mediates the regulation of stomatal aperture by abscisic acid and acts upstream of reactive oxygen species production.** *Plant Cell* 2002, 14:3089-3099.
 41. Yoshida R, Hobo T, Ichimura K, Mizoguchi T, Takahashi F, Alonso J, Ecker JR, Shinozaki K: **ABA-activated SnRK2 protein kinase is required for dehydration stress signaling in *Arabidopsis*.** *Plant Cell Physiol* 2002, 43:1473-1483.
 42. Guo Y, Xiong L, Song CP, Gong D, Halfter U, Zhu JK: **A calcium sensor and its interacting protein kinase are global regulators of abscisic acid signaling in *Arabidopsis*.** *Dev Cell* 2002, 3:233-244.
 43. Himmelbach A, Hoffmann T, Leube M, Hohener B, Grill E: **Homeodomain protein ATHB6 is a target of the protein phosphatase ABI1 and regulates hormone responses in *Arabidopsis*.** *EMBO J* 2002, 21:3029-3038.
- A homeodomain transcription factor ATHB6 was shown to interact with ABI1 protein phosphatase and to function as a negative regulator in ABA signal transduction.
44. Kim JC, Lee SH, Cheong YH, Yoo CM, Lee SI, Chun HJ, Yun DJ, Hong DJ, Lee SY, Lim CO, Cho MJ: **A novel cold-inducible zinc finger protein from soybean, SCOF-1, enhances cold tolerance in transgenic plants.** *Plant J* 2001, 25:247-259.
 45. Ishitani M, Xiong L, Stevenson B, Zhu JK: **Genetic analysis of osmotic and cold stress signal transduction in *Arabidopsis*: interactions and convergence of abscisic acid-dependent and abscisic acid-independent pathways.** *Plant Cell* 1997, 9:1935-1949.
 46. Xiong L, Lee BH, Ishitani M, Lee H, Zhang C, Zhu JK: **FIERY1 encoding an inositol polyphosphate 1-phosphatase is a negative regulator of abscisic acid and stress signaling in *Arabidopsis*.** *Genes Dev* 2001, 15:1971-1984.
 47. Xiong L, Lee H, Ishitani M, Tanaka Y, Stevenson B, Koiwa H, Bresson RA, Hasegawa PM, Zhu JK: **Repression of stress-responsive genes by FIERY2, a novel transcriptional regulator in *Arabidopsis*.** *Proc Natl Acad Sci USA* 2002, 99:10899-10904.

48. Koiwa H, Barb AW, Xiong L, Li F, McCully MG, Lee B, Sokolchik I, Zhu J, Gong Z, Reddy M *et al.*: **C-terminal domain phosphatase-like family members (AtCPLs) differentially regulate *Arabidopsis thaliana* abiotic stress signaling, growth, and development.** *Proc Natl Acad Sci USA* 2002, **99**:10893-10898.
 49. Lee H, Xiong L, Gong Z, Ishitani M, Stevenson B, Zhu JK: **The *Arabidopsis HOS1* gene negatively regulates cold signal transduction and encodes a RING finger protein that displays cold-regulated nucleocytoplasmic partitioning.** *Genes Dev* 2001, **15**:912-924.
 50. Guo Y, Xiong L, Ishitani M, Zhu JK: **An *Arabidopsis* mutation in translation elongation factor 2 causes superinduction of CBF/DREB1 transcription factor genes but blocks the induction of their downstream targets under low temperatures.** *Proc Natl Acad Sci USA* 2002, **99**:7786-7791.
 51. Lee H, Guo Y, Ohta M, Xiong L, Stevenson B, Zhu JK: ***LOS2*, a genetic locus acquired for cold-responsive gene transcription encodes a bi-functional enolase.** *EMBO J* 2002, **21**:2692-2702.
 52. Xiong L, Ishitani M, Lee H, Zhu JK: **The *Arabidopsis LOS5/ABA3* locus encodes a molybdenum cofactor sulfurase and modulates cold stress- and osmotic stress-responsive gene expression.** *Plant Cell* 2001, **13**:2063-2083.
 53. Xiong L, Lee H, Ishitani M, Zhu JK: **Regulation of osmotic stress-responsive gene expression by the *LOS6/ABA1* locus in *Arabidopsis*.** *J Biol Chem* 2002, **277**:8588-8596.
 54. Xiong L, Gong Z, Rock CD, Subramanian S, Guo Y, Xu W, Galbraith D, Zhu JK: **Modulation of abscisic acid signal transduction and biosynthesis by an Sm-like protein in *Arabidopsis*.** *Dev Cell* 2001, **1**:771-781.
 55. Narusaka Y, Nakashima K, Shinwari ZK, Sakuma Y, Furihata T, Abe H, Narusaka M, Shinozaki K, Yamaguchi-Shinozaki K: **Interaction between two cis-acting elements, ABRE and DRE, in ABA-dependent expression of *Arabidopsis rd29A* gene in response to dehydration and high-salinity stresses.** *Plant J* 2003, **34**:137-148.
- The authors discuss crosstalk between DRE/CRT and ABRE because DRE functions as a coupling element for ABRE.
56. Kizis D, Pages M: **Maize DRE-binding proteins DBF1 and DBF2 are involved in *rab17* regulation through the drought-responsive element in an ABA-dependent pathway.** *Plant J* 2002, **30**:679-689.
 57. Kim HJ, Kim YK, Park JY, Kim J: **Light signaling mediated by phytochrome plays an important role in cold-induced gene expression through the C-repeat/dehydration responsive element (C/DRE) in *Arabidopsis thaliana*.** *Plant J* 2002, **29**:693-704.
- Cold-induced gene expression through C/DRE is greatly enhanced by light signaling, in which phytochrome B functions as a photoreceptor. Crosstalk between light and low-temperature-stress signaling pathways occurs on the C/DRE cis-acting element.
58. Takahashi S, Katagiri T, Hirayama T, Yamaguchi Shinozaki K, Shinozaki K: **Hyperosmotic stress induced a rapid and transient increase in inositol 1,4,5-triphosphate independent of abscisic acid in *Arabidopsis* cell culture.** *Plant Cell Physiol* 2001, **42**:214-222.