


C-oxidation BDE Energy Report for: 085_citalopram-out

This report covers the results for bond dissociation enthalpies (BDE) and solvent accessible surface area (SASA) calculations performed for 085_citalopram-out. Oxidation propensity is established using C-H BDE. The lower the C-H BDE values the higher the propensity for C-oxidation. Details for the density functional theory (DFT) calculations and overall workflow are explained at the end of this document.


BDE and SASA



Atom	BDE (kcal/mol)	Propensity	SASA (Å ²)
C2	114.14	Low	5.32
C3	111.92	Low	1.37
C4	113.87	Low	0.02
C9	80.22	High	0.00
C12	97.61	Low	0.00
C14	110.39	Low	0.00
C15	112.58	Low	0.00
C16	114.73	Low	0.00
C17	114.57	Low	0.00
C19	97.42	Low	3.98
C20	91.18	Moderate	0.00
C22	92.67	Moderate	3.65
C23	92.9	Moderate	3.41

Missing Sites:

None

Risk Scale:**Calculation Details**

Conformational search calculations were performed only for the base ground state molecule. The lowest energy conformer was selected to generate radicals and run optimization DFT calculations. DFT calculations were performed using Gaussian with B3LYP level of theory and 6-31G(d,p) basis set. The BDE protocol was adapted from: *Lienard, P., Gavartin, J., Boccardi, G., & Meunier, M. (2015). Predicting drug substances autoxidation. Pharmaceutical research, 32, 300-310.*