


C-oxidation BDE Energy Report for: 447_pranidipine-out

This report covers the results for bond dissociation enthalpies (BDE) and solvent accessible surface area (SASA) calculations performed for 447_pranidipine-out. Oxidation propensity is established using C-H BDE. The lower the C-H BDE values the higher the propensity for C-oxidation. Details for the density functional theory (DFT) calculations and overall workflow are explained at the end of this document.

BDE and SASA



Atom	BDE (kcal/mol)	Propensity	SASA (Å ²)
C1	82.02	High	7.92
C5	114.03	Low	1.78
C6	112.06	Low	9.79
C12	112.49	Low	29.63
C14	88.33	Moderate	26.25
C17	87.97	High	27.06
C20	114.91	Low	19.02
C22	73.83	High	22.92
C23	98.03	Low	28.76
C26	107.76	Low	8.88
C27	101.36	Low	9.90
C29	112.39	Low	21.40
C30	111.17	Low	12.31
C31	112.09	Low	34.62

C32	112.0	Low	31.93
C33	112.58	Low	35.89

Missing Sites:

None

Risk Scale:



Calculation Details

Conformational search calculations were performed only for the base ground state molecule. The lowest energy conformer was selected to generate radicals and run optimization DFT calculations. DFT calculations were performed using Gaussian with B3LYP level of theory and 6-31G(d,p) basis set. The BDE protocol was adapted from: *Lienard, P., Gavartin, J., Boccardi, G., & Meunier, M. (2015). Predicting drug substances autoxidation. Pharmaceutical research, 32, 300-310.*