


C-oxidation BDE Energy Report for: 167_indinavir-out

This report covers the results for bond dissociation enthalpies (BDE) and solvent accessible surface area (SASA) calculations performed for 167_indinavir-out. Oxidation propensity is established using C-H BDE. The lower the C-H BDE values the higher the propensity for C-oxidation. Details for the density functional theory (DFT) calculations and overall workflow are explained at the end of this document.

BDE and SASA



Atom	BDE (kcal/mol)	Propensity	SASA (Å ²)
C2	78.25	High	0.03
C4	113.23	Low	6.90
C5	90.92	Moderate	0.11
C7	85.0	High	1.70
C8	112.29	Low	12.04

C9	112.21	Low	10.59
C12	112.19	Low	13.46
C13	89.57	Moderate	0.00
C15	86.61	High	0.88
C16	98.43	Low	0.05
C18	88.62	Moderate	0.00
C19	111.67	Low	7.33
C20	112.82	Low	7.29
C21	90.12	Moderate	0.03
C23	111.87	Low	10.96
C24	111.86	Low	11.99
C26	112.27	Low	12.38
C27	87.38	High	0.00
C28	89.62	Moderate	0.26
C29	91.76	Moderate	0.18
C31	90.97	Moderate	0.29
C35	80.75	High	0.77
C38	102.9	Low	3.98
C39	102.95	Low	3.82
C40	102.79	Low	4.19
C41	112.5	Low	8.05
C42	106.04	Low	10.36
C43	112.43	Low	13.98
C45	107.06	Low	17.74

Missing Sites:

None

Risk Scale:



Calculation Details

Conformational search calculations were performed only for the base ground state molecule. The lowest energy conformer was selected to generate radicals and run optimization DFT calculations. DFT

calculations were performed using Gaussian with B3LYP level of theory and 6-31G(d,p) basis set. The BDE protocol was adapted from: *Lienard, P., Gavartin, J., Boccardi, G., & Meunier, M. (2015). Predicting drug substances autoxidation. Pharmaceutical research, 32, 300-310.*