


C-oxidation BDE Energy Report for: iloperidone


This report covers the results for BDE calculations performed for: iloperidone. Oxidation propensity is established using C-H Bond Dissociation Enthalpies (BDE). The lower the C-H BDE values the higher the propensity for C-oxidation. Details for the DFT calculations and overall workflow are explained at the end of this document

Bond Dissociation Energies (kcal/mol)



Atom	BDE (kcal/mol)	Propensity
C3	112.99	Low
C4	115.68	Low
C6	117.79	Low
C11	85.15	High
C12	100.0	Low
C13	90.81	Moderate
C15	91.34	Moderate
C16	99.66	Low
C17	91.02	Moderate
C18	98.3	Low
C19	94.49	Low
C22	111.93	Low
C23	111.41	Low
C25	113.89	Low
C29	97.24	Low
C31	95.21	Low

Risk Scale:



Calculation details and output files

Conformational search calculations were performed only for the base ground state molecule. The lowest energy conformer was selected to generate radicals and run optimization DFT calculations. DFT calculations were performed using Gaussian with B3LYP level of theory and 6-31G(d,p) basis set. The BDE protocol was adapted from: *Lienard, P., Gavartin, J., Boccardi, G., & Meunier, M. (2015). Predicting drug substances autoxidation. Pharmaceutical research, 32, 300-310.*