


# C-oxidation BDE Energy Report for: 219\_onapristone-out

This report covers the results for BDE calculations performed for: 219\_onapristone-out. Oxidation propensity is established using C-H Bond Dissociation Enthalpies (BDE). The lower the C-H BDE values the higher the propensity for C-oxidation. Details for the DFT calculations and overall workflow are explained at the end of this document


## Bond Dissociation Energies (kcal/mol)



Atom	BDE (kcal/mol)	Propensity
C1	90.11	Moderate
C3	90.11	Moderate
C5	110.28	Low
C6	111.27	Low
C8	110.07	Low
C9	110.24	Low
C10	72.03	High
C12	66.56	High
C13	82.51	High
C15	94.81	Low
C18	83.34	High
C19	98.37	Low
C20	81.89	High
C21	88.04	Moderate
C23	111.45	Low
C26	94.76	Low

C27	95.55	Low
C28	100.35	Low
C29	96.37	Low
C31	96.4	Low
C32	94.44	Low

Risk Scale:



## Calculation details and output files

Conformational search calculations were performed only for the base ground state molecule. The lowest energy conformer was selected to generate radicals and run optimization DFT calculations. DFT calculations were performed using Gaussian with B3LYP level of theory and 6-31G(d,p) basis set. The BDE protocol was adapted from: *Lienard, P., Gavartin, J., Boccardi, G., & Meunier, M. (2015). Predicting drug substances autoxidation. Pharmaceutical research, 32, 300-310.*