


# C-oxidation BDE Energy Report for: 018\_CORTISOL-out

This report covers the results for bond dissociation enthalpies (BDE) and solvent accessible surface area (SASA) calculations performed for 018\_CORTISOL-out. Oxidation propensity is established using C-H BDE. The lower the C-H BDE values the higher the propensity for C-oxidation. Details for the density functional theory (DFT) calculations and overall workflow are explained at the end of this document.

## BDE and SASA



Atom	BDE (kcal/mol)	Propensity	SASA (Å <sup>2</sup> )
C1	94.26	Low	0.00
C2	91.11	Moderate	0.41
C4	95.22	Low	3.83
C5	90.15	Moderate	1.82
C6	96.69	Low	5.98
C8	97.96	Low	6.03
C9	100.94	Low	17.56
C10	96.24	Low	45.58
C12	88.7	Moderate	8.67
C13	80.78	High	15.18
C17	98.22	Low	16.48
C18	100.74	Low	16.85
C20	75.06	High	19.93
C22	111.55	Low	9.65
C23	89.18	Moderate	27.14

**Missing Sites:**

None

**Risk Scale:****Calculation Details**

Conformational search calculations were performed only for the base ground state molecule. The lowest energy conformer was selected to generate radicals and run optimization DFT calculations. DFT calculations were performed using Gaussian with B3LYP level of theory and 6-31G(d,p) basis set. The BDE protocol was adapted from: *Lienard, P., Gavartin, J., Boccardi, G., & Meunier, M. (2015). Predicting drug substances autoxidation. Pharmaceutical research, 32, 300-310.*