


C-oxidation BDE Energy Report for: carvedilol

This report covers the results for bond dissociation enthalpies (BDE) and solvent accessible surface area (SASA) calculations performed for carvedilol. Oxidation propensity is established using C-H BDE. The lower the C-H BDE values the higher the propensity for C-oxidation. Details for the density functional theory (DFT) calculations and overall workflow are explained at the end of this document.

BDE and SASA



Atom	BDE (kcal/mol)	Propensity	SASA (Å ²)
C6	113.0	Low	9.85
C8	114.37	Low	14.92
C9	112.36	Low	8.52
C11	113.19	Low	15.04
C12	112.75	Low	13.38
C13	112.13	Low	14.73
C14	95.37	Low	0.77
C15	112.39	Low	15.30
C16	88.98	Moderate	0.00
C17	92.9	Moderate	0.39
C20	97.15	Low	0.68
C21	95.12	Low	1.29
C25	112.79	Low	8.36
C26	111.61	Low	9.60
C28	113.0	Low	14.13
C29	112.42	Low	14.71
C30	97.71	Low	8.84

Missing Sites:

None

Risk Scale:



Calculation Details

Conformational search calculations were performed only for the base ground state molecule. The lowest energy conformer was selected to generate radicals and run optimization DFT calculations. DFT calculations were performed using Gaussian with B3LYP level of theory and 6-31G(d,p) basis set. The BDE protocol was adapted from: *Lienard, P., Gavartin, J., Boccardi, G., & Meunier, M. (2015). Predicting drug substances autoxidation. Pharmaceutical research, 32, 300-310.*