

MATH 135: Introduction to the Theory of Sets

Jad Damaj

Fall 2022

Contents

1	Introduction	4
1.1	August 25	4
1.1.1	Introduction	4
1.1.2	Basics	4
2	Axioms and Operations	6
2.1	August 30	6
2.1.1	Zermelo Fraenkel Axioms of Set Theory	6
3	Relations and Functions	8
3.1	September 1	8
3.1.1	Relations and Functions	8
3.2	September 6	9
3.2.1	Functions and Relations	9
3.2.2	Infinite Cartesian Products	9
4	Naturals, Rationals, Reals	11
4.1	September 8	11
4.1.1	Natural Numbers	11
4.2	September 13	12
4.2.1	Operations on the Natural Numbers	12
4.2.2	Integers	13
4.2.3	Rationals	13
4.3	September 15	13
4.3.1	Reals (Dedekind Cuts)	13
5	Cardinal Numbers and the Axiom of Choice	14
5.1	September 15	14
5.1.1	Cardinality	14
5.2	September 20	15
5.2.1	Cardinality	15
5.3	September 22	16
5.3.1	Cardinals	16
5.4	September 27	17
5.4.1	Schroder-Bernstein Theorem	17
5.5	September 29	18
5.5.1	Zorn's Lemma	18
5.6	October 4	19

5.6.1	Axiom of Choice	19
5.7	October 6	20
5.7.1	Axiom of Choice	20
5.7.2	Applications of Axiom of Choice	20
5.8	October 11	20
5.8.1	Countable Sets	20
5.9	October 13	21
5.9.1	Cardinal Arithmetic	21
6	Orderings and Ordinals	23
6.1	October 25	23
6.1.1	Orderings	23
6.2	October 27	24
6.2.1	Induction and Recursion	24
6.3	November 1	25
6.3.1	The Replacement Axiom	25
6.4	November 3	25
6.4.1	Ordinals	25
6.5	November 8	27
6.5.1	Cumulative Hierarchy	27
6.6	November 10	29
6.6.1	Transfinite Recursion	29
6.7	November 15	29
6.7.1	Ordinals	29
6.8	November 17	30
6.8.1	Zorn's Lemma	30

Chapter 1

Introduction

1.1 August 25

1.1.1 Introduction

Foundations of Mathematics: language, axioms, formal proofs

- We focus on the axioms in set theory
- We use ZFC (Zermelo-Fraenkel + Choice)
- There is only one primitive notion : \in
- Within the ZFC universe, everything is a set

Course Outline:

- Basic axioms
- Operations, relations, functions
- $\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}$
- cardinals
- AC
- ordinals

1.1.2 Basics

Principle of Extensionality: Two sets A, B are the same \leftrightarrow they have the same elements $\forall x(x \in A \leftrightarrow x \in B)$

Example 1.1.1. $2, 3, 5 = \{5, 2, 4\} = \{2, 5, 2, 3, 3, 2\}$

Definition 1.1.2. There is a set with no elements, denoted \emptyset

- $\emptyset \neq \{\emptyset\}$
- $A \subseteq B$: A is a subset of $B \leftrightarrow$ each element of A is in B (use \subsetneq to denote proper subset)

- $\{2\} \subseteq \{2, 3, 5\}$ but $\{2\} \notin \{2, 3, 5\}$
- Power set operation: $\mathcal{P}(A) = \{B \mid B \subseteq A\}$

We can define a hierarchy:

$$V_0 = \emptyset, V_1 = \mathcal{P}(\emptyset) = \{\emptyset\}, V_2 = \mathcal{P}\mathcal{P}(\emptyset) = \{\emptyset, \{\emptyset\}\}$$

$$V_3 = \mathcal{P}(V_2) = \{\emptyset, \{\emptyset\}, \{\{\emptyset\}\}, \{\emptyset, \{\emptyset\}\}\}, V_4, \dots$$

$$V_\omega = \bigcup_{n \in \mathbb{N}} V_n, \mathcal{P}(V_\omega), \mathcal{P}\mathcal{P}(V_\omega), \dots, V_{\omega+\omega}, \dots, V_{\omega+\omega+\dots}, \dots, V_{\omega \times \omega}, \dots, V_{\omega^\omega}$$

Chapter 2

Axioms and Operations

2.1 August 30

2.1.1 Zermelo Fraenkel Axioms of Set Theory

Setting: in ZFC all objects are sets

Language: contains vocabulary (\in), logical symbols ($=, \wedge, \vee, \exists, \forall, \neg$), variables (x, y, A, B , etc.)

Axiom 2.1.1 (Extensionality Axiom). Two sets are the same if they have the same elements
 $\forall A, B (\forall x (x \in A \leftrightarrow x \in B) \rightarrow A = B)$

Axiom 2.1.2 (Empty Set Axiom). There is a set with no members, denoted \emptyset
 $\exists A \forall x (x \notin A)$

Axiom 2.1.3 (Pairing Axiom). For any sets u, v there is a set whose elements are u and v , denoted $\{u, v\}$
 $\forall u, v \exists A \forall x (x \in A \leftrightarrow x = u \vee x = v)$

Axiom 2.1.4 (Union Axiom (Preliminary Form)). For any sets a, b there is a set whose elements are elements of a and elements of b , denoted $a \cup b$
 $\forall a, b \exists A \forall x (x \in A \leftrightarrow x \in a \vee x \in b)$

Axiom 2.1.5 (Powerset Axiom). Each set A , has a power set $\mathcal{P}(A)$.
 $\forall A \exists B \forall x (x \in B \iff x \subseteq A)$ where $x \subseteq A$ stands for $\forall y (y \in x \rightarrow y \in A)$

Axiom 2.1.6 (Union Axiom). For any set A , there is a set $\bigcup A$ whose members are members of the members of A .
 $\forall A \exists B \forall x (x \in B \leftrightarrow \exists y \in A (x \in y))$

Idea for the subset axiom: For any set A , there is a set B whose members are members of A satisfying some property.

eg. $B = \{x \in A \mid x \text{ satisfies property } P\} \subseteq A$

Example 2.1.7. $B = \{n \in \mathbb{N} \mid n \text{ cannot be described in less than 20 words}\}$

- let b be the smallest element in B , then b is the smallest element that cannot be described in 20 words.
- Paradox : need to use formal language to express property P .

Example 2.1.8. Let $B = \{x \mid x \notin x\}$

Question: $B \in B$? $B \in B \leftrightarrow B \notin B$: need to have property be contained in some larger set.

We can now restate the axiom more formally:

Axiom 2.1.9 (Subset Axiom (Scheme)). For each formula $\phi(x)$, there is an axiom:
 $\forall A \exists B \forall x (x \in B \leftrightarrow x \in A \wedge \phi(x))$

Example 2.1.10. Suppose there is a set of all sets A . Consider $B = \{x \in A \mid x \notin x\}$. Then $B \in B \leftrightarrow B \notin B$, contradiction. So there can be no such set A .

The language of 1st order logic for ZFC:

The following are formulas:

- $x = y, x \in y$ atomic formulas
- $(\varphi \wedge \psi), (\varphi \vee \psi), \neg \varphi$ where φ, ψ are formulas
- $\exists v \varphi, \forall x \varphi$

Example 2.1.11. $\varphi(v, w) := (\exists v (v \in x \wedge \neg v = w)) \rightarrow (\forall y (\neg y \in y))$ is a formula

Chapter 3

Relations and Functions

3.1 September 1

3.1.1 Relations and Functions

Ordered Pair: $\langle a, b \rangle = \langle c, d \rangle \leftrightarrow a = c, b = d$

Definition 3.1.1. $\langle a, b \rangle = \{\{a\}, \{a, b\}\}$

Cartesian product of A and B , denoted $A \times B = \{\langle x, y \rangle \mid x \in A, y \in B\}$

Using the subset axiom $A \times B = \{z \in \mathcal{PP}(A \cup B) \mid \exists x \in A \exists y \in B z = \langle x, y \rangle\}$

Observation: $\langle x, y \rangle \in \mathcal{PP}(C)$ for $x, y \in C$

$\{x\}, \{x, y\} \in \mathcal{P}(C)$ so $\{\{x\}, \{x, y\}\} \subseteq \mathcal{P}(C)$ so $\{\{x\}, \{x, y\}\} \in \mathcal{PP}(C)$

Definition 3.1.2. A binary relation is a set R whose elements are ordered pairs.

If $R \subset A \times B$ then R is a relation from $A \rightarrow B$.

Definition 3.1.3. Given a relation R , $\text{dom } R = \{x \in \bigcup \bigcup R \mid \exists y \langle x, y \rangle \in R\}$,
 $\text{range } R = \{y \in \bigcup \bigcup R \mid \exists x \langle x, y \rangle \in R\}$, $\text{field } (R) = \text{dom}(R) \cup \text{range}(R)$

Example 3.1.4. $R = \{\langle a, b \rangle, \langle c, d \rangle, \langle e, f \rangle\} = \{\{\{a\}, \{a, b\}\}, \{\{c\}, \{c, d\}\}, \{\{e\}, \{e, f\}\}\}$

$\bigcup R = \{\{a\}, \{a, b\}, \{c\}, \{c, d\}, \{e\}, \{e, f\}\}$

$\bigcup \bigcup R = \{a, b, c, d, e, f\}$

n -ary relations: define n -tuple by $\langle a, b, c \rangle = \langle \langle a, b \rangle, c \rangle$ etc.

Definition 3.1.5. A function is a relation F such that $\forall x, y, z \langle x, y \rangle \in F$ and $\langle x, z \rangle \in F \rightarrow y = z$

$\forall x \in \text{dom } (F)$ there is y such that $\langle x, y \rangle \in F$. If $A = \text{dom}(F)$, $B \supseteq \text{range}(F)$ then F is said to be a function from A to B , $f : A \rightarrow B$

We say that $f : A \rightarrow B$ is onto if $B = \text{range}(F)$

Definition 3.1.6. F is injective if $\forall x, y, z \langle x, z \rangle \in F \wedge \langle y, z \rangle \in F \rightarrow x = y$.

Definition 3.1.7. For a set A , relations F, G

- (a) inverse $F^{-1} = \{\langle y, x \rangle \mid \langle x, y \rangle \in F\}$
- (b) composition: $G \circ F = \{\langle x, z \rangle \mid \exists y \langle x, y \rangle \in F, \langle y, z \rangle \in G\}$
- (c) restriction: $F \upharpoonright_A = \{\langle x, y \rangle \in F \mid x \in A\}$
- (d) image of A under F , $F[A] = \{y \mid \exists x \in A \langle x, y \rangle \in F\} = \text{range}(F \upharpoonright_A)$

Example 3.1.8. If F is a function, F^{-1} may not be a function. F^{-1} is a function $\leftrightarrow F$ is one to one.

Example 3.1.9. $F^{-1} \circ F = \{\langle x, x \rangle \mid x \in \text{dom}(F)\}$ if F is one to one

More generally, $F^{-1} \circ F = \{\langle x, z \rangle \mid \exists y \in \text{range } F \langle x, y \rangle, \langle z, y \rangle \in F\}$.

3.2 September 6

3.2.1 Functions and Relations

Theorem 3.2.1. Let $F : A \rightarrow B$ with $A \neq \emptyset$

- (a) There is a function $G : B \rightarrow A$ such that $G \circ F = \text{id}_A \leftrightarrow F$ is one to one.
- (b) There is a function $G : B \rightarrow A$ such that $F \circ G = \text{id}_B \leftrightarrow F$ is onto.

Proof. (a) Suppose there is such a G . Take a_1, a_2 such that $F(a_1) = F(a_2)$, then $a_1 = G \circ F(a_1) = G \circ F(a_2) = a_2$

Conversely, suppose F is one to one. We want to define $G : B \rightarrow A$ given $b \in B$, let $G(b)$ = the unique $a \in A$ such that $F(a) = b$ if $b \in \text{range}(F)$. If $b \notin \text{range}(F)$, let $G(b) = a_0$ with $a_0 \in A$ arbitrary (exists since A nonempty)

- (b) Suppose that $G : B \rightarrow A$, with $F \circ G = \text{id}_B$ Want to show $\forall b \in B \exists a F(a) = b$ Take $a = G(b) \rightarrow F(a) = F(G(b)) = b$

Conversely, suppose F is onto. We want to define G , given $b \in B$ want to define $G(b)$ such that $F(G(b)) = b$, equivalently, want $G(b) \in F^{-1}(\{b\})$. Since F is onto $F^{-1}(\{b\})$ is nonempty. Let $G(b)$ be any element of $F^{-1}(b)$, equivalently $G \subseteq F^{-1}$ and $\text{dom}(G) = B = \text{dom}(F^{-1})$.

Example 3.2.2. Suppose $A = \mathbb{N}$, let $G = \{(b, a) \in B \times \mathbb{N} : a \text{ is least satisfying } f(a) = b\}$

- Don't have a method to specify such elements in gneral.

Axiom 3.2.3 (Axiom of Choice - Form I). For every relation R , there is a function $G \subseteq R$ with $\text{dom}(G) = \text{dom}(R)$

3.2.2 Infinite Cartesian Products

$$A \times B = \{\langle x, y \rangle \in \mathcal{PP}(A \cup B) \mid x \in A \wedge y \in B\}$$

Definition 3.2.4. Let M be a function with domain I such that for every $i \in I$, $H(i)$ is a set. Let

$$\times_{i \in I} H(i) = \{f : I \rightarrow \bigcup_{i \in I} H(i) \mid f(i) \in H(i) \forall i \in I\}$$

Example 3.2.5. Let ω_g be $\{G \subseteq \mathbb{R} \mid 0 \neq G, G \cup \{0\} \text{ is closed under addition}\}$
 $\times_{G \in \omega_g} H(G)$ is a function such that for each $G \in \omega_g$, you get an element of G .

Observation: If one of the $H(i)$ is \emptyset , then $\times_{i \in I} H(i) = \emptyset$

Axiom 3.2.6 (Axiom of Choice - Form II). If H is a function with domain I such that $H(i) \neq \emptyset \forall i \in I$, then $\times_{i \in I} H(i) \neq \emptyset$

(ACI) \rightarrow (ACII): We are given H with $H(i) \neq \emptyset$ for all i . Want $f : I \rightarrow \bigcup_{i \in I} H(i)$ with $f(i) \in H(i) \forall i \in I$. Let $R = \{\langle i, h \rangle \in I \times \bigcup_{i \in I} H(i) \mid h \in H(i)\}$. $\text{dom}(R) = I$, since $H(i) \neq \emptyset$ there is $h \in H(i)$ so $\langle i, h \rangle \in R$. BY ACI, there is $F \subseteq R$ with $\text{dom}(F) = \text{dom}(R) = I$. $\forall i, \langle i, f(i) \rangle \in F \subseteq R$ so $f(i) \in H(i)$

Chapter 4

Naturals, Rationals, Reals

4.1 September 8

4.1.1 Natural Numbers

Idea: each natural number is the set of all the previous numbers.

$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots, n = \{0, 1, \dots, n-1\}, \dots$

Definition 4.1.1. The successor of a set a is defined as $a^+ = a \cup \{a\}$

Definition 4.1.2. A set I is inductive if $\emptyset \in I$ and $\forall a \in I, a^+ \in I$

Definition 4.1.3. a is a natural number if it belongs to all inductive sets, $\forall I (I \text{ inductive} \rightarrow a \in I)$

If I is any inductive set, let $\omega = \{a \in I \mid a \text{ belongs to all inductive sets}\}$ = the minimal inductive set.

Observation: ω is inductive because \emptyset is in all inductive sets and if n belongs to all inductive sets then so does n^+

Axiom 4.1.4 (Infinity Axiom). There is an inductive set.

Inductivon Principle: If $A \subseteq \omega$ is inductive set $A = \omega$

Example 4.1.5. Every natural number is 0 or the successor of some natural number.

Let $A = \{n \in \omega \mid n = 0 \vee \exists m \in \omega n = m^+\}$. A is inductive so $A = \omega$

Definition 4.1.6. A set A is transitive if one of the following equivalent conditions holds:

- if $x \in a \in A$, then $x \in A$
- $\bigcup A \subseteq A$
- if $a \in A$, then $a \subseteq A$
- $A \in \mathcal{P}(A)$

Example 4.1.7. Transitive sets include \emptyset , each natural number, ω , V_ω

Claim: $A = \{n \in \omega \mid n \text{ is transitive}\}$ is inductive (implies all natural numbers are transitive)

- Base: $0 \in A$ since \emptyset is transitive
- Inductive Step: Suppose $n \in A$ transitive, want to show n^+ is transitive.
Consider $x \in a \in n^+ = n \cup \{n\}$. If $a = n$, $x \in n \subseteq n^+$. If $a \in n$, $x \in a \in n$ so by transitivity $x \in n^+$ so $x \in n^+$

Theorem 4.1.8. If a is transitive, then $\bigcup a^+ = a$

Proof. $(\supseteq) a = \bigcup \{a\} \subseteq \bigcup (a \cup \{a\}) = \bigcup a^+ \quad (a \in a^+ \text{ so } a \subseteq \bigcup a^+)$
 (\subseteq) Take $x \in \bigcup a^+$, then let $b \in a^+$ with $x \in b$. If $b = a$, $x \in a$. If $b \in a$, $x \in b \in a$ so $x \in a$.

- If a, b transitive and $a^+ = b^+$ then $a = \bigcup a^+ = \bigcup b^+ = b$ so successor function is one to one on transitive sets, more specifically ω .

Fix a number $k \in \omega$. Consider the following functions:

- $A_k : \omega \rightarrow \omega$ by $A_k(0) = 0, A_k(n^+) = A_k(n)^+$
- $M_k : \omega \rightarrow \omega$ by $M_k(0) = 0, M_k(n^+) = A_k(M_k(n))$

4.2 September 13

4.2.1 Operations on the Natural Numbers

Theorem 4.2.1. Let A be a set, $a \in A$ and $F : A \rightarrow A$. Then there is a unique function $h : \omega \rightarrow A$ such that:

1. $h(0) = a$
2. $h(n^+) = F(h(n))$ for all $n \in \omega$

Proof. Let $h = \{\langle n, b \rangle \in \omega \times A \mid \text{there is } g : n^+ \rightarrow A \text{ such that } g(0) = a, g(i^+) = F(g(i)) \wedge g(n) = b\}$

Claim 1: For all n there is a $g : \{0, \dots, n\} \rightarrow A$ such that $g(0) = a, g(i^+) = F(g(i))$

Claim 2: Such a g is unique.

Proof of Claim 1. Let $I = \{n \in \omega \mid \text{such a } g \text{ exists}\}$. Want to show that I is inductive.

1. $0 \in I$: let $g : \{0\} \rightarrow A$ be such that $g(0) = a$ eg. $g = \{\langle 0, a \rangle\}$
2. Suppose $n \in I$, we know such a g exists for n , $g : \{0, \dots, n\} \rightarrow A$. We want $\tilde{g} : \{0, \dots, n, n^+\} \rightarrow A$.
Let $\tilde{g} = g \cup \{\langle n^+, F(g(n)) \rangle\}$

□

Proof of Claim 2. Suppose $g, \tilde{g} : \{0, \dots, n\} \rightarrow A$ such that $g(0) = a = \tilde{g}(0), g(i^+) = F(g(i)), \tilde{g}(i^+) = F(\tilde{g}(i)), i < n$. We want to show $g(i) = \tilde{g}(i) \forall i \leq n$. $g(0) = \tilde{g}(0), g(i^+) = F(g(i)) = F(\tilde{g}(i)) = \tilde{g}(i^+)$

Can formally show this by induction using $I = \{i \in \omega \mid i \in n^+ \wedge g(i) = \tilde{g}(i) \vee i \notin n^+\}$ □

Claim 3: $\forall n \in \omega, h(n^+) = F(H(n))$

Definition 4.2.2. Given $k \in \omega$, define $A_k : \omega \rightarrow \omega$ by $A_k(0) = k, A_k(n^+) = (A_k(n))^+$. Define $n+k = A_k(n)$. Define $M_k : \omega \rightarrow \omega$ by $M_k(0) = 0, M_k(n^+) = A_k(M_k(n))$, let $n \times k = M_k(n)$.
Let $m < n$ if $m \in n$

Theorem 4.2.3. We can show the associativity of addition: $\forall a, b, v \in \omega((a + b) + c = a + (b + c))$, commutativity of addition: $\forall a, b \in \omega a + b = b + a$, etc.

4.2.2 Integers

Let \sim be the following equivalence relation on $\omega \times \omega$ by $\langle a, b \rangle \sim \langle c, d \rangle \leftrightarrow a + d = b + c$

Define $\mathbb{Z} = \omega \times \omega / \sim$. $0_{\mathbb{Z}} = [\langle 0, 0 \rangle]$, $1_{\mathbb{Z}} = [\langle 1, 0 \rangle]$

Let $[\langle a, b \rangle] +_{\mathbb{Z}} [\langle c, d \rangle] = [\langle a+c, b+d \rangle]$. One needs to show this is well defined eg. if $\langle a, b \rangle \sim \langle a', b' \rangle, \langle c, d \rangle \sim \langle c', d' \rangle$ then $\langle a+c, b+d \rangle \sim \langle a'+c', b'+d' \rangle$

Let $[\langle a, b \rangle] \times_{\mathbb{Z}} [\langle c, d \rangle] = [\langle ac+bd, ad+bc \rangle]$

Let $E : \omega \rightarrow \mathbb{Z}$ by $E(n) = [\langle n, 0 \rangle]$

4.2.3 Rationals

Let \sim be the following equivalence relation on $\mathbb{Z} \times \mathbb{Z} \setminus \{0\}$. $\langle a, b \rangle \sim \langle c, d \rangle \iff a \times_{\mathbb{Z}} d = b \times_{\mathbb{Z}} c$

Define $\mathbb{Q} = \mathbb{Z} \times \mathbb{Z} \setminus \{0\} / \sim$. $0_{\mathbb{Q}} = [\langle 0, 1 \rangle]$, $1_{\mathbb{Q}} = [\langle 1, 1 \rangle]$

Let $[\langle a, b \rangle] \times_{\mathbb{Q}} [\langle c, d \rangle] = [\langle a \times c, b \times d \rangle]$

Let $[\langle a, b \rangle] +_{\mathbb{Q}} [\langle c, d \rangle] = [\langle ad+bc, bd \rangle]$

$E : \mathbb{Z} \rightarrow \mathbb{Q}$ by $E(z) = [\langle z, 1 \rangle]$

4.3 September 15

4.3.1 Reals (Dedekind Cuts)

Definition 4.3.1. A dedekind cut is a subset $D \subseteq \mathbb{Q}$ such that

- $\emptyset \neq D \neq \mathbb{Q}$
- D is closed downwards, if $d \in D, c < d \rightarrow c \in D$
- D has no greatest element.

Let $\mathbb{R} = \{D \in \mathcal{P}(\mathbb{Q}) \mid D \text{ is a dedekind cut}\}$

$\sqrt{2} = \{q \in \mathbb{Q} \mid q \times_{\mathbb{Q}} q < 2\}$, $e = \{q \in \mathbb{Q} \mid \exists n \in \omega q <_{\mathbb{Q}} (1 + \frac{1}{n})^N\}$ For $r \in \mathbb{R}$, $-r = \{q \in \mathbb{Q} \mid -q \in r\} \setminus \{-\sup(r)\}$

For $r_1, r_2 \in \mathbb{R}$, $r_1 \leq_{\mathbb{R}} r_2 \iff r_1 \subseteq r_2$

$r_1 \times r_2 = \{q \in \mathbb{Q} \mid \exists q \leq 0 \in r \exists b \leq 0 \in r_2 q, a \times_{\mathbb{Q}} b \text{ if } r_1, r_2 > 0, \dots$

Theorem 4.3.2. $(\mathbb{R}, 0, 1, +, \times, \leq)$ is an ordered field.

$E : \mathbb{Q} \rightarrow \mathbb{R}$ is a field embedding.

Chapter 5

Cardinal Numbers and the Axiom of Choice

5.1 September 15

5.1.1 Cardinality

Definition 5.1.1. A is equinumerous to B (written $A \approx B$) if there is a bijection $f : A \rightarrow B$

Theorem 5.1.2. For every A, B, C

- $A \approx A$
- If $A \approx B$, $B \approx B$
- If $A \approx B$, $B \approx C$ then $A \approx C$

Lemma 5.1.3. $\mathbb{Z} \approx \omega$

Proof. For $z \in \mathbb{Z}$, $f(z) = \begin{cases} -2z & z \leq 0 \\ 2z + 1 & z > 0 \end{cases}$

Lemma 5.1.4. $\mathbb{Q} \approx \omega$

Proof. $f : \omega \rightarrow \mathbb{Z} \times \mathbb{Z}^+$, $\mathbb{Q} = \mathbb{Z} \times \mathbb{Z}^+ / \sim$
 $f' : \omega \rightarrow \mathbb{Q}$, $f'(n) = \text{least } i \in \omega \text{ } g(i) \notin \{f(1), \dots, f(n-1)\}$

Lemma 5.1.5. $\mathbb{R} \approx (0, 1)_{\mathbb{R}}$

5.2 September 20

5.2.1 Cardinality

- Lemma 5.2.1.** 1. $\mathbb{N} \not\approx \mathbb{R}$
 2. For any set A , $A \not\approx \mathcal{P}(A)$

Proof. 1. Let $f : \omega \rightarrow \mathbb{R}$, claim f is not onto. Want $r \notin \text{ran}(f)$, $\forall n \in \omega r \neq f(n)$. Choose A_0 such that $f(0) \notin A_0$. Given A_n such that $f(0), \dots, f(n) \notin A_n$. Divide A_n by 2, take half that does not contain $f(n+1)$ to be A_{n+1} , then $A_0 \supset A_1 \supset A_2 \supset \dots$, $\bigcap_{n \in \omega} A_n \neq \emptyset$ and for each n , $f(n) \notin A_n$ so $f(n) \notin \bigcap A_n$

2. let $f : A \rightarrow A$. Claim f is not onto. Let $B = \{b \in A \mid b \notin f(b)\}$. Claim $B \notin \text{range}(f)$. Suppose for contradiction that $B = f(b)$ for $b \in A$, $b \in B \leftrightarrow b \notin f(b) \iff b \notin B$, contradiction.

Definition 5.2.2. A set A is finite if $\exists n \in \omega (A \approx n)$ eg. $\exists \text{next} : n \rightarrow A$ bijection. $A = \{f(0), f(1), \dots, f(n-1)\}$

Lemma 5.2.3 (Pigeonhole Principle). No finite set is equinumerous to a finite subset of itself.

Lemma 5.2.4. If B is a proper subset of $n \in \omega$ there is $m < n$ such that $B \approx m$

Proof. Use induction on n . Let $A = \{n \in \omega \mid \forall B \in n \exists m \in n B \approx m\}$.
 Claim A is inductive. $0 \in A$ trivial, $1 \in A$. $B \subsetneq \{\emptyset\} \rightarrow B = \emptyset \rightarrow B \approx 0$.
 Suppose $n \in A$, want to show $n^+ \in A$. Take $B \subsetneq n^+ = n \cup \{n\}$. If $n \in B$, $B \cap n \subseteq n$ so $\exists m < n B \cap n \approx m$ so $B \approx m^+ < n^+$. If $n \notin B$, either $B \cap n = n$ so $B \approx n < n^+$ or $B \cap n \subsetneq n$ so $\exists m < n B \cap n \approx m$ so $B \approx m^+ < n^+$.

Proof (Pigeonhole Principle). Take n , $B \subsetneq n$, $B \approx n$. Then $B \approx m$ for some $m < n$ so $m \approx n$. Let $A = \{n \mid \exists m < n m \approx n\}$. Claim A is inductive. $0 \in A$, suppose $n \in A$, want to show $n^+ \in A$. Idea: turn a bijection for $n^+ \approx m^+$ so a bijection $n \approx m$

Corollary 5.2.5. • No finite set is equinumerous to a proper subset

- ω is not finite ($\omega \approx \omega \setminus \{0\}$ by $n \mapsto n+1$)
- Every finite set is equinumerous to a unique natural number.
We call that number the cardinality of A , $\text{card}(A)$
- A subset of a finite subset is finite

Definition 5.2.6. A set κ is said to be a cardinal if

- κ is transitive (if $x \in a, a \in \kappa \rightarrow x \in \kappa$)
- \in is a linear order on κ ($\forall x, y, x \in y$ or $y \in x$ or $x = y$)
- $\forall x \in \kappa, x \not\approx \kappa$

Theorem 5.2.7. For every set A , there is a unique cardinal κ such that $A \approx \kappa$. We call this κ $\text{card}(A)$

Example 5.2.8. • $n = \{0 \in 1 \in 2 \in \dots \in n-1\}$ is a cardinal

- $\omega = \{0 \in 1 \in 2 \in \dots\}$ is a cardinal
- $\omega^+ = \{0, 1, 2, \dots\} \cup \{\omega\} \approx \omega$ is not a cardinal

Notation: $\omega = \aleph_0$, $\text{card}(\mathbb{R}) = 2^{\aleph_0}$, smallest cardinal greater than $\aleph_0 = \aleph_1$

5.3 September 22

5.3.1 Cardinals

Definition 5.3.1. Given cardinals κ and λ let

- $\kappa + \lambda = \text{card}(K \cup L)$ where K and L are disjoint sets of cardinality κ and λ
- $\kappa \cdot \lambda = \text{card}(K \times L)$ where K and L are sets of cardinality κ and λ
- $\kappa^\lambda = \{f \text{ function } L \rightarrow K\} = \text{card}({}^L K)$ where K and L are sets of cardinality κ and λ

Notation: ${}^A B = \{f : f \text{ is a function } A \rightarrow B\}$

Theorem 5.3.2. Let κ, λ, μ be cardinals

- $\kappa + (\lambda + \mu) = (\kappa + \lambda) + \mu$

Proof. Let K, L, M be disjoint sets of size κ, λ, μ . $K \cup (L \cup M) = (K \cup L) \cup M$

- $\kappa + \lambda = \lambda + \kappa$
- $\kappa \cdot (\lambda \cdot \mu) = (\kappa \cdot \lambda) \cdot \mu$

Proof. $(K \times L) \times M \rightarrow K \times (L \times M)$ by $\langle \langle k, l \rangle, m \rangle \rightarrow \langle k, \langle l, m \rangle \rangle$

- $\kappa \cdot \lambda = \lambda \cdot \kappa$
- $\kappa \cdot (\lambda + \mu) = \kappa \cdot \lambda + \kappa \cdot \mu$

Proof. $K \times (L \cup M) \approx (K \times L) \cup (K \times M)$

- $\kappa^{\lambda+\mu} = \kappa^\lambda \cdot \kappa^\mu$

$$\bullet \kappa^{\lambda \cdot \mu} = (\kappa^\lambda)^\mu$$

Proof. $F : {}^{L \times M}K \rightarrow {}^M L K$, $f : {}^{L \times M}K$, $F(g)$ = the function that maps m to $g_m : L \rightarrow K$ where $g_m(l) = g(l, m)$
 $F^{-1}(h)$ with $h : M \rightarrow ({}^L K)$ is g such that $g(l, m) = h(m)(l)$

Definition 5.3.3. A is dominated by B (written $A \leq B$) if there is a one to one function from $A \rightarrow B$

$$A \leq B \iff \text{card}(A) \leq \text{card}(B)$$

Example 5.3.4. $\bullet A \subseteq B \iff A \leq B$

$$\bullet \mathbb{N} \approx \mathbb{N} \approx \mathbb{Q} \leq \mathbb{R} \leq \mathbb{C}$$

Example 5.3.5. $\mathbb{R} \approx (0, 1)_{\mathbb{R}} \leq {}^\omega 2 \leq \mathbb{R}$

$\bullet (0, 1)_{\mathbb{R}} \leq {}^\omega 2$. Given r , let $f_r : \omega \rightarrow \{0, 1\}$ be $f_r(n)$ = n th digit of binary representation of r avoiding representations that end in all 1s.

$$\bullet {}^\omega 2 \leq \mathbb{R}, f : \omega \rightarrow 2 \mapsto \sum_{i \in \omega} f(i) \cdot 10^{-i}$$

Observation: ${}^2\omega \approx \mathcal{P}(\omega)$ $\text{card}({}^2\omega) = 2^{\aleph_0}$

5.4 September 27

5.4.1 Schroder-Bernstein Theorem

Example 5.4.1. Show that $\mathbb{R} \cup \{*\}$ and \mathbb{R} are equinumerous.

We define f by $f(*) = 0$, $f(r) = \begin{cases} r + 1 & r \in \mathbb{N} \\ r & r \in \mathbb{R} \setminus \mathbb{N} \end{cases}$

Lemma 5.4.2. If A is finite, then $\omega \leq A$

Proof. $A \neq \emptyset$ so $\exists a_0 \in A$. Let $f(0) = a_0$, $A \setminus \{a_0\} \neq \emptyset$ since $A \neq 1$ so $a_1 \in A \setminus \{a_0\}$ Let $f(1) = a_1$.
 We want $G : \{\text{finite subsets of } A\} \rightarrow A$ such that $G(F) \in A \setminus F$. Let $R = \{\langle F, a \rangle \mid F \text{ finite } a \in A \setminus F\}$
 Observation: $\text{dom}(R) = \{\text{all finite subsets of } A\}$. Since A is not finite $A \setminus F \neq \emptyset$ for all finite sets, $F \subseteq A$. Use AC to get a function $G \subseteq R$ such that $\text{dom}(G) = \text{dom}(R)$.
 Define $f : \omega \rightarrow A$ by recursion. $f(0) = a_0$, $f(n) = G(\{f(0), \dots, f(n-1)\}) \in A \setminus \{f(0), \dots, f(n-1)\}$.

Corollary 5.4.3. A set A is infinite $\leftrightarrow A$ is equinumerous to some proper subset of itself.

If A is infinite, there is 1 to 1 $f : \omega \rightarrow A$. We define a bijection $h : A \rightarrow A \setminus \{f(0)\}$ by $h(a) = \begin{cases} a & a \notin \text{dom}(f) \\ f(n+1) & a = f(n) \end{cases}$

Theorem 5.4.4 (Schroder Bernstein Theorem). If $A \leq B$, $B \leq A$, then $A \approx B$

Proof. Let $f : A \rightarrow B$ 1 to 1, $g : A \rightarrow B$ 1 to 1. We want $h : A \rightarrow B$ bijection.

Let $C_0 = A \setminus \text{ran}(g)$, let $D_0 = f[C_0]$, $C_1 = [D_0]$. $C_0 \cap C_1 = \emptyset$ because $C_0 = A \setminus \text{ran}(g)$ and $C_1 \subseteq \text{ran}(g)$. We recursively define $C_{n+1} = g[D_n]$, $D_{n+1} = [C_{n+1}]$. We see that C_n disjoint, D_n disjoint. Define

$$h(a) = \begin{cases} g(a) & a \in \bigcup_{n \in \omega} C_n \\ g^{-1} & a \in A \setminus \bigcup_{n \in \omega} C_n \end{cases} \quad . \quad f \rightarrow \bigcup_{n \in \omega} \text{ is a bijection } \bigcup C_n \rightarrow \bigcup D_n. \quad g \rightarrow \bigcup_{n \in \omega} D_n \text{ is a bijection } B \setminus \bigcup_{n \in \omega} D_n \rightarrow A \setminus A \setminus \bigcup_{n \in \omega} C_n$$

- Follows that $\mathbb{R} \approx \mathcal{P}(\omega)$

5.5 September 29

5.5.1 Zorn's Lemma

Theorem 5.5.1. For every A, B either $A \leq B$ or $B \leq A$.

Zorn's Lemma: Let \mathcal{A} be a collection of sets such that for every chain $\mathcal{C} \subseteq \mathcal{A}$, $\bigcup \mathcal{C} \in \mathcal{A}$. Then \mathcal{A} has a maximal element.

Definition 5.5.2. \mathcal{C} is a chain if for every $C, D \in \mathcal{C}$ either $C \subseteq D$ or $D \subseteq C$
 $B \in \mathcal{A}$ is maximal if there is no $C \in \mathcal{A}$ with $B \subsetneq C$

We prove the following theorem to get some practice with Zorn's Lemma

Theorem 5.5.3. Every vector space has a basis.

Proof. Let V be a vector space over a field k . $B \subseteq V$ is linearly independent if for every $v_1, \dots, v_n \in B$, distinct, k_1, \dots, k_n such that $\sum k_i v_i = 0$, $k_1 = k_2 = \dots = 0$. B is a basis if B is linearly independent and $\langle B \rangle = V$ where $\langle B \rangle = \{ \sum_{i=1}^n k_i v_i \mid v_1, \dots, v_n \in B, k_1, \dots, k_n \in k \}$

Let $\mathcal{A} = \{ B \subseteq V \mid B \text{ is linearly independent} \}$. We need to show that if $\mathcal{C} \subseteq \mathcal{A}$ is a chain then $\bigcup \mathcal{C} \in \mathcal{A}$. Consider a chain \mathcal{C} consisting of linearly independent sets. To prove that $\bigcup \mathcal{C}$ is linearly independent assume we have $v_1, \dots, v_n \in \bigcup \mathcal{C}$, $k_1, \dots, k_n \in k$ with $\sum_{i=1}^n v_i k_i = 0$. For each v_i , there is $C_i \in \mathcal{C}$ with $v_i \in C_i$. One C_i contains all the others, say C_{i_0} . $v_1, \dots, v_n \in C_{i_0}$. C_{i_0} is linearly independent so all $k_i = 0$. Now we apply Zorn's Lemma to get a maximal element $B \in \mathcal{A}$. B is a maximal linearly independent set in V . $\langle B \rangle = V$ since if there is some $v \in V \setminus \langle B \rangle$ then $B \cup \{v\}$ is linearly independent, contradicting the maximality of B .

Lemma 5.5.4. Let \mathcal{C} be a collection of functions. Then

- (i) $\bigcup \mathcal{C}$ is a function
- (ii) $\text{dom}(\bigcup \mathcal{C}) = \bigcup \{ \text{dom } f : f \in \mathcal{C} \}$
- (iii) $\text{ran}(\bigcup \mathcal{C}) = \bigcup \{ \text{ran } f : f \in \mathcal{C} \}$
- (iv) if all functions in \mathcal{C} are 1 to 1, then $\bigcup \mathcal{C}$ is one to one.

Proof. (ii): $\text{dom}(\bigcup \mathcal{C}) = \{a \mid \exists b \langle a, b \rangle \in \bigcup \mathcal{C}\} = \{a \mid \exists b \exists f \in \mathcal{C} \langle a, b \rangle \in f\} = \{a \mid \exists f (\exists b \langle a, b \rangle \in f)\} = \{a \mid \exists f \in \mathcal{C} a \in \text{dom } f\} = \bigcup \{\text{dom } f : f \in \mathcal{C}\}$

(i): $\bigcup \mathcal{C}$ is a relation. Want to show it is a function. Suppose $\langle a, b \rangle \in \bigcup \mathcal{C}$ and $\langle a, c \rangle \in \bigcup \mathcal{C}$. $\exists f \in \mathcal{C}$, $\langle a, b \rangle \in f$, $\exists g \in \mathcal{C}$ $\langle a, c \rangle \in g$. Since \mathcal{C} a chain, either $f \subseteq g$ or $g \subseteq f$. If $f \subseteq g$, $\langle a, b \rangle, \langle a, c \rangle \in g$, a function, $b = c$.

(iv): $\bigcup \mathcal{C}$ is a function. Want to show it is one to one. Suppose $\langle a, b \rangle \in \bigcup \mathcal{C}$ and $\langle c, b \rangle \in \bigcup \mathcal{C}$. $\exists f \in \mathcal{C}$, $\langle a, b \rangle \in f$, $\exists g \in \mathcal{C}$ $\langle c, b \rangle \in g$. Since \mathcal{C} a chain, either $f \subseteq g$ or $g \subseteq f$. If $f \subseteq g$, $\langle a, b \rangle, \langle c, b \rangle \in g$, a one to one, $a = c$.

5.6 October 4

5.6.1 Axiom of Choice

Theorem 5.6.1. For all set C and D , we have $C \leq D$ or $D \leq C$

Proof. Let $\mathcal{A} = \{f \subseteq C \times D \mid f \text{ is a one to one function}\}$. If $\mathcal{C} \subseteq \mathcal{A}$ is a chain $\bigcup \mathcal{C}$ is a function with $\text{dom}(\bigcup \mathcal{C}) = \bigcup \{\text{dom } f : f \in \mathcal{C}\} \subseteq C$, $\text{ran}(\bigcup \mathcal{C}) = \bigcup \{\text{ran } f : f \in \mathcal{C}\} \subseteq D$ so $\bigcup \mathcal{C} \in \mathcal{A}$. By Zorn's lemma, \mathcal{A} has a maximal element, call it F , a one to one function with $\text{dom } F \subseteq C$, $\text{ran } F \subseteq D$

Claim $\text{dom } F = C$ or $\text{ran } F = D$. If not there is $c \in C \setminus \text{dom } F$ and $d \in D \setminus \text{ran } F$. Let $G = F \cup \{\langle c, d \rangle\}$.

We see that G is a one to one function, $G \subseteq C \times D$ so $G \in \mathcal{A}$, $F \subsetneq G$, contradicting the maximality of F .

If $\text{dom } F = C$, we have $F : C \rightarrow D$ and $C \leq D$.

If $\text{dom } F = D$, we have $F : D \rightarrow C$ and $D \leq C$

Theorem 5.6.2. The following are equivalent:

1. For any relation R , there is a function $F \subseteq R$ and $\text{dom } F = \text{dom } R$
2. If H is a function, $I = \text{dom } H$, $\forall i \in I H(i) \neq \emptyset$, then $\times_{i \in I} H(i) \neq \emptyset$
3. For every set A there is a function $F : \mathcal{P}(A) \setminus \{\emptyset\} \rightarrow A$ such that $\forall B \subseteq A$, $F(B) \in B$
4. For every set \mathcal{A} of nonempty disjoint sets, there is a set C such that $\forall A \in \mathcal{A}$, $\text{ord}(C \cap A) = 1$
5. Cardinal comparability: For any sets C, D , $C \leq D$ or $D \leq C$
6. Zorn's Lemma

Proof. $1 \rightarrow 2$) Let H be a function such that $\forall i \in I$, $H(i) \neq \emptyset$. Let $R = \{\langle i, h \rangle \in I \times \bigcup H(i) \mid i \in I, h \in H(i)\}$. By (1) there is a function $F \subseteq R$ with $\text{dom } F = \text{dom } R = I$. $\forall i \in I$, $\langle i, F(i) \rangle \in F \subseteq R \rightarrow F(i) \in H(i)$ so $F \in \times_{i \in I} H(i)$

$2 \rightarrow 4$) We have a collection \mathcal{A} of disjoint nonempty subsets. We want to define H such that $H(A)$ is nonempty for $A \in \mathcal{A}$. Let $I = \mathcal{A}$, for $A \in I$, $H(A) = A$. Then $\times_{A \in I} H(A) = \times_{A \in \mathcal{A}} A$, by (2) there is $f \in \times_{A \in \mathcal{A}} A$. We claim that $C = \text{ran } f$ is as wanted. For all $A \in \mathcal{A}$, $f(A) \in A$ and if $A' \neq A$, $F(A') \in A'$ disjoint from A so $\text{ran}(f) \cap A = \{f(A)\}$

$6 \rightarrow 1$) Let R be a relation. Let $\mathcal{A} = \{f \subseteq R \mid f \text{ is a function}\}$. $\mathcal{A} \neq \emptyset$ since $\emptyset \in \mathcal{A}$. If $\mathcal{C} \subseteq \mathcal{A}$ is chain, $\bigcup \mathcal{C}$ is a function, $\bigcup \mathcal{C} \subseteq R$ so $\bigcup \mathcal{C} \in \mathcal{A}$. By (6) there is a maximal $F \in \mathcal{A}$, $F \subseteq R$ is a function.

Claim $\text{dom } F = \text{dom } R$. If not, then there is $d \in \text{dom } R \setminus \text{dom } F$. Let r be such that $\langle d, r \rangle \in R$. Then $F \cup \{\langle d, r \rangle\} \in \mathcal{A}$, $F \subsetneq F \cup \{\langle d, r \rangle\}$, contradicting maximality.

5.7 October 6

5.7.1 Axiom of Choice

Proof (Pf (cont)). $4 \rightarrow 3$) We have a set A . We want $f : \mathcal{P}(A) \setminus \{\emptyset\} \rightarrow A$, $F(B) \in B$. $B^* = \{\langle B, b \rangle : b \in B\} \approx B$. $B \neq C \rightarrow B^* \cap C^* = \emptyset$. Let $\mathcal{A} = \{B^* : B \subseteq A, B \neq \emptyset\}$. By (4) there is a set C such that $\forall B^* \in \mathcal{A}, |C \cap B^*| = 1$.

$3 \rightarrow 1$) Let R be a relation. For $a \in \text{dom } R$, we want to pick an element in $R_a = \{b \in \text{ran } R \mid \langle a, b \rangle \in R\}$. Let $A = \text{ran } R$. By (3) there is $f : \mathcal{P}(A) \setminus \{\emptyset\} \rightarrow A$, $f(B) \in B$ for all $B \subseteq A$. $F = \{\langle a, f(R_a) \rangle \mid a \in \text{dom } R\}$

5.7.2 Applications of Axiom of Choice

Want to define a measure on \mathbb{R} with the following properties.

1. $m([0, 1]) = 1$
2. $m(A + r) = m(A)$
3. $m(\bigcup_{i \in \omega} A_i) = \sum_{i \in \omega} m(A_i)$

Theorem 5.7.1. There is no $m : \mathcal{P}(\mathbb{R}) \rightarrow (\mathbb{R}^{<0} \cup \infty)$ satisfying the above conditions.

Proof. For $r, s \in [0, 1]$, let $r \sim s$ if $r - s \in \mathbb{Q}$. Let $[r] = \{s \in [0, 1] \mid r - s \in \mathbb{Q}\}$. Let $\mathcal{A} = \{[r] : r \in [0, 1]\}$. \mathcal{A} is a family of disjoint sets so by AC there is a set C such that $|C \cap [r]| = 1$ for each $[r] \in \mathcal{A}$. Assume $C \subseteq [0, 1]$.

Consider $C + q$ for $q \in \mathbb{Q}$

- disjoint since if $p \neq q \in \mathbb{Q}$, $(C + p) \cap (C + q) = \emptyset$
- $\bigcup_{q \in \mathbb{Q}} (C + q) = \mathbb{R}$
- $\bigcup_{q \in \mathbb{Q} \cap [0, 1]} (C + q) \subseteq [0, 2]$

$m(\mathbb{R}) = m(\bigcup_{q \in \mathbb{Q}} (C + q)) = \sum_{q \in \mathbb{Q}} m(C + q) = \sum_{q \in \mathbb{Q}} m(C)$, so $m(C) > 0$ since $m(\mathbb{R}) > 1$. Also, $2 = m([0, 2]) \geq m(\bigcup_{q \in \mathbb{Q} \cap [0, 1]} (C + q)) = \sum_{q \in \mathbb{Q} \cap [0, 1]} m(C + q) = \sum_{q \in \mathbb{Q} \cap [0, 1]} m(C) = \infty$, a contradiction.

5.8 October 11

5.8.1 Countable Sets

Definition 5.8.1. A set is countable if $A \leq \omega \leftrightarrow$ either $A = \emptyset$ or there is an onto function $f : \omega \rightarrow A$, ie. $A = \{f(0), f(1), f(2), \dots\}$

Observation: $\omega \times \omega \approx \omega \sqcup \omega \approx \omega$

Theorem 5.8.2. let \mathcal{A} be a countable collection of countable sets. Then $\bigcup \mathcal{A}$ is countable.

Proof. We want to define an onto function $\omega \times \omega \rightarrow \bigcup \mathcal{A}$. Since \mathcal{A} is countable there is a function $g : \omega \rightarrow \mathcal{A}$. For each $A \in \mathcal{A}$, there is an onto function $f_A : \omega \rightarrow A$. Want $F : \mathcal{A} \rightarrow$ functions from ω to $\bigcup \mathcal{A}$, $A \rightarrow f_A$, such that $f_A : \omega \rightarrow A$ is onto. $R = \{\langle A, f \rangle \mid A \in \mathcal{A} \text{ } f \text{ onto function } \omega \rightarrow A\}$ for each $A \in \mathcal{A}$ we know there is at least one f such that $\langle A, f \rangle \in R$ because A is countable so $\text{dom}(R) = \mathcal{A}$. By (AC1) there is a function $F \subseteq R$, $\text{dom}(F) = \text{dom}(R) = \mathcal{A}$. Define $H : \omega \times \omega \rightarrow \bigcup \mathcal{A}$ by $H(n, m) = f_{g(n)}(m) = F(g(n))(m) \in \bigcup \mathcal{A}$

Observation: If A is countable $A^{<\omega} = A^1 \cup A^2 \cup A^3 \cup \dots$ is countable

- $r \in \mathbb{R}$ is algebraic if it is the root of a polynomial in $\mathbb{Z}[X]$ $\{r \in \mathbb{R} : \text{algebraic}\}$ is countable.

Theorem 5.8.3. For every infinite cardinal κ , $\kappa + \kappa = \kappa$

Proof. Let K have size κ . Let $\mathcal{A} = \{f \in \mathcal{P}(\kappa \sqcup \kappa) \times \kappa \mid f \text{ is a function and there is } A \subseteq \kappa, \text{dom}(f) = A \sqcup A, \text{dom}(f) = A, f \text{ is one to one}\}$. To check the conditions for Zorn's Lemma, take a chain $\mathcal{C} \subseteq \mathcal{A}$. By the lemma, $\bigcup \mathcal{C}$ is a one to one function, $\text{dom}(\bigcup \mathcal{C}) = \bigcup \{\text{dom } f : f \in \mathcal{C}\} = \bigcup \{\text{ran } f \sqcup \text{ran } f : f \in \mathcal{C}\} = \bigcup \{\text{ran } f : f \in \mathcal{C}\} \cup \bigcup \{\text{ran } f \sqcup \text{ran } f : f \in \mathcal{C}\} = \text{ran}(\bigcup \mathcal{C}) \sqcup \text{ran}(\bigcup \mathcal{C})$ so $\bigcup \mathcal{C} \in \mathcal{A}$.

By Zorn's lemma, there is a maximal $F \in \mathcal{A}$. For this F there is $A \subseteq \kappa$, $\text{dom}(F) = A \sqcup A$, $\text{ran}(F) = A$.

- If $\kappa \setminus A$ is finite, $\text{card}(\kappa) = \text{card}(A)$ and $F : A \sqcup A \rightarrow A$ is a bijection, using a bijection $\kappa \rightarrow A$, we can build a bijection $\kappa \sqcup \kappa \rightarrow \kappa$.
- If $\kappa \setminus A$ is infinite, let $D \subseteq \kappa \setminus A$ be a countable set, let $h : D \sqcup D \rightarrow D$ be a bijection. Let $G : (A \cup D) \sqcup (A \cup D) \rightarrow A \cup D$, $G \upharpoonright_{A \sqcup A} = F$, $G \upharpoonright_{D \sqcup D} = h$. $F \subsetneq G$ contradicting that F was maximal.

5.9 October 13

5.9.1 Cardinal Arithmetic

Theorem 5.9.1. $\kappa \cdot \kappa = \kappa$ for all infinite cardinals κ

Proof. Let $\mathcal{A} = \{f \in \mathcal{P}((\kappa \times \kappa) \times \kappa) \mid f \text{ is a function } \text{dom}(f) = \text{ran } f \times \text{ran } f, \text{one to one}\}$. If $A = \text{ran } f$, f is a bijection $A \times A \rightarrow A$. We need to show \mathcal{A} satisfies the conditions to apply Zorn's Lemma. Let $\mathcal{C} \subseteq \mathcal{A}$ be a chain, we want to show $\bigcup \mathcal{C} \in \mathcal{A}$. By the lemma, $\bigcup \mathcal{C}$ is a function, $\text{dom}(\bigcup \mathcal{C}) = \bigcup \{\text{dom } f : f \in \mathcal{C}\}$, $\text{ran}(\bigcup \mathcal{C}) = \bigcup \{\text{ran } f : f \in \mathcal{C}\}$. By Zorn's Lemma, there is a maximal $F \in \mathcal{A}$. Let $A = \text{ran}(F)$, $F : A \times A \rightarrow A$ bijection. Note that A must be infinite or else $A \times A \not\approx A$.

If $A \approx \kappa$, then $\kappa \times \kappa \approx A \times A \xrightarrow{F} A \approx \kappa$ so $\kappa \times \kappa \approx \kappa$, as wanted.

If not, we want to get a contradiction with the maximality of F . If $A < \kappa$, then $A < \kappa \setminus A$, otherwise $\kappa = (\kappa \setminus A) \cup A \leq A \sqcup A \approx A < \kappa$. Take $D \subseteq \kappa \setminus A$ if size A . $A \times D \cup D \times D \cup D \times A \approx A \times A \sqcup A \times A \sqcup A \times A \xrightarrow{F} A \sqcup A \sqcup A \approx A \approx D$ so there is a bijection $h : A \times D \cup D \times D \cup D \times A \rightarrow D$, then $F \cup h \in \mathcal{A}$, contradicting the maximality of F . \square

Corollary 5.9.2. If κ and λ are infinite cardinals, $\kappa + \lambda = \kappa \times \lambda = \max(\kappa, \lambda)$

Proof. If $\kappa = \max(\kappa, \lambda)$, $\kappa \leq \kappa + \lambda \leq \kappa + \kappa = \kappa$, $\kappa \leq \kappa \times \lambda \leq \kappa \times \kappa = \kappa$.

- $\text{card}\{f : \mathbb{R} \rightarrow \mathbb{R}\} = 2^{2^{\aleph_0}}$, $\text{card}\{f : \mathbb{R} \rightarrow \mathbb{R} : f \text{ cont}\} = 2^{\aleph_0}$ since if $f, g : \mathbb{R} \rightarrow \mathbb{R}$ continuous, then $f = g \leftrightarrow f \upharpoonright_{\mathbb{Q}} = g \upharpoonright_{\mathbb{Q}}$ so $\leq {}^{\mathbb{Q}}\mathbb{R} = 2^{\aleph_0}$ and $2^{\aleph_0} \leq$ since have a constant function for each real number.

Theorem 5.9.3. For κ infinite and λ such that $2 \leq \lambda \leq 2^\kappa$, $\lambda^\kappa = 2^\kappa$

Proof. $2^\kappa \leq \lambda^\kappa \leq (2^\kappa)^\kappa = 2^{\kappa \cdot \kappa} = 2^\kappa$

Continuum Hypothesis (CH): Every uncountable subset of \mathbb{R} is equinumerous to \mathbb{R} .

Thm(Gödel): CH can't be refuted in ZFC

Thm(Cohen): CH can't be proved in ZFC

Generalized Continuum Hypothesis (GCH) : For every infinite cardinal κ , there is no λ with $\kappa < \lambda < 2^\kappa$

Chapter 6

Orderings and Ordinals

6.1 October 25

6.1.1 Orderings

Definition 6.1.1. A partial ordering is a pair $p = (D, <)$ where $< \subseteq D \times D$ and satisfies transitivity and irreflexivity. ie $\forall a, b, c \in D \ a < b \wedge b < c \rightarrow c < a$ and $\forall a \not< a$

Example 6.1.2. • $(\mathcal{P}(C), \subset)$

- $(\mathbb{N}, |)$ $a|b$ if a divides b and $a \neq b$
- $(\mathbb{R}, <)$
- $(\mathbb{N}, \triangleleft)$ where $m \triangleleft n \leftrightarrow \begin{cases} m \text{ even } n \text{ odd} \\ m, n \text{ odd } m <_{\mathbb{N}} n \\ m, n \text{ even } m <_{\mathbb{N}} n \end{cases}$
 $(0 \triangleleft 2 \triangleleft 3 \triangleleft \dots) \triangleleft (1 \triangleleft 3 \triangleleft \dots)$

Definition 6.1.3. A well ordering is a linear ordering $\langle A, < \rangle$ such that every nonempty set has a least element.

Example 6.1.4. • $(\mathbb{N}, <)$ well ordering

- $(\mathbb{N} \times \mathbb{N}, <_{\text{lex}})$ $(a, b) <_{\text{lex}} (c, d) \leftrightarrow a < c$ or $a = c$ and $b < d$ is well ordered.

Proof. Take $B \subseteq \mathbb{N} \times \mathbb{N}$ nonempty. Want to show B has $<_{\text{lex}}$ least element. Take $B_0 = \{a \mid \exists b \langle a, b \rangle \in B\} \subseteq \mathbb{N}$. Let $a_0 = \text{least in } B_0$. Let $B_1 = \{b \mid \langle a_0, b \rangle \in B\}$. Let b_0 be least in B_1 . $\langle a, b \rangle <_{\text{lex}}$ least in B .

- $(\mathbb{Z}, <)$ not well ordering
- $(\mathbb{N}, <) + (\mathbb{Z}, <)$ has least element but is not a well ordering since \mathbb{Z} has no least element.
- $[0, 1] \cap \mathbb{Q}$ not well ordering since $\{1/n : n \in \mathbb{N}\}$ has no least element.
- $A = \{a, b, \dots, z\}$, consider $(A^<, <_{\text{lex}})$. Not a well ordering

Example 6.1.5. $\mathbb{N}[x]$ the set of polynomials with coefficients in \mathbb{N} , $p(x) \triangleleft q(x)$ if $\lim_{x \rightarrow \infty} p(x) - q(x) > 0$ is a well ordering

Lemma 6.1.6. Let (A, \leq) be a linear ordering. The following are equivalent.

1. Every nonempty subset has a least element
2. There is no infinite decreasing sequence ie. no $a_0 > a_1 > a_2 > \dots \in A$

Proof. $1 \rightarrow 2$) If (2) is false, and there is a sequence $a_0 > a_1 > a_2 > \dots \in A$ $\{a_0, a_1, a_2, \dots\}$ has no least element so (1) is false.

$2 \rightarrow 1$) If (1) is false, there is nonempty $B \subseteq A$ with no least element. Let $b_0 \in B$ since b_0 is not the least element, have $b_1 \in B$ with $b_1 < b_0$, $b_2 \in B$ with $b_2 < b_1 < b_0$, \dots end up with $b_0 > b_1 > b_2 > \dots$

6.2 October 27

6.2.1 Induction and Recursion

Notation: For $t \in A$, $\text{seg } t = \{s \in A \mid s < t\}$

Theorem 6.2.1 (Transfinite Induction Principle). Let $(A, <)$ be a well ordering. Let $B \subseteq A$. If $\forall t \in a[\forall s < t (s \in B) \rightarrow t \in B]$ then $B = A$.

Proof. Take $B \subseteq A$, suppose $\forall t \in a[\text{seg } t \subseteq B \rightarrow t \in B]$. We want to show $B = A$. If not $A \setminus B$ is nonempty so it has a least element b . Since b is least in $A \setminus B$, $\forall s < b$ $s \in B$ so $\text{seg } b \subseteq B$ so $b \in B$, a contradiction.

Example 6.2.2. If $A = \omega \times \omega, <_{\text{lex}}$. Want to define $F : \omega \times \omega \rightarrow \mathbb{R}$, $F(n, m) = \sup\{F(a, b) + 2^{-b} : (a, b) < (n, m)\}$. We get $F(0, 0) = 0, F(0, 1) = 1, F(0, 2) = \frac{3}{2}, F(0, 3) = \frac{7}{4}, \dots, F(1, 0) = 2, F(1, 1) = 3, \dots, F(2, 0) = 4, \dots$

Theorem 6.2.3 (Transfinite Recursion Principle). Let $(A, <)$ be a well ordering. Given ${}^{<A}B \rightarrow B$ there is a unique function $F : A \rightarrow B$ such that $\forall t \in A$ $F(t) = G(F \upharpoonright_{\text{seg } t})$

- We define ${}^{<A}B = \{f \mid f \text{ is a function, } \text{dom } (f) = \text{seg } t \text{ for some } t \in A, \text{ran } f \subseteq B\}$

Let $A = \omega + \omega, 0, 1, 2, \dots, \omega, \omega + 1, \omega + 2, \dots$

$V_0 = \emptyset, \dots, V_{n+1} = \mathcal{P}(V_n), \dots, \bigcup_{n \in \omega} V_n = V_\omega, \dots, V_{\omega+n+1} = \mathcal{P}(V_{\omega+n})$, ie $V_\alpha = \bigcup\{\mathcal{P}(V_\beta) : \beta < \alpha\}$.

Axiom 6.2.4 (Replacement Axiom). For each first order formula $\varphi(x, y)$ if $\varphi(x, y)$ is function like on a set A then there is a set B such that $\forall y (y \in B \leftrightarrow \exists x \in A \varphi(x, y))$

Definition 6.2.5. $\varphi(x, y)$ is function like on A if $\forall x \in A \exists! y \varphi(x, y)$

6.3 November 1

6.3.1 The Replacement Axiom

- Every well ordering has a least element, every element $t \in A$ has a successor $s(t) \in A$, $s(t)$ least element $\triangleright t$
- Some elements have a predecessor (called successor element) and some don't (limit elements)

Definition 6.3.1. A formula $\varphi(x, y)$ is function like if $\forall x \exists! y \varphi(x, y)$, function like on A if $\forall x \in A \exists! y \varphi(x, y)$

Example 6.3.2. If $\varphi(x, y)$ is $y = \mathcal{P}(x)$ ie. $(\forall z(z \in y \leftrightarrow z \subseteq x))$ or $y = \text{ran}(x)$ ie. $\forall z(z \in y \leftrightarrow \exists w \langle w, z \rangle \in x)$

Axiom 6.3.3. For any formula $\varphi(x, y)$ we have an axiom $\forall A$ if φ is function like on A , $\exists B$ such that $\forall x \in A \exists y \in B \varphi(x, y)$

Let $\gamma(x, y)$ be the function $y = f(x)$. For a well ordering (A, \triangleleft) define a function E with domain A by transfinite recursion, $\forall t \in A E(t) = \text{ran}(E \upharpoonright_{\text{seg } t})$

Consider this function over the polynomials in $\mathbb{N}[x]$:

- $\text{seg}(0) = \emptyset$ so $E \upharpoonright_{\text{seg } 0} = \emptyset$ ie. $E(0) = 0$
- $\text{seg}(1) = \{0\}$, $E \upharpoonright_{\text{seg } 1} = \{\langle 0, \emptyset \rangle\}$ so $E(1) = \{\emptyset\}$
- $\text{seg}(2) = \{0, 1\}$, $E \upharpoonright_{\text{seg } 2} = \{\langle 0, \emptyset \rangle, \langle 1, \{\emptyset\} \rangle\}$ so $E(2) = \{\emptyset, \{\emptyset\}\}$
- Continuing we get $0_{\mathbb{N}}, 1_{\mathbb{N}}, 2_{\mathbb{N}}$, $E(x) = \omega$, $E(x+1) = \omega^+$, $E(x+2) = \omega^{++}, \dots$, $E(2x) = \omega + \omega$

We call $\text{ran}(E)$ the ε -image of (A, \triangleleft)

Theorem 6.3.4. Let (A, \triangleleft) be a well ordering, let $\alpha = \text{ran}(E)$. Let $\varepsilon_\alpha = \{\langle a, b \rangle \in \alpha \times \alpha \mid a \in b\}$, then E is an isomorphism $(A, \triangleleft) \rightarrow (\alpha, \varepsilon_\alpha)$, ie it is a bijection and $a \triangleleft b \leftrightarrow E(a) \in_\alpha E(b)$. Given well orderings (A, \triangleleft_A) and (B, \triangleleft_B) with ε images α and β , $(A, \triangleleft_A) \cong (B, \triangleleft_B) \leftrightarrow \alpha = \beta$

6.4 November 3

6.4.1 Ordinals

Theorem 6.4.1. For $s, t \in A$, $s \triangleleft t \leftrightarrow E(s) \in E(t)$

Theorem 6.4.2. • $\forall t \in A, E(t) \notin E(t)$

- E is one to one
- α is transitive.

It follows that E is an isomorphism $(A, \triangleleft) \rightarrow (\alpha, \varepsilon_\alpha)$

Theorem 6.4.3. Given well orderings (A, \triangleleft_A) and (B, \triangleleft_B) with ε images α and β , $(A, \triangleleft_A) \cong (B, \triangleleft_B) \leftrightarrow \alpha = \beta$.

Proof. \leftarrow) If $\alpha = \beta$ then $(A, \triangleleft_A) \cong (\alpha, \in_\alpha) = (\beta, \in_\beta) \cong (B, \triangleleft_B)$
 \rightarrow) Suppose $f : A \rightarrow B$ is an isomorphism. $E_A : A \rightarrow \alpha$, $E_B : B \rightarrow \beta$. Claim $\forall t \in A$, $E_A(t) = E_B(f(t))$.
 Use transfinite induction. Let $T = \{t \in A \mid E_A(t) = E_B(f(t))\}$, want to show $T = A$. It is enough to prove that $\forall t \in A (\text{seg } t \subseteq T \rightarrow t \in T)$. $E_A(t) = \{E_A(s) : s \in A, s \triangleleft t\} = \{E_B(f(s)) : s \in A, s \triangleleft t\} = \{E_B(s) : s \in B, s \triangleleft_B f(t)\} = E_B(f(t))$.

Definition 6.4.4. α is an ordinal if it is the ε image of some well ordering.

Theorem 6.4.5. If α is transitive, well ordered by \in , then α is the ε image of some well ordering.

Proof. If α is transitive, (α, \in_α) is a well ordering, then we claim α is the ε -image of itself, ie. the map $E : \alpha \rightarrow \alpha$ is the identity. Use transfinite induction to show that $\forall t \in \alpha$ $E(t) = t$. $E(t) = \{E(s) \mid s \in \alpha, s \in \text{seg } t\} = \{E(s) : s \in t\} = \{s \mid s \in t\} = t$

Theorem 6.4.6. Given well orderings (A, \triangleleft_A) and (B, \triangleleft_B) either

- $(A, \triangleleft_A) \cong (B, \triangleleft_B)$
- $\exists a \in A$ $(\text{seg } a, \triangleleft_A) \cong (B, \triangleleft_B)$
- $\exists b \in B$ $(A, \triangleleft_A) \cong (\text{seg } b, \triangleleft_B)$

Proof. Define $f : A \rightarrow B$, $f(a) = \min(B \setminus \text{ran}(f \upharpoonright_{\text{seg } a})) = \min\{b \in B \mid \forall s \triangleleft_A a, f(s) \neq b\}$. f is order preserving and one to one. If $B \setminus \text{ran}(f)$ is nonempty, then it has minimal element b and f is an isomorphism from (A, \triangleleft_A) to $(\text{seg } b, \triangleleft_B)$. If $B \setminus \text{ran}(f)$ is empty, $\text{dom } f = A$, then $A \cong B$. If f is not longer defined for some $a \in A$, then it is defined on $\text{seg } a$ so $(\text{seg } a, \triangleleft_A) \cong (B, \triangleleft_B)$

Theorem 6.4.7. For any ordinals α, β, γ

- Every member of α is an ordinal
 If $\alpha = \text{ran } E_A$, $a \in \alpha$ then $a = E(t)$ for some $t \in A$ so $a = \text{ran}(E \upharpoonright_{\text{seg } t})$ so a is the ε image of $\text{seg } t$
- $\alpha \in \beta \in \gamma \rightarrow \alpha \in \gamma$
- $\alpha \notin \alpha$
- Exactly one of the following holds: $\alpha \in \beta$ or $\alpha = \beta$ or $\beta \in \alpha$
- Every nonempty set of ordinals has a least element

Proof. If S is a nonepty set of ordinals take $\alpha \in S$, $S \cap \alpha \subseteq \alpha$ has a least element if nonempty. If it has a least element, then such an element is the least element of S . If it is empty, then α is the least element of S .

Theorem 6.4.8 (Burali-Forti Paradox). There is not set that contains all ordinals.

Observation:

- \emptyset is an ordinal, $n \in \omega$ and ω are ordinals.
- If α is an ordinal so is $\alpha^+ = \alpha \cup \{\alpha\}$
- If S is a set of ordinals, then $\bigcup S$ is an ordinal.

6.5 November 8

6.5.1 Cumulative Hierarchy

Want to formalize hierarchy by defining $V_\alpha = \bigcup \{\mathcal{P}(V_\beta) : \beta \in \alpha\}$. Want to define this using transfinite recursion but can't do this directly. Need approximate this function since can't have a domain ORD. For $\delta \in \text{ORD}$, $F_\delta(\alpha) = \bigcup \{\mathcal{P}(F_\delta(\beta)) : \beta \in \alpha\}$.

Theorem 6.5.1. For any ordinal δ there exists an F_δ

Proof. Transfinite recursion on (δ, \in_δ) with $y = \bigcup \{P(z) : z \in \text{ran}(x)\}$. To check that this gives our desired function we see $F(\alpha) = \bigcup \{P(z) : z \in \text{ran}(F \upharpoonright_{\text{seg } \alpha})\} = \bigcup \{\mathcal{P}(F(\beta)) : \beta \in \alpha\}$. Given δ_1, δ_2 with $\delta_1 \in \delta_2$ we claim that $F_{\delta_1}(\alpha) = F_{\delta_2}(\alpha)$ for $\alpha \in \delta_1$. Follows from the uniqueness of transfinite recursion since $F_{\delta_2} \upharpoonright_{\delta_1}$ satisfies the recursive conditions so must have $F_{\delta_2} \upharpoonright_{\delta_1} = F_{\delta_1}$

Definition 6.5.2. $V_\alpha = F_\delta(\alpha)$ for any $\delta > \alpha$

Observation

- (i) $V_\alpha = \bigcup \{\mathcal{P}(V_\beta) : \beta \in \alpha\}$
- (ii) V_α is a transitive set.

Proof. By induction: $V_\alpha = \bigcup \{\mathcal{P}(V_\beta) : \beta \in \alpha\}$. $x \in y \in V_\alpha$ so $y \in \mathcal{P}(V_\beta)$ for some $\beta \in \alpha$, $y \subseteq V_\beta$ so $x \in V_\beta$ so $x \subseteq V_\beta$ so $x \in \mathcal{P}(V_\beta)$ and so $x \in V_\alpha$

- (iii) $\alpha \in \beta \rightarrow V_\alpha \subseteq V_\beta$

Proof. $V_\beta = \bigcup \{\mathcal{P}(V_\gamma) : \gamma \in \beta\}$, $V_\alpha = \bigcup \{\mathcal{P}(V_\gamma) : \gamma \in \alpha\}$

Theorem 6.5.3. (a) $V_0 = \emptyset$

(b) $V_\alpha^+ = \mathcal{P}(V_\alpha)$ for all α

(c) If λ is a limit ordinal, then $V_\lambda = \bigcup_{\beta \in \lambda} V_\beta$

Proof. (ii) $V_{\alpha^+} = \bigcup \{\mathcal{P}(V_\beta) : \beta \in \alpha^+\} = \bigcup \{\mathcal{P}(V_\beta) : \beta \in \alpha\} \cup \mathcal{P}(V_\alpha) = V_\alpha \cup \mathcal{P}(V_\alpha) = \mathcal{P}(V_\alpha)$

(iii) if $x \in V_\lambda = \bigcup \{\mathcal{P}(V_\beta) : \beta \in \lambda\}$, then $x \in \mathcal{P}(V_\beta)$ for some $\beta \in \lambda$ so $x \in V_{\beta^+}$ and $\beta^+ \in \lambda$ so $x \in \bigcup_{\beta \in \lambda} V_\beta$.
If $x \in \bigcup_{\beta \in \lambda} V_\beta$, then $x \in V_\beta$ for $\beta \in \lambda$ so $x \subseteq V_\beta$ so $x \in \mathcal{P}(V_\beta)$ so $x \in V_{\lambda}$

Definition 6.5.4. A set S is grounded if there is some α such that $S \subseteq V_\alpha$. If S is grounded, $\text{rank}(S)$ is the least α such that $S \subseteq V_\alpha$

Observation:

(i) If A is grounded, then so are all $a \in A$ and $\text{rank}(a) \in \text{rank}(A)$

Proof. $A \subseteq V_\alpha = \bigcup \{\mathcal{P}(V_\beta) : \beta \in \alpha\}$, $a \in A \rightarrow a \in \mathcal{P}(V_\beta)$ for $\beta \in \alpha$ so $a \subseteq V_\beta$ for $\beta \in \alpha$

(ii) $\text{rank}(A) =$ the least ordinal greater than $\text{rank}(a)$ for $a \in A$

Proof. Consider A and consider $\bigcup \{\text{rank}(a)^+ : a \in A\} = \alpha$. $\alpha \leq \text{rank}(A)$ since $\text{rank}(A)$ is an upper bound for $\{\text{rank}(a)^+ : a \in A\}$. Further, $\text{rank}(A) \leq \alpha$ since for $a \in A$, $a \subseteq V_{\text{rank}(a)}$ so $a \in V_{\text{rank}(a)^+}$ and so $a \in V_\alpha$ and $A \subseteq V_\alpha$

Theorem 6.5.5. The following are equivalent

(i) (Regularity) For any nonempty set A , there is some $m \in A$ such that $A \cap m = \emptyset$

(ii) There does not exist a function f with domain ω such that $f(n^+) \in f(n)$ for all n .

(iii) Every set is grounded.

Proof. i \rightarrow ii) Suppose (ii) is false, then look at $\text{ran}(f) = A$. For any $a \in A$, $a = f(n)$ for some n but $f(n^+) \in f(n)$ so $A \cap a \neq \emptyset$

ii \rightarrow iii) Suppose there is some non grounded set a_0 , a_0 must have some non grounded element a_1 , similarly, there is $a_2 \in a_1$ non grounded, \dots

Note: to make this more formal, need to use the transitive closure, and use choice

iii \rightarrow i) For nonempty A , A is grounded. Consider $\{\text{rank}(a) : a \in A\}$, a nonempty set of ordinals so it has some least element α . Pick $m \in A$ with $\text{rank}(m) = \alpha$, then $A \cap m = \emptyset$ since any elements of m must have strictly smaller rank.

6.6 November 10

6.6.1 Transfinite Recursion

Theorem 6.6.1 (Transfinite Recursion). Let (A, \triangleleft) be a well ordering and $\gamma(x, y)$ a function like formula, then there is a function F with domain A such that $\forall t \in A \gamma(F \upharpoonright_{\text{seg } t} F(t))$. Moreover F is unique.

Proof. Let $B = \{t \in A : \exists \text{ a function } f \text{ with } \text{dom}(f) = \text{seg } t, \forall s \triangleleft t, \gamma(f \upharpoonright_{\text{seg } s}, f(s))\}$. Want to show $B = A$. Pick $t \in A$. We show that if $\text{seg } t \subseteq B, t \in B$

Lemma: For $r \triangleleft r' \in A$ if f_r has $\text{dom } f_r = \text{seg } r$ and $f_{r'}$ has $\text{dom } f_{r'} = \text{seg } r'$ then $f_r = f_{r'} \upharpoonright_{\text{seg } r}$

Proof. Let $I = \{s \in \text{seg } r \mid f_r(s) = f_{r'}(s)\}$. Want to show $I = \text{seg } r$ by transfinite induction. Take $s \in \text{seg } r$, if we have that $\forall s' \triangleleft s f_r(s') = f_{r'}(s')$, then $f_r \upharpoonright_{\text{seg } s} = f_{r'} \upharpoonright_{\text{seg } s}$. $f_r(s)$ is the unique w such that $\gamma(f_r \upharpoonright_{\text{seg } s}, w)$ and similarly for $f_{r'}(s)$ so it follows that $f_r(s) = f_{r'}(s)$ and so $I = \text{seg } r$

Case 1: $t = \text{succ}(t')$ and $t' \in B$. Let $f_t = f_{t'} \cup \{\langle t', w \rangle\}$ where w is the unique w satisfying $\gamma(f_{t'}, w)$
 Case 2: t is a limit. Let $f_t = \bigcup_{r \triangleleft t} \{f_r \mid r \in \text{seg } t\}$ (set by replacement). Well defined since by IH since $\forall r \in \text{seg } t, \exists! f$ satisfying conditions.

6.7 November 15

6.7.1 Ordinals

Let ω_1 be the set of all countable ordinals. Why is this a set?

Consider $W = \{(A, R) \in (\omega + 1) \times \mathcal{P}(\omega \times \omega) \mid (A, R) \text{ is a well ordering}\}$, a set by subset axiom. For each $R \in W$ there is a unique cardinal α , $(A, R) \cong (\alpha, \in_\alpha)$, namely the \in image of (A, R) . If α is a countable ordinal, (say infinite), there is a bijection $f : \omega \rightarrow \alpha$. Define $R = \{\langle a, b \rangle \mid f(a) \in_\alpha f(b)\}$, we get the ε image of (ω, R) is α .

- If $\alpha \in \beta \in \omega_1$, α is an ordinal, countable because $\alpha \subseteq \beta$ so $\alpha \in \omega_1$
- If $\alpha, \beta \in \omega_1$, since α, β are ordinals $\alpha \in \beta$ or $\beta \in \alpha$ or $\alpha = \beta$ so \in is a linear ordering.

It follows that ω_1 is an ordinal and $\omega_1 \notin \omega_1$ so ω_1 is not countable. For any ordinal γ , either $\gamma \in \omega_1$ so γ is countable or $\omega_1 \in \gamma$, $\omega_1 \subseteq \gamma$ so γ is uncountable so ω_1 is the least uncountable ordinal.

Theorem 6.7.1 (Hartog's Lemma). For any set A there is an ordinal α such that $\alpha \not\leq A$

Proof. Consider $\alpha = \{\beta \mid \beta \text{ cardinal}, \beta \leq A\}$. this is a set because the set of \in images of $W = \{(B, R) \in \mathcal{P}(A) \times \mathcal{P}(A \times A) \mid (B, R) \text{ is a well ordering}\}$ is a set by replacement

- α is an ordinal since if $\beta \in \alpha$, β is an ordinal, if $\gamma \in \beta \in \alpha$, γ is an ordinal, $\gamma \subseteq \beta \leq A$ so α is transitive, \in_α is a linear order so α is an ordinal
- $\alpha \notin \alpha \rightarrow \alpha \not\leq A$

Lemma 6.7.2. If S is a transitive set of ordinals, then S is an ordinal.

Given A , let A^+ be the least ordinal α , $\alpha \not\leq A$. $\gamma(x, y) \equiv y$ is the least ordinal such that $y \not\leq x$. γ is function like.

Theorem 6.7.3. If $\gamma(x, y)$ is a function like function, there is another function like $\theta(x, y)$ on the ordinals such that $\forall \alpha$ if $F = \{(\beta, \gamma) : \beta \in \alpha, \theta(\beta, \gamma)\}$ then the unique z such that $\theta(\alpha, z)$ satisfies $\gamma(F, z)$

Example 6.7.4. alephs, $\aleph_0 = \omega$, $\aleph_1 = \aleph_0^+$, \dots , $\aleph_{\alpha+1} = \aleph_\alpha^+$, $\aleph_\lambda = \bigcup_{\alpha < \lambda} \aleph_\alpha$ if λ limit

6.8 November 17

6.8.1 Zorn's Lemma

Theorem 6.8.1. The following are equivalent

1. For every relation R , there is a function $F \subseteq R$, $\text{dom } F \subseteq \text{dom } R$
3. For every set A , there is a function $F : \mathcal{P}(A) \setminus \{\emptyset\} \rightarrow A$. $\forall B \subseteq A$, $F(B) \in B$
5. For any sets C, D either $C \leq D$ or $D \leq C$
6. Zorn's Lemma
7. For every set A there is a relation \triangleleft on A such that (A, \triangleleft) is well ordered.

Proof. CC \rightarrow WO) Take a set A , Use Hartog's Lemma to get $\alpha \not\leq A$. By CC, $A \leq \alpha$ so there is a one to one function $f : A \rightarrow \alpha$. Define \triangleleft on A by $a \triangleleft b \leftrightarrow f(a) \in f(b)$. Then $(A, \triangleleft) \cong (f[A], \in)$.

WO \rightarrow 3) Take A , by WO, there is a well ordering \triangleleft on A . Define $F : \mathcal{P}(A) \setminus \{\emptyset\} \rightarrow A$ by $F = \{\langle B, b \rangle \in (\mathcal{P}(A) \setminus \{\emptyset\}) \times A : b \text{ is the } \triangleleft\text{-least element of } B\}$

1 \rightarrow 6) Consider such an \mathcal{A} . Suppose that \mathcal{A} has no maximal element. For any set $A \in \mathcal{A}$, let $F(A)$ be a set in \mathcal{A} , $A \subsetneq F(A)$. The definition of F using (1) is given by $R = \{\langle A, B \rangle \in \mathcal{A} \times \mathcal{A} : A \subsetneq B\}$ since \mathcal{A} has no maximal element, $\text{dom}(R) = \mathcal{A}$. Use (1) to get a function $F \subseteq R$, $\text{dom}(F) = \mathcal{A}$, $\forall A \in \mathcal{A}$ $F(A) \supsetneq A$.

Now, use Hartog's theorem to get an ordinal $\alpha \not\leq \mathcal{A}$. We define a function $h : \alpha \rightarrow \mathcal{A}$ by transfinite recursion. For $\beta \in \mathcal{A}$, define $H(\beta)$ using $H \upharpoonright_{\text{seg } \beta}$. We split into 3 cases:

- $\beta = 0$. $H(0) = A_0$ (since $\mathcal{A} \neq \emptyset$)
- $\beta = \gamma^+$, $H(\beta) = F(H(\gamma))$
- β limit, $H(\beta) = \bigcup_{\gamma \in \beta} H(\gamma) \in \mathcal{A}$ because $\{H(\gamma) : \gamma \in \beta\}$ is a chain.

Now, $\forall \gamma \in \beta \in \alpha$, $H(\gamma) \subsetneq H(\beta)$ so H is a one to one function, contradicting $\alpha \not\leq \mathcal{A}$.

Theorem 6.8.2. For every set A , there is a unique cardinal κ such that $\kappa \approx A$

Observation: If κ_1 and κ_2 are cardinals and $\kappa_1 \approx \kappa_2$, then $\kappa_1 = \kappa_2$

Proof. BY WO, there is a well ordering \triangleleft on A . Let α be the \in image of (A, \triangleleft) . α is an ordinal, $\alpha \cong A$.
 Let κ be the least ordinal $\cong A$.

Let γ be a formula such that \exists ordinal $\alpha, \gamma(\alpha)$. Claim: there is a least ordinal satisfying γ .
 Let $G = \{\beta \in \alpha^+ : \gamma(\beta)\}$, $\alpha \in G$, $G \subseteq \alpha^+$ so G has a least element.