

# MATH 142: Elementary Algebraic Topology

Jad Damaj

Fall 2022

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	August 24 . . . . .	3
1.1.1	What is Algebraic Topology . . . . .	3
1.1.2	Continuity . . . . .	3
1.2	August 26 . . . . .	4
1.2.1	Continuity . . . . .	4
1.2.2	Topology . . . . .	4

# Chapter 1

## Introduction

### 1.1 August 24

#### 1.1.1 What is Algebraic Topology

Recall Metric Spaces:  $(X, d)$ ,  $X$  is a set,  $d$  is a metric on  $X$  (ie.  $d : X \times X \rightarrow \mathbb{R}$ )

1.  $d(x, y) = 0$  exactly if  $x = y$
2.  $d(x, y) = d(y, x)$
3.  $d(x, z) \leq d(x, y) + d(y, z)$

Let  $V$  be a vector space, let  $\|\cdot\|$  be a norm on  $V$ , let  $d(v, w) = \|v - w\|$

- $\mathbb{R}^n$ :  $\|(r_j)\|_2 = (\sum |r_j|^2)^{\frac{1}{2}}$  - Euclidean Norm,  $\|(r_j)\|_1 = \sum |r_j|$ ,  $\|(r_j)\|_\infty = \max |r_j|$

If  $(X, d)$  is a metric space and if  $Y \subseteq X$ , let  $d^Y$  be the restriction of  $d$  to  $Y \times Y$ . Then  $(Y, d^Y)$  is a metric space.

Metric spaces  $\leftrightarrow$  geometry: length, area, size of angles.

Let  $X$  be a balloon on  $\mathbb{R}^3$

- Two natural metrics: inherited metric from  $\mathbb{R}^3$ , path-length metric (eg. length of shortest path on surface between two points)
- Consider a deformation: (Insert Figure)  
the shapes have different Euclidean distances but still have an underlying commonality
- We also observe that the balloon cannot be continuously deformed into: (Insert Figure)

We want to be able to prove such things without embedding into a metric space. This is done by attaching algebraic objects to topological spaces such that their isomorphism classes don't change under continuous deformation.

#### 1.1.2 Continuity

Let  $(X, d^X)$  and  $(Y, d^Y)$  be two metric spaces. Let  $f : X \rightarrow Y$  be a function. Let  $x_0 \in X$ . We say  $f$  is continuous at  $x_0$  if for all  $\varepsilon > 0$  there exists  $\delta > 0$  such that if  $d^X(x, x_0) < \delta$  then  $d^Y(f(x), f(x_0)) < \varepsilon$ .

- Let  $(X, d)$  be a metric space. By the open ball of radius  $r$  about  $x_0$ , we mean  $B(x_0, r) = \{x \in X : d(x, x_0) < r\}$  (closed ball is  $\{x \in X : d(x, x_0) \leq r\}$ )
- the above definition can be rephrased as: for any  $B(f(x_0), \varepsilon)$  there is an open ball  $B(x_0, \delta)$  such that if  $x \in B(x_0, \delta)$  then  $f(x) \in B(f(x_0), \varepsilon)$ .  
eg. For every open ball  $B_1$  about  $f(x_0)$  there is an open ball  $B_2$  about  $x_0$  such that if  $x \in B_2$  then  $f(x) \in B_1$

**Definition 1.1.1.** For  $(X, d)$  a metric space, by a neighborhood of a point  $x \in X$ , we mean any subset of  $X$  that contains an open ball about  $x$ .

- rephrasing the definition again we get: For any neighborhood  $N_{f(x_0)}$  of  $f(x_0)$  there is a neighborhood  $N_{x_0}$  of  $x_0$  such that if  $x \in N_{x_0}$  then  $f(x) \in N_{f(x_0)}$

**Definition 1.1.2.**  $f : X \rightarrow Y$  is continuous if it is continuous at each points of  $X$ .

## 1.2 August 26

### 1.2.1 Continuity

Recall: Given  $(X, d^X)$ ,  $(Y, d^Y)$  and  $f : X \rightarrow Y$ ,  $f$  is continuous at  $x_0$  if for any open ball  $B_1$  about  $f(x_0)$  there is an open ball  $B_2$  about  $x_0$  such that if  $x \in B_2$  then  $f(x) \in B_1$ , ie.  $B_2 \subseteq f^{-1}(B_1)$

**Definition 1.2.1.** Let  $(X, d)$  be a metric space. Let  $U \subseteq X$ . We say that  $U$  is open if for every  $x \in U$  there is an open ball  $B$  about  $x$  such that  $B \subseteq U$ , ie.  $U$  is a neighborhood of each point it contains.

We say  $f : X \rightarrow Y$  is continuous if it is continuous at each point of  $X$ .

Let  $U$  be an open set in  $Y$ ,  $x \in X$  with  $f(x) \in U$ . For each ball  $B_1$  in  $U$  about  $f(x)$ , there is an open ball about  $x$   $B_2 \subseteq X$  such that if  $x' \in B_2$  then  $f(x') \in B_1$ , ie.  $B_2 \subseteq f^{-1}(B_1) \subseteq f^{-1}(U)$

ie. if  $x \in f^{-1}(U)$  then there is an open ball  $B_2$  about  $x$  with  $B_2 \subseteq f^{-1}(U)$

ie.  $f^{-1}(U)$  is open

Conversely, if the preimage  $f^{-1}(U)$  of every open set  $U$  in  $Y$  is open, then  $f$  is continuous. This is because if  $x_0 \in X$ ,  $B_1$  an open ball about  $f(x_0)$ , then  $f^{-1}(B_1)$  is open in  $X$ .  $f(x_0) \in B_1$  so we have an open ball  $B_2 \subseteq X$  about  $x_0$  such that  $B_2 \subseteq f^{-1}(B_1)$  so  $f$  is continuous at  $x_0$ .

Thus,  $f : X \rightarrow Y$  is continuous exactly if for any open  $U$  in  $Y$ ,  $f^{-1}(U)$  is open in  $X$ .

### 1.2.2 Topology

Let  $(X, d)$  be a metric space. Let  $J$  be the collection of open subsets in  $X$  of  $d$ .  $J$  has the following properties:

1.  $X \in J$ ,  $\emptyset \in J$
2. an arbitrary, maybe infinite, union of open sets is open
3. a finite intersection of open sets is open.

*Proof of (3).* If  $U_1, \dots, U_n$  are open sets and  $x \in U_1 \cap \dots \cap U_n$  then there are  $r_1, \dots, r_n \in \mathbb{R}$  such that  $B(x, r_j) \subseteq U_j$  for  $j = 1, \dots, n$ . Let  $r = \min\{r_1, \dots, r_n\}$ , then  $B(x, r) \subseteq U_j$  for each  $j$  so  $B(x, r) \subseteq U_1 \cap \dots \cap U_n$ . Thus,  $U_1 \cap \dots \cap U_n$  is open.  $\square$

Note: This does not hold for infinite intersections, consider  $\bigcap_{i \in \mathbb{N}} B(x, \frac{1}{n}) = \{x\}$  in the plane.

This motivates the following definition:

**Definition 1.2.2.** Let  $X$  be a set. By a topology on  $X$  we mean a collection,  $\mathcal{T}$ , of subsets of  $X$  (called the open sets of the topology) satisfying **1**, **2**, and **3** above.

**Definition 1.2.3.** If  $(X, \mathcal{T}^X)$ ,  $(Y, \mathcal{T}^Y)$  are topological spaces,  $f : X \rightarrow Y$  is continuous if for every  $U \in \mathcal{T}^Y$ ,  $f^{-1}(U) \in \mathcal{T}^X$ .

**Example 1.2.4.** Given  $X$ , let  $\mathcal{T}_X$  be all subsets of  $X$ . This is called the discrete topology on  $X$ .

- This topology can also be given by the metric  $d(x, y) = 1$  if  $x \neq y$ .

**Definition 1.2.5.** If  $\mathcal{T}_1, \mathcal{T}_2$  are topologies on  $X$ , we say  $\mathcal{T}_1$  is bigger, or finer, than  $\mathcal{T}_2$  if  $\mathcal{T}_1 \supseteq \mathcal{T}_2$ .

- the discrete topology is the biggest topology on  $X$ .

**Example 1.2.6.**  $\mathcal{T} = \{X, \emptyset\}$ , called the indiscrete topology on  $X$ .

Note: this topology can not be given by a metric if  $X$  has 2 or more points.