

The FREQ Procedure

Table of PRIMARY_MORPHOLOGY by HEMISPHERE				
		HEMISPHERE(Hemisphere with respect to equator (0=South, 1=North))		Total
		0	1	
PRIMARY_MORPHOLOGY(Crater has a classifiable primary morphology (1) or does not (0))				
0	Frequency	207584	132134	339718
	Percent	54.01	34.38	88.39
	Row Pct	61.10	38.90	
	Col Pct	88.92	87.57	
1	Frequency	25865	18760	44625
	Percent	6.73	4.88	11.61
	Row Pct	57.96	42.04	
	Col Pct	11.08	12.43	
Total	Frequency	233449	150894	384343
	Percent	60.74	39.26	100.00

Statistics for Table of PRIMARY_MORPHOLOGY by HEMISPHERE

Statistic	DF	Value	Prob
Chi-Square	1	163.5012	<.0001
Likelihood Ratio Chi-Square	1	162.4070	<.0001
Continuity Adj. Chi-Square	1	163.3693	<.0001
Mantel-Haenszel Chi-Square	1	163.5007	<.0001
Phi Coefficient		0.0206	
Contingency Coefficient		0.0206	
Cramer's V		0.0206	

Fisher's Exact Test	
Cell (1,1) Frequency (F)	207584
Left-sided Pr <= F	1.0000
Right-sided Pr >= F	1.794E-37
Table Probability (P)	2.516E-38
Two-sided Pr <= P	3.441E-37

Sample Size = 384343