

AFRE 802

Statistical Methods for Agricultural, Food, & Resource Economists



Sampling distributions & the Central Limit Theorem – Part 2 of 2

(WMS Ch. 7.3, 7.5-7.6)

October 26, 2017

Nicole Mason

Michigan State University

Fall 2017

GAME PLAN

- Sampling distributions & the Central Limit Theorem (Part 2 of 2)

1. Review of the sampling distribution of the sample mean when the underlying RVs \sim normal
2. Review of chi-squared, t, and F distributions
3. The Central Limit Theorem
4. The Law of Large Numbers
5. The normal approximation to the binomial distribution

Review

MICHIGAN STATE
UNIVERSITY

Sampling distributions related to the sample mean & sample variance of a random sample of normal RVs:

If Y_1, Y_2, \dots, Y_N is a **random sample from a normal distribution** with mean, μ , and variance, σ^2 , then:

- Sample mean**

$$\bar{Y} = \frac{1}{N} \sum_{i=1}^N Y_i \sim \text{Normal}(\mu, \frac{\sigma^2}{N})$$

$$Z = \frac{\bar{Y} - \mu}{\sigma / \sqrt{N}} \sim N(0, 1)$$

- Sample variance**

$$S^2 = \frac{1}{N-1} \sum_{i=1}^N (Y_i - \bar{Y})^2$$

$$\frac{(N-1)S^2}{\sigma^2} \sim \chi^2$$

with $(N-1)$ d.f.

2

Review: t distribution & replacing σ with S in our standardized statistic for the sample mean


- We just reviewed that **when our random sample is from a normal distribution**, then:

$$Z = \frac{\bar{Y} - \mu}{\sigma / \sqrt{N}} \sim N(0, 1)$$


- If we don't know σ (which we often don't), we can replace it with S to get a new statistic, T :

$$T = \frac{\bar{Y} - \mu}{S / \sqrt{N}} \sim t \text{ distribution with } N-1 \text{ d.f.}$$

We'll use this later to test hypotheses related to μ .




See Table 5 in Appendix 3 for table of probabilities for the t distribution.



Review: Comparing the variances of 2 normal populations and the F distribution

- When **testing hypotheses about the means of 2 (potentially different) normal populations**, we often **need to compare the variances of those two populations**.
- We use **F statistics** to do this (assuming we have **independent random samples** from the two normal populations):



4

Two of the most famous theorems in probability

1. Central Limit Theorem

2. Law of Large Numbers

5

The Central Limit Theorem (CLT)

- So far we've focused on random samples $(Y_i, i=1, \dots, N)$ drawn from normal distributions
- But we often don't know the distribution from which our random sample is drawn (or our sample if from a non-normal population)
- The **Central Limit Theorem** allows us to approximate the sampling distribution of \bar{Y} regardless of the distribution of Y_i
- **CLT**: The sampling distribution of the sample mean will be approximately normal as $N \rightarrow \infty$ regardless of the distribution of Y_i


6

CLT – YouTube videos – simulation & review



- <https://www.youtube.com/watch?v=BwE2a18Th4c&feature=youtu.be>
 - Watch from 8:00 to end
- <https://www.youtube.com/watch?v=sZ0DsE4vhqk>
 - Watch from 4:44 to end

7

Another CLT example before we state it formally



What is the mean and variance of Y ?



The Central Limit Theorem (CLT) - formally

Let Y_1, Y_2, \dots, Y_N be i.i.d. distributed RVs with

$E(Y_i) = \mu$ and $V(Y_i) = \sigma^2 < \infty$, then

the distribution of $\frac{\bar{Y}_N - \mu}{\sigma / \sqrt{N}}$

converges to the standard normal as $N \rightarrow \infty$

See WMS section 7.4 for the proof if interested

Simpler statement:

\bar{Y}_N is asymptotically normally distributed

with mean μ and variance $\frac{\sigma^2}{N}$

- “Large” sample size: roughly **$N > 30$**
- Note: CLT applies to a random sample from **ANY distribution** with finite mean & variance & large N

CLT: example #1

EXAMPLE 7.8

Achievement test scores of all high school seniors in a state have mean 60 and variance 64. A random sample of $n = 100$ students from one large high school had a mean score of 58. Is there evidence to suggest that this high school is inferior? (Calculate the probability that the sample mean is at most 58 when $n = 100$.)

CLT Let Y_1, Y_2, \dots, Y_N be i.i.d. distributed RVs with $E(Y_i) = \mu$ and $V(Y_i) = \sigma^2 < \infty$, then

the distribution of $\frac{\bar{Y}_N - \mu}{\sigma / \sqrt{N}}$

converges to the standard normal as $N \rightarrow \infty$

10

CLT: example #2

EXAMPLE 7.9

The service times for customers coming through a checkout counter in a retail store are independent random variables with mean 1.5 minutes and variance 1.0. Approximate the probability that 100 customers can be served in less than 2 hours of total service time.

CLT Let Y_1, Y_2, \dots, Y_N be i.i.d. distributed RVs with $E(Y_i) = \mu$ and $V(Y_i) = \sigma^2 < \infty$, then

the distribution of $\frac{\bar{Y}_N - \mu}{\sigma / \sqrt{N}}$

converges to the standard normal as $N \rightarrow \infty$

11

The Law of Large Numbers

- We often want to know the probability that the sample mean (based on a given sample size, N) is within or outside of a certain distance of the population mean:


$$P(|\bar{Y}_N - \mu| < \varepsilon) \quad \text{or} \quad P(|\bar{Y}_N - \mu| > \varepsilon) \quad \text{for some } \varepsilon > 0$$

- What happens to the latter probability as the sample size (N) gets large?*

$$P(|\bar{Y}_N - \mu| > \varepsilon) \rightarrow 0 \text{ as } N \rightarrow \infty$$

for any $\varepsilon > 0$.

The “**law of large numbers**”:
as the sample size goes to infinity, the sample mean converges (in probability) to the population mean.



12


Another illustration of the Law of Large Numbers

- What is the population mean of the number obtained on a roll of a fair die?*

$$(1+2+3+4+5+6)/6 = 21/6 = 3.5$$

Would the sample mean be exactly 3.5 if you rolled the die only a small number of times?

What if you rolled it 1,000 times? A million times?



Normal approximation to the binomial distribution

- Recall that a binomial RV, Y , is the # of successes in n trials, where the $P(\text{success})$ on one trial is p

$$p(y) = \binom{n}{y} p^y (1-p)^{n-y}$$

- OR think of as Y as the sum of n binary variables

$$Y = \sum_{i=1}^n X_i, \quad X_i = \begin{cases} 1, & \text{if the } i\text{th trial results in success,} \\ 0, & \text{otherwise.} \end{cases}$$

- Divide both sides by n : $\frac{Y}{n} = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}$

- As n gets large, by the CLT:

$$\frac{Y}{n} = \bar{X} \sim \text{Normal}\left(p, \frac{pq}{n}\right)$$

Why is
this the
mean &
variance?

Note: This approximation works well if:

$$n > 9 \left(\frac{\text{larger of } p \text{ and } q}{\text{smaller of } p \text{ and } q} \right)$$

Normal approximation to the binomial

$$\frac{Y}{n} = \bar{X} \sim \text{Normal}\left(p, \frac{pq}{n}\right)$$

This is the distribution of Y/n . What is the distribution of Y ?

$$\frac{Y}{n} = \bar{X} \sim \text{Normal}\left(p, \frac{pq}{n}\right)$$

Normal approximation to the binomial – example

EXAMPLE 7.10

Candidate A believes that she can win a city election if she can earn at least 55% of the votes in precinct 1. She also believes that about 50% of the city's voters favor her. If $n = 100$ voters show up to vote at precinct 1, what is the probability that candidate A will receive at least 55% of their votes?

16

Summary of Chapter 7

- To make inferences about population parameters from info in a random sample from that population, we need to know the probability distributions for the statistics we are using
 - **Statistic** = a function of observable RVs in the sample and known constants
- The probability distributions of statistics are called **sampling distributions**
- We studied several sampling distributions related to the normal distribution, the sample mean, & the sample variance: **standard normal, chi-square, t , and F**
- Also studied the **law of large numbers** and the **central limit theorem** (CLT), and used the CLT to come up with the **normal approximation to the binomial**
- Up next: **estimating population parameters** from the info in a sample. After that: **hypothesis testing**

17

Use remaining class time (if any) to go over extra in-class exercises in previous lecture

18

Homework:

- WMS Ch. 7 (part 2 of 2)
 - Central limit theorem: 7.43, 7.44, 7.45, 7.48, 7.49
 - Normal approximation to the binomial: 7.71, 7.75
- **Ch. 7 HW is due on Tuesday

Next class

- Estimation (Part 1 of 2) – hurray!

Reading for next class:

- WMS Ch. 8 (sections 8.1-8.4)

19

Extra slides

20

Normal approx. to the binomial - continuity correction

- FYI – not something I will test you on
- Binomial is discrete RV, normal is continuous RV
- → If using a continuous RV to approximate a discrete one, often have to make “continuity correction”
- EX) If want to know $P(Y \leq 3)$, where $Y \sim \text{binomial}$, use $P(Y \leq 3.5)$ when convert Z . (Recall that binomial RV Y is the # of successes in N trials, so it's an integer.)

See WMS pp. 380-382
and these links for details if you
are interested:

[http://www.statisticshowto.com/
what-is-the-continuity-
correction-factor/](http://www.statisticshowto.com/what-is-the-continuity-correction-factor/)

[http://courses.wcupa.edu/rbove/
Berenson/10th%20ed%20CD-
ROM%20topics/section6_5.pdf](http://courses.wcupa.edu/rbove/Berenson/10th%20ed%20CD-ROM%20topics/section6_5.pdf)

