

Sheet 7: Hyperbolic geometry

J. Evans

The final mark out of 10 (which counts towards your grade) will be calculated as your mark on Q1 plus your mark on Q2 plus your best mark from Q3–5. I will also award stars: silver for a total mark of 12 or more on any four questions, gold for a total mark of 15 or more on all questions.

Question 1. (2 marks)

Suppose that $a, b, c, d \in \mathbf{R}$ and $z \in \mathbf{C}$. Find the imaginary part of $\frac{az+b}{cz+d}$.

Answer 1. We have

$$\frac{az+b}{cz+d} = \frac{(az+b)(c\bar{z}+d)}{|cz+d|^2} = \frac{ac|z|^2 + bd + bc\bar{z} + adz}{|cz+d|^2}.$$

The denominator is real and the imaginary part of the numerator $ac|z|^2 + bd + bc\bar{z} + adz$ is $(ad - bc) \operatorname{Im}(z)$, so the imaginary part of $(az+b)/(cz+d)$ is

$$\frac{(ad - bc) \operatorname{Im}(z)}{|cz+d|^2}.$$

Question 2. (5 marks)

- (a) Find the subgroup of $PSL(2, \mathbf{R})$ consisting of Möbius transformations which fix the point i in the upper half-plane.
- (b) Show that the subgroup you found in (a) is isomorphic to $SO(2)$, the group of rotations in 2-dimensional space.
- (c) If g is a Möbius transformation with fixed point set P and h is another Möbius transformation, what is the fixed point set of hgh^{-1} ?
- (d) Let $t_b(z) = z + b$ and consider the group $\{t_b \in PSL(2, \mathbf{R}) : b \in \mathbf{R}\}$ be the group of translations of the upper half-plane; these all have precisely one fixed point (∞) . Conjugate t_b by $h(z) = -1/z$ to get a subgroup of isometries of the hyperbolic upper half-plane which fix 0. Show that, as b varies, the orbit (under the action of this subgroup) of a point ri on the imaginary axis is a Euclidean circle centred at $ri/2$ with radius $r/2$. [Such a circle is called a horocycle.]

Answer 2. (a) Suppose $i = (ai + b)/(ci + d)$. Then $ai + b = -c + di$ so (since a, b, c, d are real) $a = d$ and $c = -b$. Moreover, as the original Möbius transformation is in $PSL(2, \mathbf{R})$, we can take $ad - bc = 1$, which becomes $a^2 + b^2 = 1$, so the most general element of $PSL(2, \mathbf{R})$ fixing i is the Möbius transformation $(z \cos \theta - \sin \theta)/(z \sin \theta + \cos \theta)$.

- (b) There is an obvious map $F: SO(2) \rightarrow G$ where G is the group of Möbius transformations in (a), namely $F \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} = \frac{z \cos \theta - \sin \theta}{z \sin \theta + \cos \theta}$. We have seen that this is a homomorphism (it's just the restriction of the usual map $SL(2, \mathbf{R}) \rightarrow PSL(2, \mathbf{R})$ to the subgroup $SO(2)$). When is a matrix in the kernel of this map? When $\frac{z \cos \theta - \sin \theta}{z \sin \theta + \cos \theta} = z$ for all z , which means $z \cos \theta - \sin \theta = z^2 \sin \theta + z \cos \theta$, or $(z^2 + 1) \sin \theta = 0$, which implies $\sin \theta = 0$. Therefore the kernel consists of the identity and minus the identity, so the image of the homomorphism is isomorphic to $SO(2)/\{\pm 1\}$. But this is again isomorphic to $SO(2)$ as there is an isomorphism $SO(2)/\{\pm 1\} \rightarrow SO(2)$ given by

$$G \left[\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \right] = \begin{pmatrix} \cos(2\theta) & -\sin(2\theta) \\ \sin(2\theta) & \cos(2\theta) \end{pmatrix}.$$

More directly, one could use the map

$$H \left[\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \right] = \frac{z \cos(\theta/2) - \sin(\theta/2)}{z \sin(\theta/2) + \cos(\theta/2)}.$$

This is still a homomorphism (same proof as for F) and it's clearly surjective, but it's now injective as well: the kernel consists of those matrices $\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ for which $\cos(\theta/2) = \pm 1$, i.e. $\theta \in 2\pi\mathbf{Z}$, but this is just the identity matrix.

- (c) The fixed point set of hgh^{-1} is $h(P)$: if $x = h(y) \in h(P)$ then $hgh^{-1}h(y) = hg(y) = h(y)$ as $y \in P$ and conversely.

- (d) When we conjugate by h , we get a subgroup of Möbius transformations of the form $ht_b h^{-1}(z) = -1/(-1/z + b) = z/(1 - bz)$. If we apply this to ri then we get $ri/(1 - bri) = ri(1 + bri)/(1 + b^2 r^2) = \frac{-br^2 + ri}{1 + b^2 r^2}$. The real part is $x = -br^2/(1 + b^2 r^2)$ and the imaginary part is $y = r/(1 + b^2 r^2)$. Therefore $x^2 + y^2 = \frac{b^2 r^2 + 1}{(1 + b^2 r^2)^2} r^2 = ry$, which gives $x^2 + (y - r/2)^2 = r^2/4$, which is the equation of a circle centred at $r/2$ with radius $r/2$.

Question 3. (3 marks)

Consider the upper half-plane model of hyperbolic 2-space. Let C_1 , C_2 and C_3 be three hyperbolic lines. We say that C_3 is a common perpendicular if C_3 intersects C_1 and C_2 orthogonally at points Q and R .

- (a) Assume there exists a common perpendicular for C_1 and C_2 . By considering the hyperbolic triangle PQR that would be formed, prove that C_1 and C_2 cannot intersect at a third point P in the upper half-plane.
- (b) When C_1 and C_2 are ultraparallel show that there is always a common perpendicular and that this is unique.

Answer 3. (a) The triangle PQR would have internal angles $\pi/2, \pi/2, \epsilon$ for some $\epsilon > 0$ and these sum to $\pi + \epsilon > \pi$ which contradicts the Gauss-Bonnet theorem.

- (b) WLOG take C_1 to be the imaginary axis. Then C_3 must be a semicircle centred at 0. There is a 1-parameter family of such semicircles, an interval of which intersect C_2 . At one end of this interval, the angle of intersection between C_3 and C_2 is zero (when they are first tangent), then it starts to increase, until it reaches π (when they are last tangent). Since this angle depends continuously on the radius of C_3 , it takes on every value between 0 and π (by the intermediate value theorem). Therefore there exists such a C_3 . If there were two, they (along with C_1 and C_2) would form a hyperbolic quadrilateral with all internal angles equal to $\pi/2$, which would contradict Gauss-Bonnet.

Question 4. (3 marks)

Consider the upper half-plane model of hyperbolic space. Let I denote the imaginary axis, so that $r_I(z) = -\bar{z}$ defines the reflection in I . Let C denote the upper unit semicircle $C = \{z \in \mathbf{C} : |z| = 1, \operatorname{Im}(z) > 0\}$. By conjugating r_I by an appropriate Möbius transformation, find the formula for the reflection r_C in the hyperbolic line C . Now let D be the upper semicircle centred at $a \in \mathbf{R}$ with radius r . By conjugating r_C by an appropriate Möbius transformation, deduce that the reflection r_D in the hyperbolic line D is given by $r_D(z) = a + \frac{r^2}{\bar{z} - a}$.

Answer 4. The Möbius transformation $g(z) = (z - 1)/(z + 1)$ sends 0 to -1 , ∞ to 1 and i to i , so it sends I to C . Therefore $r_C = g \circ r_I \circ g^{-1}$ (because this is the unique Möbius transformation other than the identity which fixes C pointwise, i.e. the unique nontrivial element in the stabiliser of C under the action of isometries on hyperbolic lines). We have $g^{-1}(z) = (z + 1)(1 - z)$, so computing, we get

$$r_C(z) = g(r_I(g^{-1}(z))) = \frac{-\frac{\bar{z}-1}{\bar{z}+1} + 1}{\frac{\bar{z}-1}{\bar{z}+1} + 1} = \frac{1}{\bar{z}}.$$

Now the Möbius transformation $h(z) = \frac{z-a}{r}$ sends a to 0 and D to C , so $r_D = h^{-1} \circ r_C \circ h$. We have $h^{-1}(z) = rz + a$, so

$$r_D(z) = h^{-1}(r_C(h(z))) = a + r^2/(\bar{z} - a)$$

as required.

Question 5. (3 marks)

Find the area of a convex hyperbolic n -gon with interior angles $\alpha_1, \dots, \alpha_n$. Prove that for every $0 < \alpha < (1 - \frac{2}{n})\pi$ there is a regular convex hyperbolic n -gon with interior angles equal to α . Sketch this n -gon in both the disc and upper-half plane models. (The disc model is probably easier!)

Answer 5. By subdividing the polygon into $n - 2$ triangles, we apply the Gauss-Bonnet theorem to each triangle and deduce that

$$\text{area}(P) = (n - 2)\pi - \sum \alpha_i.$$

Let μ be an n th root of unity and consider the points $r\mu^k$, $k = 0, 1, \dots, n - 1$. When $r < 1$, these form the vertices of a regular convex hyperbolic n -gon $P(r)$ in the disc model. As $r \rightarrow 1$, the interior angles go to zero and the area goes to $(n - 2)\pi$. As $r \rightarrow 0$ the area goes to zero, so $\alpha = (\text{area}(P(r)) - (n - 2)\pi)/n \rightarrow (1 - \frac{2}{n})\pi$. As the area and the angles change continuously in r , α takes every possible value in between 0 and $(1 - \frac{2}{n})\pi$ (by the intermediate value theorem).