

Sheet 6: More Möbius maps

J. Evans

The final mark out of 10 (which counts towards your grade) will be calculated as your mark on Q1 plus your mark on Q2 plus your best mark from Q3–5. I will also award stars: silver for a total mark of 12 or more on any four questions, gold for a total mark of 15 or more on all questions.

Question 1. (2 marks)

Suppose that z_1, z_2, z_3, z_4 are distinct points in $\mathbb{C} \cup \{\infty\}$. For which $w \in \mathbb{C} \cup \{\infty\}$ is it impossible that $[z_1, z_2; z_3, z_4] = w$?

Answer 1. Without loss of generality, we can take $z_1 = 0, z_2 = \infty, z_4 = 1$ which gives $w = [0, \infty; z, 1] = z$ and z can be anything except $0, 1, \infty$.

Question 2. (5 marks)

Let $S: S^2 \rightarrow \mathbb{C} \cup \{\infty\}$ denote the stereographic projection and let $x = (x_1, x_2, x_3)$ be a point on S^2 .

- (a) Show that $|S(x)| \leq 1$ if and only if $x_3 \leq 0$ (so that the southern hemisphere projects stereographically to the unit disc).
- (b) What is the stereographic projection of the spherical triangle with vertices $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$?
- (c) Suppose that $z = S(x_1, x_2, x_3)$ and let θ be the angle between $(0, 0, 1)$ and (x_1, x_2, x_3) . Show that $|z| = \tan(\theta/2)$. [This is the geometric origin of the famous “ $\tan(x/2)$ substitution”.]
- (d) Prove that four points in $\mathbb{C} \cup \{\infty\}$ lie on a circle if and only if their cross-ratio is real.
- (e) Deduce that if A, B, C, D are points on a circle (cyclically ordered) then the internal angles in the quadrilateral $ABCD$ at B and at D add up to π and that the angle between AB and AC equals the angle between DB and DC .

[Hint: Consider the arguments of the cross-ratios $[D, B; A, C]$ and $[A, B; C, D]$.]

Answer 2. (a) We have $|S(x)|^2 = \left| \frac{x_1 + ix_2}{1 - x_3} \right|^2 = \frac{x_1^2 + x_2^2}{(1 - x_3)^2}$ and since $x_1^2 + x_2^2 = 1 - x_3^2$ we get

$$|S(x)| = \frac{1 - x_3^2}{(1 - x_3)^2} = \frac{1 + x_3}{1 - x_3}$$

which is ≤ 1 if and only if $1 + x_3 \leq 1 - x_3$ i.e. $x_3 \leq 0$.

- (b) The vertices map to $1, i$ and ∞ . Since spherical circles project to circles and straight lines, the edges map to: the segment of the unit circle connecting 1 and i , the half-lines $\{i + it : t \geq 0\}$ and $\{1 + t : t \geq 0\}$. So the whole triangle projects to

$$\{x + iy : x^2 + y^2 \geq 1, x \geq 0, y \geq 0\}.$$

- (c) We have $x_3 = \cos \theta$ and $x_3 = \frac{1 - |z|^2}{1 + |z|^2}$ so $|z| = \sqrt{\frac{1 - \cos \theta}{1 + \cos \theta}}$ but

$$\begin{aligned} 1 - \cos \theta &= 1 - \cos^2(\theta/2) + \sin^2(\theta/2) \\ &= 2 \sin^2(\theta/2) \\ 1 + \cos \theta &= 1 + \cos^2(\theta/2) - \sin^2(\theta/2) \\ &= 2 \cos^2(\theta/2) \\ \therefore \frac{1 - \cos \theta}{1 + \cos \theta} &= \frac{\sin^2(\theta/2)}{\cos^2(\theta/2)} \\ \therefore \sqrt{\frac{1 - \cos \theta}{1 + \cos \theta}} &= \tan(\theta/2). \end{aligned}$$

- (d) Let T be the Möbius transformation taking z_1, z_2, z_3 to $0, 1, \infty$ and let $z = Tz_4$. Then the cross-ratio of the four points is z . If z_1, z_2, z_3, z_4 lie on a circle C then $0, 1, \infty, z$ lie

on a circle/straight line TC , but $0, 1, \infty$ lie on a unique circle/straight line, namely $\mathbf{R} \cup \{\infty\}$, hence $z \in \mathbf{R}$. Conversely, if $z \in \mathbf{R}$ then $0, 1, \infty, z$ lie on a common straight line and hence z_1, z_2, z_3, z_4 lie on a common circle ($T^{-1}(\mathbf{R} \cup \{\infty\})$).

- (e) The internal angle between DA and DC is the argument of $(D - C)/(D - A)$. The internal angle between BC and BA is the argument of $(B - A)/(B - C)$. The quotient $(D - A)(B - C)/(D - C)(B - A)$ is the cross-ratio $[D, B; A, C]$ which is real because A, B, C, D lie on a line, hence the difference between the arguments of $(D - A)/(D - C)$ and $(B - A)/(B - C)$ is an integer multiple $n\pi$ (the argument of a real number is an integer multiple of π). Moreover, because $ABCD$ are cyclically ordered, if D, B, C map to $0, \infty, 1$ then A goes to a negative number, which means that n is odd. Similarly, by considering the cross-ratio $[A, B; C, D]$ we get the other identity.

Question 3. (3 marks)

Let $u, v \in \mathbf{C}$ and consider the points $\pi(u), \pi(v) \in S^2 \subset \mathbf{R}^3$ where $\pi: \mathbf{C} \rightarrow S^2$ is the inverse of stereographic projection. Prove that the Euclidean distance $|\pi(u) - \pi(v)|$ in \mathbf{R}^3 is

$$|\pi(u) - \pi(v)| = \frac{2|u - v|}{\sqrt{(1 + |u|^2)(1 + |v|^2)}}.$$

[Courage - this will probably get worse before it gets better.]

Answer 3. We have $\pi(u) = \left(\frac{2u}{1+|u|^2}, \frac{1-|u|^2}{1+|u|^2} \right)$ so

$$\begin{aligned} |\pi(u) - \pi(v)| &= \left| \frac{2u}{1+|u|^2} - \frac{2v}{1+|v|^2} \right|^2 + \left(\frac{1-|u|^2}{1+|u|^2} - \frac{1-|v|^2}{1+|v|^2} \right)^2 \\ &= \frac{1}{(1+|u|^2)^2(1+|v|^2)^2} \left(4|u(1+|v|^2) - v(1+|u|^2)|^2 + \right. \\ &\quad \left. + ((1-|u|^2)(1+|v|^2) - (1+|u|^2)(1-|v|^2))^2 \right) \\ &= \frac{4}{(1+|u|^2)^2(1+|v|^2)^2} \left(|u|^2(1+|v|^2)^2 + |v|^2(1+|u|^2)^2 - \right. \\ &\quad \left. - (\bar{u}v + u\bar{v})(1+|u|^2)(1+|v|^2) + (|v|^2 - |u|^2)^2 \right) \\ &= \frac{4}{(1+|u|^2)^2(1+|v|^2)^2} \left((|u|^2 + |v|^2)(1+|u|^2)(1+|v|^2) - \right. \\ &\quad \left. - (\bar{u}v + u\bar{v})(1+|u|^2)(1+|v|^2) \right) \\ &= \frac{4|u - v|^2}{(1+|u|^2)(1+|v|^2)} \end{aligned}$$

and taking square roots gives the answer.

Question 4. (3 marks) Consider a Möbius map of the form $Tz = \frac{az+b}{-bz+\bar{a}}$ with $|a|^2 + |b|^2 = 1$. Show that if $u, v \in \mathbf{C}$ and $\pi: \mathbf{C} \cup \{\infty\} \rightarrow S^2$ is the inverse of stereographic projection then $|\pi(Tu) - \pi(Tv)| = |\pi(u) - \pi(v)|$. (You may assume the expression for $|\pi(u) - \pi(v)|$ from the previous question).

Answer 4. We have

$$|\pi(u) - \pi(v)| = \frac{2|u - v|}{\sqrt{(1 + |u|^2)(1 + |v|^2)}}.$$

so

$$\begin{aligned} |\pi(Tu) - \pi(Tv)| &= \frac{2|Tu - Tv|}{\sqrt{(1 + |Tu|^2)(1 + |Tv|^2)}} \\ &= \frac{2 \left| \frac{au+b}{-bu+\bar{a}} - \frac{av+b}{-bv+\bar{a}} \right|}{\sqrt{\left(1 + \left| \frac{au+b}{-bu+\bar{a}} \right|^2\right) \left(1 + \left| \frac{av+b}{-bv+\bar{a}} \right|^2\right)}} \\ &= \frac{2|(au+b)(-\bar{b}v+\bar{a}) - (av+b)(-\bar{b}u+\bar{a})|}{\sqrt{(|-\bar{b}u+\bar{a}|^2 + |au+b|^2)(|-\bar{b}v+\bar{a}|^2 + |av+b|^2)}} \\ &= \frac{2||a|^2u - |b|^2v - a\bar{b}uv + \bar{a}b - |a|^2v + |b|^2u - b\bar{a} + a\bar{b}uv|}{\sqrt{(|\bar{b}u|^2 + |\bar{a}|^2 + |au|^2 + |b|^2)(|\bar{b}v|^2 + |\bar{a}|^2 + |av|^2 + |b|^2)}} \\ &= \frac{2|u - v|}{\sqrt{(1 + |u|^2)(1 + |v|^2)}} \end{aligned}$$

where we have used $|a|^2 + |b|^2 = 1$ in the final step.

Question 5. (3 marks)

By considering the Möbius transformation $z \mapsto 1 - z$ (and its action on $0, 1, \infty$), show that, if A, B, C, D are four points in \mathbb{C} , then $[D, B; A, C] = 1 - [C, B; A, D]$. Deduce that if A, B, C, D lie ordered cyclically on a circle then

$$|A - B||C - D| + |D - A||B - C| = |D - B||A - C|.$$

Deduce that, in a regular pentagon $ABCDE$, the ratio of the length of a diagonal to the length of a side is $\frac{1+\sqrt{5}}{2}$ (i.e. the golden ratio).

[Hint: You may use the results in Q.2(e) to prove, for example, that $\frac{(D-A)(B-C)}{(D-C)(B-A)} = -\frac{|D-A||B-C|}{|D-C||B-A|}$ or $\frac{(D-B)(A-C)}{(D-C)(A-B)} = \frac{|D-B||A-C|}{|D-C||A-B|}$.]

Answer 5. Let f be the Möbius transformation sending D, B, C, A to $0, \infty, 1, z$ (where $z = [D, B; A, C]$) and let $g(z) = 1 - z$. Then $g \circ f$ sends D, B, C, A to $1, \infty, 0, 1 - z$, hence $[C, B; A, C] = 1 - z$, proving the first part. Now we see that

$$\frac{(D-A)(B-C)}{(D-C)(B-A)} = 1 - \frac{(D-B)(A-C)}{(D-C)(A-B)}$$

Q.2(e) gives us that we may replace all differences with their magnitudes (at the cost of introducing a sign for the term on the left-hand side), from which we deduce the required equality.

Take A, B, C, D, E to be the vertices of a regular pentagon. Then $x = |D - A| = |D - B| = |A - C| = |A - D|$ is the length of a diagonal and $y = |B - A| = |C - B| = |D - C|$ is the length of a side. The equation now gives $y^2 + xy = x^2$ so x/y satisfies $\tau^2 - \tau - 1 = 0$, which is solved by $(1 \pm \sqrt{5})/2$. Only one of these solutions is positive, namely $\frac{1+\sqrt{5}}{2}$.