

Sheet 3: Quaternions and rotations

J. Evans

The final mark out of 10 (which counts towards your grade) will be calculated as your mark on Q1 plus your mark on Q2 plus your best mark from Q3–5. I will also award stars: silver for a total mark of 12 or more on any four questions, gold for a total mark of 15 or more on all questions.

Question 1. (2 marks)

In each case, find unit quaternions q such that the map $\text{Im}(\mathbf{H}) \rightarrow \text{Im}(\mathbf{H}), x \mapsto qxq^{-1}$, is the specified orthogonal transformation:

(a) $\begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$

(b) $\begin{pmatrix} 1/\sqrt{2} & 0 & -1/\sqrt{2} \\ 0 & 1 & 0 \\ 1/\sqrt{2} & 0 & 1/\sqrt{2} \end{pmatrix}.$

Answer 1. (a) This first matrix is a $\pi/2$ -rotation around the z -axis, so should be effected by the quaternion $\cos(\pi/4) + k \sin(\pi/4) = \frac{1}{\sqrt{2}}(1 + k)$. We can check that this is indeed the case:

$$\begin{aligned} \frac{1}{\sqrt{2}}(1 + k)(ix + jy + kz) \frac{1}{\sqrt{2}}(1 - k) &= \frac{1}{2}((i + j)x + (j - i)y + (k - 1)z)(1 - k) \\ &= \frac{1}{2}((i + j + j - i)x + (j - i - i - j)y + (k - 1 + 1 + k)z) \\ &= -iy + jx + kz \end{aligned}$$

so $(1, 0, 0)$ is sent to $(0, 1, 0)$, $(0, 1, 0)$ is sent to $(-1, 0, 0)$ and $(0, 0, 1)$ is sent to $(0, 0, 1)$, as required.

(b) This matrix corresponds to a $-\pi/4$ -rotation around the y -axis, so should be effected by the quaternion $\cos(\pi/8) - j \sin(\pi/8)$.

Question 2. (5 marks)

- (a) Let $q_1 = t_1 + ix_1 + jy_1 + kz_1$ and $q_2 = t_2 + ix_2 + jy_2 + kz_2$. Show that

$$\operatorname{Re}(\bar{q}_1 q_2) = t_1 t_2 + x_1 x_2 + y_1 y_2 + z_1 z_2.$$

- (b) Suppose that $t_1 = t_2 = 0$. Show that $q_1 q_2 = -q_1 \cdot q_2 + q_1 \times q_2$, i.e. the real part equals $-(x_1 x_2 + y_1 y_2 + z_1 z_2)$ and the imaginary part equals $i(y_1 z_2 - y_2 z_1) + j(z_1 x_2 - z_2 x_1) + k(x_1 y_2 - x_2 y_1)$.
- (c) Show that $A: G \times G \rightarrow \operatorname{Maps}(\mathbf{H}, \mathbf{H})$, $A(g, h)x = gxh^{-1}$ defines an action of $G \times G$ on \mathbf{H} .
- (d) Show that $A(g): x \mapsto gxh^{-1}$ is an isometry of \mathbf{H} when $g, h \in G$.
- (e) Parts (c) and (d) imply that we have a homomorphism $G \times G \rightarrow O(4)$. Show that the kernel of this homomorphism is $\{(1, 1), (-1, -1)\}$. [Hint: Consider the effect of the isometry $x \mapsto gxh^{-1}$ on the quaternions $x = 1, x = i, x = j$]

Answer 2. (a) We have

$$(t_1 - ix_1 - jy_1 - kz_1)(t_2 + ix_2 + jy_2 + kz_2) = t_1 t_2 - i^2 x_1 x_2 - j^2 y_1 y_2 - k^2 z_1 z_2 + i(\dots) + j(\dots) + k(\dots).$$

- (b) We have

$$(ix_1 + jy_1 + kz_1)(ix_2 + jy_2 + kz_2) = -(x_1 x_2 + y_1 y_2 + z_1 z_2) + ijx_1 y_2 + jiy_1 x_2 + \dots$$

and $ij = -ji = k$, etc.

- (c) We need to show that $A(1, 1)(x) = 1x1 = x$ and that

$$\begin{aligned} A(g_1, h_1)A(g_2, h_2)x &= A(g_1, h_1)g_2 x h_2^{-1} \\ &= g_1 g_2 x h_2^{-1} h_1^{-1} \\ &= g_1 g_2 x (h_1 h_2)^{-1} \\ &= A(g_1 g_2, h_1 h_2)x. \end{aligned}$$

- (d) We have $|gxh^{-1}| = |g||x||h^{-1}| = |x|$ since $|g| = |h| = |h^{-1}| = 1$.

- (e) (g, h) is in the kernel of this homomorphism if $gxh^{-1} = x$ for all x , i.e. $gx = xh$. For $x = 1$ this implies $g = h$. For $x = i$ this implies $gi = ig$. If $g = a + ib + jc + kd$ then $gi - ig = ai - b - kc + jd - ia + b - kc + jd = 2(jd - kc)$ so $gi - ig = 0$ implies $d = c = 0$. Similarly, $gj = jg$ implies $b = 0$. Therefore g is real and has length 1, so it must be ± 1 . This leaves the two possibilities $(1, 1)$ and $(-1, -1)$.

Question 3. (3 marks)

Suppose that y is a unit quaternion and let $H \subset \mathbf{H}$ be the plane orthogonal to y . Show that the reflection in the plane H is given in terms of quaternionic multiplication by $z \mapsto -y\bar{z}y$.

Answer 3. We know that $z \cdot y = \text{Re}(\bar{z}y)$ so

$$\begin{aligned} z - 2(z \cdot y)y &= z - 2\text{Re}(\bar{z}y)y \\ &= z - y(\bar{z}y + \bar{y}z)y \\ &= z - y\bar{z}y + y\bar{y}z \text{ as } y \text{ commutes with real numbers} \\ &= -y\bar{z}y \text{ as } |y| = 1. \end{aligned}$$

Question 4. (3 marks) Consider the 2-by-2 complex matrices:

$$\sigma_1 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}.$$

Show that $\sigma_1^2 = \sigma_2^2 = \sigma_3^2 = -1$ and that $\sigma_a \sigma_b = \sigma_c = -\sigma_b \sigma_a$ where a, b, c is a cyclic permutation of 1, 2, 3.

Consider the group of 2-by-2 complex matrices

$$SU(2) = \{A : A^\dagger = A^{-1}, \det A = 1\} \text{ where } \dagger \text{ denotes conjugate transpose.}$$

Show that if $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SU(2)$ then $c = -\bar{b}$, $d = \bar{a}$ and $|a|^2 + |b|^2 = 1$. If $a = t + ix$ and $b = y + iz$ then show that

$$\begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix} = t\mathbf{1} + x\sigma_1 + y\sigma_2 + z\sigma_3$$

and deduce that $SU(2)$ is isomorphic to the group of unit quaternions.

Answer 4. The first part is just a matrix computation. For the second part, observe that

$$A^\dagger = \begin{pmatrix} \bar{a} & \bar{c} \\ \bar{b} & \bar{d} \end{pmatrix}, \quad A^{-1} = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$

so if $A \in SU(2)$ then $c = -\bar{b}$ and $d = \bar{a}$. Moreover, $\det A = ad - bc = a\bar{a} + b\bar{b} = |a|^2 + |b|^2 = 1$. The next line is just obtained by substituting the matrices σ_m in and expanding. Finally, we use the isomorphism

$$G \rightarrow SU(2), \quad t + ix + jy + kz \mapsto t\mathbf{1} + x\sigma_1 + y\sigma_2 + z\sigma_3.$$

This is clearly a bijection between the unit quaternions and the matrices in $SU(2)$ (the unit condition becomes the determinant one condition) so we just need to show it is a homomorphism. The multiplication

$$(t_1 + x_1\sigma_1 + y_1\sigma_2 + z_1\sigma_3)(t_2 + x_2\sigma_1 + y_2\sigma_2 + z_2\sigma_3)$$

is clearly bilinear in the coefficients and therefore it suffices to check that it gives the correct answer on a basis. This is what we did in the first part of the question.

Question 5. (3 marks)

We define the *exponential* of a purely imaginary quaternion q by the formula

$$e^q := \sum_{m=0}^{\infty} \frac{1}{m!} q^m.$$

If $q = u\theta$ for $|u| = 1$, prove that $e^q = \cos \theta + u \sin \theta$.

Check that

$$e^{i\pi/2} e^{j\pi/2} \neq e^{(i+j)\pi/2}$$

(i.e. the usual law of logarithms fails).

[This is just the beginning of a much more general story of exponential maps for groups. For more, see the 4th year course **Lie groups and Lie algebras**.]

Answer 5. We have

$$\begin{aligned} e^{u\theta} &= 1 + u\theta + \frac{1}{2!} u^2 \theta^2 + \frac{1}{3!} u^3 \theta^3 + \dots \\ &= \left(1 - \frac{1}{2!} \theta^2 + \dots\right) + u \left(\theta - \frac{1}{3!} \theta^3 + \dots\right) \\ &= \cos \theta + u \sin \theta. \end{aligned}$$

since $u^2 = -1$.

For the second part we have $e^{i\pi/2} = i$, $e^{j\pi/2} = j$, $ij = k$ and $e^{(i+j)\pi/2} = e^{\frac{i+j}{\sqrt{2}} \frac{\pi}{\sqrt{2}}} = \cos\left(\frac{\pi}{\sqrt{2}}\right) + \frac{i+j}{\sqrt{2}} \sin\left(\frac{\pi}{\sqrt{2}}\right)$ so $e^{i\pi/2} e^{j\pi/2} \neq e^{(i+j)\pi/2}$.