





Model 4: published SRI, published implementation (n = 7085)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100



Model 4: released classification, unsmoothed, Phillips formula (n = 7076)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100



Model 4: released classification, unsmoothed, GGIR (n = 7071)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100


Model 4: released classification, unsmoothed, sleepreg, 5 days (n = 3673)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100


Model 4: released classification, unsmoothed, sleepreg, 3 days (n = 5342)






Model 4: rolling majority, 1 h, Phillips formula (n = 7076)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100



Model 4: rolling majority, 1 h, GGIR (n = 7071)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100


Model 4: rolling majority, 1 h, sleepreg, 5 days (n = 3673)





Model 4: rolling majority, 1 h, sleepreg, 3 days (n = 5342)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100



Model 4: GGIR SRI1 smoothing, 1 h, 80%, Phillips formula (n = 7076)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100



Model 4: GGIR SRI1 smoothing, 1 h, 80%, GGIR (n = 7071)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100



Model 4: GGIR SRI1 smoothing, 1 h, 80%, sleepreg, 5 days (n = 3673)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100



Model 4: GGIR SRI1 smoothing, 1 h, 80%, sleepreg, 3 days (n = 5342)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100


Model 4: awakenings <30 min bridged, Phillips formula (n = 7076)






Model 4: awakenings <30 min bridged, GGIR (n = 7073)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100


Model 4: awakenings <30 min bridged, sleepreg, 5 days (n = 3854)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100



Model 4: awakenings <30 min bridged, sleepreg, 3 days (n = 5423)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100



Model 4: main sleep period only, Phillips formula (n = 7076)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100



Model 4: main sleep period only, GGIR (n = 7073)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100



Model 4: main sleep period only, sleepreg, 5 days (n = 3854)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100


Model 4: main sleep period only, sleepreg, 3 days (n = 5423)




Model 4: random sample, PAXPREDM (n = 997)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100


Model 4: random sample, GGIR SIB (n = 997)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100


Model 4: random sample, GGIR SPT (n = 997)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100


Model 4: random sample, SWaN (n = 997)
Mexican American
Other Hispanic
Non−Hispanic White
Non−Hispanic Black
Non−Hispanic Asian
Other, incl. multiracial
men
women
Predicted BMI (kg/m²)
SRI
20
25
30
35
40
0
20
40
60
80
100