

1 Text S1. Transparency statement for secondary data analysis (after van den Akker et al. 2021)

Prior knowledge of the data. All 48 deposits were downloaded on 28 August 2026 and inspected for file structure before the grid of decision nodes was written on the same day. No specification had been computed at that point. The published analyses were extracted from the papers on 12 September 2026, and the mapping rules of Methods 2.3 were fixed after that extraction.

Order of analyses. First complete multiverse run: 28 August 2026, with the ad hoc mapping (registry v1). Rule-based mapping (registry v2) and rerun: 12 September 2026. Statistical review and corrections (pooled within-group denominator, reproduction rule, multiple imputation): 12 September 2026. Adversarial review, reclassification of outlier rules, addition of the strict grid, long-format baseline correction, refit of the published models: 13 September 2026. Displacement analysis: 13 September 2026.

Decisions made after seeing results. Outlier rules were moved from the core grid to a sensitivity arm after a winsorising rule on a bounded outcome had produced a range of 1.08 standard deviations. The strict grid was added after the first run showed that in most trials only baseline handling could vary. The core grid was renamed from principled to estimand-preserving after the precision criterion had been applied to baseline handling. The reproduction rule (2% and 10% relative difference) was fixed before any computation. An absolute criterion introduced on 12 September was withdrawn on 13 September and the prespecified rule reinstated, with the absolute difference reported without threshold.

Registry v1 versus v2. Table S2 lists both mappings, Table S7 the effect of v1 on the results. The sign of the median effect reversed in 7 of 16 trials.

Coders. One coder mapped all trials.

2 Table S1. Exclusions at manual assessment (n = 32)

study_id	category	reason_code	reason
2016_early-sitting-in-ischemic-stroke-patients-sevel	design	binary_outcome	Primary outcome is the proportion with modified Rankin score 0-2, i.e. binary; the grid is built for continuous outcomes.
2017_predicting-outcome-in-frozen-shoulder-shoulder-c	duplicate	duplicate_data	Uses the same dataset as the adhesive capsulitis study (106 rows, Group 1/2/3, identical SPADI columns); double counting.
2020_an-age-adapted-plyometric-exercise-program-impro	design	feasibility_primary	Feasibility study; primary endpoint is the feasibility of the programme.
2021_home-based-exercise-for-people-living-with-frail	design	feasibility_primary	Primary endpoints are eligibility, recruitment, adherence and attrition; no effect estimate.
2022_dependence-of-learning-outcomes-in-flipped-and-1	design	not_clinical	Educational research on teaching formats; not a clinical study in patients.
2022_a-comparison-of-psychological-characteristics-in	design	not_rct	Cross-sectional/cohort comparison of psychological characteristics; no randomised intervention.
2025_subphenotypic-classification-of-covid-19-survivo	duplicate	duplicate_data	Latent-class secondary analysis of the same TEREKO dataset (119 rows, group_randomized, SF12 columns).
2025_isoenergetic-pre-exercise-meals-varying-in-carbo	design	crossover_design	Crossover design; the grid assumes parallel groups and would ignore the within-person structure.
2021_impact-of-assessment-and-intervention-by-a-health	data	outcome_not_identifiable	Primary outcome per abstract is ED length of stay; no column in the deposit is clearly identifiable as such.
2024_effects-of-anodal-tdcs-on-resting-state-eeg-powe	data	outcome_data_not_tabular	The table file contains only baseline characteristics; the EEG outcomes are in 7z archives.

study_id	category	reason_code	reason
2022_influence-of-the-statistical-significance-of-res	design	not_clinical	Reader study on how readers interpret abstracts with/without spin; no intervention in patients.
2021_the-impact-of-a-physician-s-recommendation-and-g	design	not_clinical	Qualtrics vignette survey on physician recommendation and gender; no clinical intervention.
2021_effect-of-an-inverted-seated-position-with-upper	design	crossover_design	Four conditions per person with rest/pre/post; crossover, no parallel groups.
2023_a-clinical-model-to-predict-the-progression-of-k	design	not_rct	Prognostic model for the course of knee osteoarthritis; no randomisation.
2025_comparative-effects-of-combined-aerobic-and-resi	data	no_ipd	Deposit contains only a results table with group differences and CIs, no individual participant data.
2025_comparing-structured-note-taking-and-multiple-ch	design	not_clinical	Educational research on note-taking techniques; not a clinical study.
2026_effects-of-modest-carbohydrate-energy-supplement	design	crossover_design	Each person receives both supplements (Supplement_1 and Supplement_2 in all 20 rows, 10 C/10 P each); crossover, not parallel groups.
2026_preliminary-efficacy-of-cognitive-multisensory-r	data	arm_not_in_data	Paper randomises to immediate CMR (Group A) vs observation (Group B), but the deposit has no group column.
2025_time-restricted-eating-without-exercise-enhances	design	crossover_design	Three phases per person (Phase 1, 2, 3) with no group variable; within-person design.
2024_eight-weeks-of-high-intensity-interval-training	data	file_not_readable	Both CSV files in the deposit cannot be read as a table (inconsistent number of columns).
2022_the-effect-of-mindfulness-on-the-inflammatory-ps	data	no_data_in_deposit	The archive linked in the availability statement contains only a README.md and no data.
2025_a-sensor-augmented-telerehabilitation-system-for	data	outcome_data_not_tabular	Deposit contains only raw EMG and IMU signals as NumPy arrays per participant/trial, no outcome table.
2026_effects-of-low-load-blood-flow-restriction-train	data	arm_not_in_data	Data held as extdata of an R package in Parquet/Excel files; the contractility table lacks a group variable.
2023_the-effects-of-creatine-supplementation-on-cogni	design	crossover_design	Supplement order per person (order_supplement, guess_first_supplement); crossover, plus four separate test files.
2025_the-effect-of-proximity-to-failure-on-perceptual	design	within_person_design	Both limbs per person receive different protocols (FAIL vs RIR); within-person design.
2026_anatomy-of-a-failure-a-retrospective-evaluation	data	arm_not_in_data	Retrospective analysis; CONDITION codes measurement timepoints (1w, 1m, 3m, 6m, 12m); no group variable found.
2025_lengthened-partial-repetitions-elicite-similar-mu	design	within_person_design	Both arms per person receive different protocols (pROM/fROM contralateral); no parallel-group control arm.
2024_pilot-randomized-controlled-trial-of-i-lymfit-i	design	feasibility_primary	Pilot feasibility RCT (n=26); primary aim is feasibility benchmarks, no efficacy primary outcome.
2022_understanding-how-individualised-physiotherapy-o	no_estimate	no_group_estimate	Exploratory network analysis (mixed graphical model) over 10 ODI items; no group-difference estimate.
2025_effects-of-an-aquatic-protocol-on-electromyograp	data	outcome_not_identifiable	No primary outcome declared; deposit has only single-muscle columns with unlabelled .A/.P suffixes.
2015_a-12-week-exercise-program-for-pregnant-women-wi	data	outcome_timepoint_not_in_deposit	Primary outcome (after documented outcome switching) needs 36-week data; deposit has only v1/v2.

study_id	category	reason_code	reason
2026_changes-in-repetitive-negative-thinking-and-stre	no_estimate	secondary_analysis_no_estimate	Mediation analysis (SEM) of the ImPuls RCT; no own group-comparison estimate; primary report is elsewhere.

3 Table S2. Trial registry, version 2 (rule-based) against version 1 (ad hoc)

study_id	report_type	outcome_v2	outcome_v1	control_v2	control_v1	covariates_v2	covariates_v1	primary_declared
physiofeedback	secondary	AP_RMS	AP_RMS	Control	Control	Age,BMI,Gender,Race	Age,BMI,Gender,Race	secondary_report
mulligan	primary	NPRS2	NPRS2	3	1		Age,Gender	registration
lytras_fms	primary	NRPSLBPpost	NRPSLBPpost	2	1		Age,Gender,BMI	registration
clusterset	primary	1RM	CMJ_mean	TS	TS	Sex	Age,Sex	sample_size
tereco	primary	X_6MWD2	X_6MWD2	2	1			registration
spadi	primary	SPADI8wks	SPADItot4wks	3	1			registration
tscs	primary	BBS_Post	BBS_Post	2	1		Age,BMI	registration
tdcs	secondary	CAR_post_all	CAR_post_all	2	1		age,BMI	secondary_report
etip	secondary	Post_PGWB	EPDS_total	2	1		Pre_preg_BMI	secondary_report
phosphatidic	primary	LBM2	LBM2	2	1		Age,Sex	none
facial	primary	T3	VM2	2	1		Age	registration
gainingmore	primary	RF50Post	RF50Post	2	1		SEX	none
etre	primary	Weight	Weight	Control	Control			paper
creatine	primary	dxa_leanmass_kg_post_1	legs_post1	Control	Control		sex	registration
free	primary	rmdq_tot	rmdq_tot	Control	Control	agegp_w0		registration
mallorca	secondary	edimburgtotal	edimburgtotal	C	C			secondary_report

4 Table S3. Summary measures per trial and grid

grid_type: principled = estimand-preserving; strict = baseline handling fixed to analysis of covariance; defensible = Type E and N nodes; outlier_arm = sensitivity arm.

study_id	grid_type	n_spec	n	d_median	d_q25	d_q75	d_min	d_max	share_same_sign	s_median
clusterset	defensible	36	36	0.0644	0.0559	0.157	0.0528	0.179	1	1.49
clusterset	outlier_arm	18	36	0.0593	0.0542	0.155	0.0528	0.171	1	1.52
clusterset	principled	18	36	0.0593	0.0542	0.155	0.0528	0.171	1	1.52
clusterset	strict	6	36	0.0583	0.0528	0.0637	0.0528	0.0637	1	1.55
creatine	defensible	48	64	0.0431	0.0414	0.196	0.0163	0.28	1	3.18
creatine	outlier_arm	9	63	0.0446	0.0439	0.256	0.0439	0.256	1	5.16
creatine	principled	9	63	0.0446	0.0439	0.256	0.0439	0.256	1	5.16
creatine	strict	3	63	0.0439	0.0439	0.0439	0.0439	0.0439	1	5.14
etip	defensible	48	91	0.132	0.0802	0.185	-0.0161	0.254	0.938	0.98
etip	outlier_arm	30	59	0.227	0.192	0.275	0.0447	0.289	1	1.6
etip	principled	9	59	0.192	0.0447	0.254	0.0447	0.254	1	1.23
etip	strict	3	59	0.192	0.192	0.192	0.192	0.192	1	1.22

study_id	grid_type	n_spec	n	d_median	d_q25	d_q75	d_min	d_max	share_same_sign	s_median
etre	defensible	18	16	-0.916	-1.24	-0.887	-1.26	-0.874	1	9.41
etre	outlier_arm	9	16	-0.903	-1.24	-0.874	-1.24	-0.874	1	9.7
etre	principled	9	16	-0.903	-1.24	-0.874	-1.24	-0.874	1	9.7
etre	strict	3	16	-0.874	-0.874	-0.874	-0.874	-0.874	1	8.97
facial	defensible	30	24	1.51	1.45	1.55	1.35	1.6	1	9.53
facial	outlier_arm	16	23	1.49	1.45	1.51	1.44	1.55	1	9.57
facial	principled	9	23	1.49	1.45	1.55	1.45	1.55	1	9.64
facial	strict	3	23	1.49	1.49	1.49	1.49	1.49	1	10.6
free	defensible	96	226	0.092	0.0752	0.0977	-0.0605	0.137	0.948	0.79
free	outlier_arm	219	214	0.0549	0.0203	0.085	-0.0309	0.117	0.913	0.424
free	principled	35	214	0.0911	0.0753	0.0952	0.0562	0.117	1	0.797
free	strict	12	214	0.0911	0.0847	0.0936	0.0753	0.105	1	0.886
gainingmore	defensible	18	39	-0.102	-0.468	-0.06	-0.499	-0.0423	1	1.07
gainingmore	outlier_arm	9	39	-0.0901	-0.468	-0.06	-0.468	-0.06	1	0.956
gainingmore	principled	9	39	-0.0901	-0.468	-0.06	-0.468	-0.06	1	0.956
gainingmore	strict	3	39	-0.0901	-0.0901	-0.0901	-0.0901	-0.0901	1	0.956
lytras_fms	defensible	9	40	-0.884	-1.01	-0.785	-1.01	-0.785	1	7.35
lytras_fms	outlier_arm	9	40	-0.884	-1.01	-0.785	-1.01	-0.785	1	7.35
lytras_fms	principled	9	40	-0.884	-1.01	-0.785	-1.01	-0.785	1	7.35
lytras_fms	strict	3	40	-0.884	-0.884	-0.884	-0.884	-0.884	1	7.7
mallorca	defensible	10	294	-0.301	-0.313	-0.295	-0.315	-0.281	1	5.82
mallorca	outlier_arm	19	263	-0.287	-0.304	-0.244	-0.315	-0.236	1	6.36
mallorca	principled	3	263	-0.315	-0.315	-0.315	-0.315	-0.315	1	6.48
mallorca	strict	3	263	-0.315	-0.315	-0.315	-0.315	-0.315	1	6.48
mulligan	defensible	18	29	-0.325	-0.329	-0.22	-0.411	-0.117	1	1.52
mulligan	outlier_arm	9	29	-0.22	-0.32	-0.117	-0.32	-0.117	1	1.03
mulligan	principled	9	29	-0.22	-0.32	-0.117	-0.32	-0.117	1	1.03
mulligan	strict	3	29	-0.22	-0.22	-0.22	-0.22	-0.22	1	1.03
phosphatidic	defensible	18	18	0.101	0.0637	0.105	0.0562	0.106	1	4.01
phosphatidic	outlier_arm	9	18	0.105	0.0637	0.106	0.0637	0.106	1	3.83
phosphatidic	principled	9	18	0.105	0.0637	0.106	0.0637	0.106	1	3.83
phosphatidic	strict	3	18	0.106	0.106	0.106	0.106	0.106	1	4.41
physiofeedback	defensible	45	276	0.0744	0.0493	0.108	-0.0194	0.141	0.956	0.762
physiofeedback	outlier_arm	157	276	0.0913	0.0704	0.108	0.0222	0.141	1	1.13
physiofeedback	principled	23	276	0.108	0.0666	0.133	0.0493	0.141	1	1.38
physiofeedback	strict	8	276	0.114	0.108	0.133	0.0899	0.133	1	1.51
spadi	defensible	24	72	-0.952	-0.975	-0.91	-1	-0.869	1	14
spadi	outlier_arm	9	71	-0.975	-1	-0.904	-1	-0.904	1	12.3
spadi	principled	9	71	-0.975	-1	-0.904	-1	-0.904	1	12.3
spadi	strict	3	71	-0.975	-0.975	-0.975	-0.975	-0.975	1	17.3
tdcs	defensible	18	20	1.41	1.39	1.43	1.38	1.46	1	7.3
tdcs	outlier_arm	41	20	1.38	1.22	1.41	0.384	1.46	1	7.94
tdcs	principled	9	20	1.42	1.41	1.46	1.41	1.46	1	7.47
tdcs	strict	3	20	1.42	1.42	1.42	1.42	1.42	1	7.59
tereco	defensible	96	119	0.758	0.655	0.808	0.537	0.917	1	16.4
tereco	outlier_arm	126	112	0.826	0.801	0.858	0.762	0.913	1	19.1
tereco	principled	18	112	0.843	0.808	0.901	0.793	0.913	1	17.9
tereco	strict	6	112	0.843	0.834	0.851	0.834	0.851	1	24.3
tscs	defensible	18	20	1.09	0.976	1.36	0.975	1.5	1	10.8

study_id	grid_type	n_spec	n	d_median	d_q25	d_q75	d_min	d_max	share_same_sign	s_median
tscs	outlier_arm	9	20	1.03	0.975	1.36	0.975	1.36	1	12.3
tscs	principled	9	20	1.03	0.975	1.36	0.975	1.36	1	12.3
tscs	strict	3	20	0.975	0.975	0.975	0.975	0.975	1	17.3

4.1 Table S3b. Wilson intervals for proportions and study-bootstrap intervals for medians

quantity	k	n	prop	wilson_lo	wilson_hi
reproduced full	11	12	0.917	0.646	0.985
reproduced full or minor	11	12	0.917	0.646	0.985
published inside principled range	9	11	0.818	0.523	0.949
published inside defensible range	11	12	0.917	0.646	0.985
sign stable principled	16	16	1	0.806	1
sign stable defensible	13	16	0.812	0.57	0.934
range includes 0 principled	0	16	0	0	0.194
spec fully representable	7	12	0.583	0.32	0.807
reproduced full or minor, denominator 48 assessed RCTs	11	48	0.229	0.133	0.365
reproduced full or minor, denominator 152 with link included, denominator 48	11	152	0.0724	0.0409	0.125
included, denominator 152	16	48	0.333	0.217	0.475
	16	152	0.105	0.0658	0.164

quantity	est	boot_lo	boot_hi
median range d, principled	0.119	0.0917	0.212
median range d, defensible	0.239	0.132	0.322
median ratio defensible/principled (finite only)	1.35	1.12	1.52
median /published d - multiverse median d/ (principled)	0.0156	0.00272	0.0564

5 Table S4. Variance partition with two-way interactions, trial-by-grid combinations with at least 30 specifications

study_id	grid_type	n_spec	n_nodes	top_additive	share_additive	top_interaction_model	share_main_in_interactions	share_all_interactions
free	principled	35	4	covars	55.3	covars	42.7	17.7
physiofeedback	defensible	45	5	transform	43.7	transform	38.2	9.31
clusterset	defensible	36	4	baseline	98.2	baseline	97.4	0.848
tereco	defensible	96	5	transform	46.6	transform	45.5	2.42
etip	defensible	48	4	transform	75.7	transform	44.9	40.7
facial	defensible	30	4	baseline	98.8	baseline	79.8	19.2
creatine	defensible	48	4	baseline	97.9	baseline	95.4	2.48

study_id	grid_type	n_spec	n_nodes	top_additive	share_additive	top_interaction_model	share_main_in_interactions	share_all_interactions
free	defensible	96	5	baseline	49.9	baseline	12.5	74.6

6 Table S5. Range drivers per trial and range with baseline handling fixed

study_id	n	baseline_imbalance_sd	cor_base_outcome	frac_missing	range_p	range_d
physiofeedback	276	0.0411	0.466	0	0.0917	0.16
mulligan	29	0.162	0.632	0	0.203	0.294
lytras_fms	40	0.217	0.379	0	0.229	0.229
clusterset	36	0.105	0.985	0	0.118	0.126
tereco	119	0.165	0.613	0.0588	0.121	0.38
spadi	72	0.108	0.486	0.0139	0.0975	0.132
tscs	20	0.409	0.882	0	0.386	0.522
tdcs	20	0.0711	0.19	0	0.0569	0.0866
etip	91	0.038	0.549	0.352	0.209	0.27
phosphatidic	18	0.0434	0.994	0	0.0419	0.0495
facial	24	0.101	0.502	0.0417	0.101	0.249
gainingmore	39	0.411	0.915	0	0.407	0.457
etre	16	0.398	0.873	0	0.366	0.386
creatine	64	0.239	0.997	0.0156	0.212	0.263
free	226	0.0237	0.227	0.0531	0.0607	0.198
mallorca	294	NA	NA	0.105	0	0.034

study_id	range_full	n_spec	range_fixed
clusterset	0.0159	6	0.0159
creatine	0.212	3	0
etip	0.244	12	0.0372
etre	0	3	0
facial	0.11	6	0.00598
free	0.148	48	0.109
gainingmore	0.407	3	0
lytras_fms	0.229	3	0
mallorca	0.0795	12	0.0795
mulligan	0.203	3	0
phosphatidic	0.0419	3	0
physiofeedback	0.0347	26	0.0347
spadi	0.0975	3	0
tdcs	1.08	9	1.03
tereco	0.152	24	0.0543
tscs	0.386	3	0

7 Table S6. Data integrity screen

p_carlisle_named: Stouffer-combined p over baseline variables named in the paper (NA where fewer than two). p_carlisle_all: name-based broad search, reported only.

study_id	n	n_baseline_named	p_carlisle_named	n_baseline_all	p_carlisle_all	frac_outcome_nonint	n_dup_rows	p_lastdigit	flag
physiofeedback	358	2	0.308	2	0.308	1	0	0.918	pass
mulligan	29	1	NA	2	0.51	0	0	NA	pass
lytras_fms	40	1	NA	5	0.873	0	0	NA	pass
clusterset	36	0	NA	8	0.333	0.194	0	NA	pass
tereco	119	1	NA	7	0.672	0	0	NA	pass
spadi	72	1	NA	5	0.859	0	0	NA	pass
tscs	20	1	NA	10	0.145	0	0	NA	pass
tdcs	20	1	NA	13	0.953	0.9	0	NA	pass
etip	91	1	NA	43	0.324	0	0	NA	pass
phosphatidic	18	1	NA	5	0.449	1	0	NA	pass
facial	24	1	NA	2	0.911	0	0	NA	pass
gainingmore	39	1	NA	9	0.573	0.974	0	0.491	pass
etre	16	0	NA	0	NA	1	0	NA	pass
creatine	64	1	NA	25	0.269	1	0	0.312	pass
free	226	1	NA	21	0.892	0	0	NA	pass
mallorca	294	0	NA	16	0.263	0	0	NA	pass

8 Table S7. Effect of the first mapping (registry v1) on the results, computed with the final engine

study_id	v1_med	v1_range	v1_sign	v2_med	v2_range	v2_sign	sign_flip
physiofeedback	0.0498	0.0173	1	0.108	0.0917	1	FALSE
mulligan	-0.218	0.193	1	-0.22	0.203	1	FALSE
lytras_fms	0.886	0.229	1	-0.884	0.229	1	TRUE
clusterset	0.376	0.0114	1	0.0593	0.118	1	FALSE
tereco	-0.851	0.105	1	0.843	0.121	1	TRUE
spadi	-0.0437	0.169	0.667	-0.975	0.0975	1	FALSE
tscs	-1	0.476	1	1.03	0.386	1	TRUE
tdcs	-1.46	0.0693	1	1.42	0.0569	1	TRUE
etip	0.191	0.0668	1	0.192	0.209	1	FALSE
phosphatidic	-0.105	0.0419	1	0.105	0.0419	1	TRUE
facial	-0.567	0.252	1	1.49	0.101	1	TRUE
gainingmore	0.075	0.448	1	-0.0901	0.407	1	TRUE
etre	-1.24	0	1	-0.903	0.366	1	FALSE
creatine	0.00914	0.207	1	0.0446	0.212	1	FALSE
free	0.0911	0.047	1	0.0911	0.0607	1	FALSE
mallorca	-0.315	0	1	-0.315	0	1	FALSE

9 Table S8. Displacement of the published result per trial and grid

study_id	grid_type	n_spec	d_published	delta_median	delta_min	delta_max	abs_delta_max	share_within_0.1	share_within_0.2	share_same_sign_as_published	universal
creatine	principled	9	0.0439	0.000689	-	0.212	0.212	0.667	0.667	1	FALSE
creatine	defensible	24	0.0439	-	-0.0227	0.236	0.236	0.667	0.667	1	FALSE
				0.0000000000000000368	0.0000000000000000368	0.0000000000000000368	0.0000000000000000368	0.0000000000000000368	0.0000000000000000368	0.0000000000000000368	
etip	principled	9	0.192	0.000000000000000006647	0.062	0.147	0.147	0.667	1	1	FALSE
etip	defensible	24	0.192	-0.00408	-0.147	0.062	0.147	0.875	1	1	FALSE
etre	principled	9	-0.903	0.0000000000000000330.337	0.0295	0.337	0.337	0.667	0.667	1	FALSE
etre	defensible	9	-0.903	0.0000000000000000330.337	0.0295	0.337	0.337	0.667	0.667	1	FALSE
gainingmore	principled	9	-0.0628	-0.0273	-0.405	0.00274	0.405	0.667	0.667	1	FALSE
gainingmore	defensible	9	-0.0628	-0.0273	-0.405	0.00274	0.405	0.667	0.667	1	FALSE
lytras_fms	principled	9	-0.785	-0.0982	-0.229	0.0000000000000000007829	0.7829	0.667	0.667	1	FALSE
lytras_fms	defensible	9	-0.785	-0.0982	-0.229	0.00000000000000000007829	0.7829	0.667	0.667	1	FALSE
mallorca	principled	3	-0.316	0.00116	0.00116	0.00116	0.00116	1	1	1	FALSE
mallorca	defensible	5	-0.316	0.00116	0.00116	0.00966	0.00966	1	1	1	FALSE
physiofeedback	defensible	22	0.0526	0.0107	-0.072	0.0291	0.072	1	1	0.909	TRUE
spadi	principled	9	-0.913	-0.0619	-0.0882	0.00933	0.0882	1	1	1	FALSE
spadi	defensible	24	-0.913	-0.0391	-0.0882	0.0437	0.0882	1	1	1	FALSE
tdcs	principled	9	1.41	0.0183	0.0000000000000000000056569	0.0569	0.0569	1	1	1	FALSE
tdcs	defensible	9	1.41	0.0183	0.000000000000000000000056569	0.0569	0.0569	1	1	1	FALSE
tereco	principled	18	0.838	0.00481	-0.0451	0.0754	0.0754	1	1	1	FALSE
tereco	defensible	48	0.838	-0.0299	-0.152	0.0792	0.152	0.792	1	1	FALSE
tscs	principled	9	0.975	0.055	0.000000000000000000005386	0.386	0.386	0.667	0.667	1	FALSE
tscs	defensible	9	0.975	0.055	0.00000000000000000000005386	0.386	0.386	0.667	0.667	1	FALSE
free	principled	35	0.139	-0.048	-0.0829	-0.0222	0.0829	1	1	1	FALSE
free	defensible	96	0.139	-0.0471	-0.2	-0.0017	0.2	0.938	1	0.948	TRUE

10 Table S9. Published models as fitted, with assumptions recorded

study_id	model_as_published	est	ci_lo	ci_hi	p	n	deviations
creatine	Linear regression (ANCOVA): change in whole-body LBM (T1->T2, 7-day wash-in) ~ group + baseline LBM (T1), complete-case (Sec. 2.3.5)	0.507	0.0569	0.958	0.0279	63	REPRODUCTION VERY CLOSE TO THE PUBLISHED VALUE (0.507 vs 0.51 kg, p=0.028 vs 0.03). The paper reports only a point estimate + p (0.51 kg, p=0.03), no CI/SE for the group difference -> our CI is model-based (t-distributio

study_id	model_as_published	est	ci_lo	ci_hi	p	n	deviations
etip	General linear model ANCOVA: PGWBI global score (late pregnancy) ~ Group + PGWBI baseline, available ITT cases (Statistical methods)	2.6	-3.77	8.97	0.417	54	REPRODUCTION VERY CLOSE TO THE PUBLISHED VALUE (estimate 2.597 vs 2.60; 95% CI [-3.77, 8.97] vs [-3.77, 8.97] in the paper; p=0.417 vs 0.42). The paper does not formally declare PGWBI as the 'primary' outcome (see regist
etre	LMM: Weight ~ Group*Time + (1/ID), REML; main estimand = Time x Group interaction (change-score contrast eTRE vs Control, per override in eTRE-minus-Control con	-2.13	-3.07	-1.18	0.000241	24	The interaction coefficient (eTRE:post) is computed directly in our eTRE-minus-Control convention and matches the sign convention in config/published_overrides.csv (published value there: -2.13 [-3.07,-1.18], p=0.001; re
gainingmore	Bayesian multilevel model (brms, gaussian family, weakly informative brms default priors): RF50 ~ Group*Time + (1/CODE); group difference-in-change = Group x Ti	-0.515	-2.73	1.76	NA	39	The paper names 'univariate multilevel regression models' within a fully Bayesian framework (presumably brms/Stan), but specifies neither the prior distributions nor the exact random-effects structure of the 'multilevel'
lytras_fms	Two-way mixed ANOVA (Group x Time) on NPRS-Lumbar; primary interaction test, followed by a Bonferroni-adjusted simple-effect t-test (group comparison at week 3,	-1.2	-2.18	-0.222	0.0175	40	Unadjusted simple-effect t-test at week 3 (Bonferroni family not defined in the paper); Group x Time interaction from the mixed ANOVA reported in the console output, not adjusted, because the SPSS procedure for 'Bonferro

study_id	model_as_published	est	ci_lo	ci_hi	p	n	deviations
mallorca	ANOVA/unadjusted group comparison (EPDS total score, 1 month postpartum) with multiple imputation (mice, PMM, m=20) for missing EPDS values, pooled by Rubin's r	-0.694	-1.23	-0.161	0.0109	294	The SPSS MI model and the exact number/selection of auxiliary variables of the original paper are not documented (only 'Multiple imputation . . . with chained equations' is stated, without m or a list of predictors). Appro
physiofeedback	LMM (original authors' code from deposit/statistical code.R): $\log(AP_RMS) \sim \text{Group} * \text{Time} + \text{Age} + \text{BMI} + \text{Gender} + \text{Race} + (1/\text{SubjectID}) + (1/\text{Site})$, REML, bobyqa optimiser, R	0.021	-0.0782	0.12	0.678	373	The deposited original code writes 'Group * Time' without an explicit factor(Time); since Time is coded numerically (1-4) in the dataset, a purely linear treatment would yield only ONE interaction term, whereas Table 3 o
spadi	Repeated-measures ANCOVA as a GLM equivalent: ANCOVA on the mean of weeks 4 and 8 ~ Group + SPADI baseline, listwise exclusion; corresponds to the between-subje	-20.8	-28.9	-12.7	0.00000255	70	The paper presumably implements the 'repeated measures ANCOVA' as an SPSS GLM (multivariate approach with a fixed repeated-measures factor, no explicit random-effects model); approximated here instead with a linear mixed
tdcs	Independent t-test (equal variances): CAR post-intervention, active a-tDCS vs sham a-tDCS (Statistical analyses)	6.85	2.27	11.4	0.00563	20	The paper names Shapiro-Wilk/Levene pre-tests to choose between a t-test (equal variances) and Mann-Whitney U, but does not specify which test was actually used for CAR; computed here directly as Student's t-test with eq

study_id	model_as_published	est	ci_lo	ci_hi	p	n	deviations
tereco	Constrained longitudinal data analysis (LMM with equality constraint on baseline means): 6MWD ~ Time + Treatment (post/follow-up only) + Centre (fixed) + (1/Pat	65.4	43.8	87.1	0.00000000913	119	The paper typically describes cLDA as a GLS/mixed model with an UNSTRUCTURED covariance matrix across the 3 timepoints; a random-intercept model (compound-symmetry assumption) is used here instead, which is usually simil
tscs	ANCOVA: BBS post-intervention ~ Group + BBS baseline; contrast RAGT+tSCS vs CPT+tSCS (Statistical analysis {20a})	7.67	5.14	10.2	0.00000641	20	Marked group imbalance (n=13 vs n=7) as intended per the protocol, but unusual for 1:1 randomisation. The numeric group coding in the .sav dataset already contained text labels ('RAGT+tSCS'/'CPT+tSCS'), which were taken
free	LMM: RMDQ ~ baseline RMDQ + week*Tment + pre-specified covariates + (1+week/GP), REML; contrast FREE vs Control at week=26 (Statistical analysis, p.7)	0.58	-0.7	1.86	0.367	221	Covariate set assembled from the baseline variables available in the deposit that are already mean-imputed (age_w0, gender_w0, NZDep_Decile, d_length_back_pain_weeks, back-pain_consistency_3f, previous_backpain_w0_imp, ps

11 Table S10. Test statistics recomputed for the four papers without a point estimate

study_id	statistic	published	recomputed	note
mulligan	Group x Time F, p (repeated-measures ANOVA, 3 arms, n = 43)	F = 0.983, p = 0.383	F(2,40) = 0.983, p = 0.383	SPSS repeated-measures ANOVA; with two occasions the univariate and multivariate solutions coincide
mulligan mulligan	Group main effect F, p within-group means pre -> post	not reported as a number Mulligan 4.79 -> 2.36; Maitland 5.07 -> 2.14; Control ?	F(2,40) = 0.290, p = 0.750 Mulligan 4.79 -> 2.36 (n=14); Maitland 5.07 -> 2.14 (n=14); Control 5.07 -> 2.80 (n=15)	


study_id	statistic	published	recomputed	note
facial	Mann-Whitney U p (SFGS composite change, day 20 - baseline)	p = 0.002	p = 0.0020 (W = 114)	exact test, as SPSS reports for small samples; the asymptotic test with continuity correction gives p = 0.0033
facial	median (IQR) change per group	Exp 59.0 (42.5-63.0); Conv 24.0 (9.0-32.0)	Experimental 59.0 (42.5-63.0), n=12; Conventional 24.0 (9.0-32.0), n=11	
clusterset	Group x Time interaction p (LMM, Type III, Satterthwaite)	p = 0.352	F(1,34.0) = 0.890, p = 0.352	as described in the paper, Methods 2.4.2
clusterset	Time main effect F, p	F(1,34) = 80.67, p < 0.001	F(1,34.0) = 80.67, p = 1.7e-10	
clusterset	Sex main effect F, p	F(1,33) = 95.13, p < 0.001	F(1,33.0) = 95.13, p = 3e-11	
phosphatidic	Group x Time interaction F, p (2x2 mixed ANOVA, n = 18)	F(1,16) = 33.30, p = 0.041 (Results); 'main effect F(1,16) = 33.30, p < 0.001' (Abstract)	F(1,16) = 4.95, p = 0.0408	internally inconsistent in the paper: F = 33.30 with df (1,16) implies p < 0.001, whereas p = 0.041 implies F = 4.9
phosphatidic	Time main effect F, p	not separately reported	F(1,16) = 30.89, p = 4.3e-05	
phosphatidic	means pre/post per group	MT 60.8+/-9.5 -> 62.7+/-10.2; PLA 61.2+/-9.7 -> 62.0+/-9.7	MT 60.8+/-9.6 -> 62.7+/-10.2 (n=8); PLA 61.2+/-9.6 -> 62.0+/-9.7 (n=10)	

12 Table S11. Trial registrations retrieved


study_id	registration_id	retrieved	registered_after_start	registered_primary_outcome	matches_our_outcome_var
physiofeedback	NCT05778604	yes	no	Change from Baseline in Fall Risk Reduction; Change from Baseline in Dynamic Balance as Measured By Timed-Up and Go (TUG) test; Change from	no
mulligan	NCT06858124	yes	unclear	Changes in neck pain intensity with NPRS (listed first); Changes in Pressure Pain Threshold (PPT); Changes in cervical ROM; Changes in Neck	yes
lytras_fms	NCT07234071	yes	yes	Changes in pain intensity with NPRS (listed first); Straight Leg Raise (SLR) Angle - Goniometry/SROM; Change in Functional Disability (RMDQ)	yes
clusterset		no			
tereco	ChiCTR2000031834	yes	no	Total distances of 6 minutes walk test (Primary indicator; sole primary outcome)	yes
spadi	NCT01570985	yes	yes	Shoulder pain and disability index (SPADI) (sole primary outcome)	yes
tscs	ChiCTR2300074090	yes	no	Balance (Berg Balance Scale / Timed Up and Go) - listed first; Mobility (Walking Index for Spinal Cord Injury WISCI II / 10-Meter Walk Test)	yes

study_id	registration_id	retrieved	registered_after_start	registered_primary_outcome	matches_our_outcome_var
tdcs	IRCT20231113060048N1	yes	no	Quadriceps muscle strength (Biodex Isokinetic Dynamometer) - the ONLY registered primary outcome. 'Central activation ratio' (CAR) is part o	no
etip	NCT01243554	yes	unclear	Weight gain during pregnancy (kg), measured from 14 weeks' gestation to birth (sole primary outcome, unchanged since version 1)	no
phosphatidic facial	CTRI/2020/02/023240	no yes	no	Sunny Brook Facial Grading System (SFGS) - listed first; House Brackmann Scale; Facial Disability Index (3 jointly registered primary outcom	yes
gainingmore	OSF preregistration https://osf.io/bztka	yes	no	No separate 'primary outcome' field in this OSF pre-registration template (open Van't Veer/Giner-Sorolla template with no primary/secondary	partial
etre creatine	ACTRN12622000040763	no yes	no	[1] Whole-body lean mass measured by dual x-ray absorptiometry (DXA) - listed first; [2] Faith's phylogenetic diversity of the gut microbiot	yes
free	ACTRN12616000888460	yes	no	Patient back pain related impairment measured with the Roland Morris Disability Questionnaire (sole primary outcome)	yes
mallorca	ISRCTN14097513	yes	yes	The incidence of epidural analgesia use during labour is determined through review of the clinical history at one month after birth (sole pr	no


13 Figure S1. Median and range of d per trial, both grids




14 Figure S2. Share of interval-width variance per node





15 Figure S3. Specification curves, clusterset_PMC13140343




16 Figure S4. Specification curves, creatine_PMC11944689




17 Figure S5. Specification curves, etip_PMC6886967





18 Figure S6. Specification curves, etre_PMC11945196



19 Figure S7. Specification curves, facial_PMC11164989




20 Figure S8. Specification curves, free_PMC6733445




Band: 95% interval per specification. Dashed line: published estimate where reported.


21 Figure S9. Specification curves, gainingmore_PMC10809978




22 Figure S10. Specification curves, lytras_fms_PMC12942207




23 Figure S11. Specification curves, mallorca_PMC8198819




24 Figure S12. Specification curves, mulligan_PMC13092435





25 Figure S13. Specification curves, phosphatidic_PMC4891923



26 Figure S14. Specification curves, physiofeedback_PMC12821712





27 Figure S15. Specification curves, spadi_PMC4880881




Band: 95% interval per specification. Dashed line: published estimate where reported.

28 Figure S16. Specification curves, tdc_s_PMC13257960




Band: 95% interval per specification. Dashed line: published estimate where reported.

29 Figure S17. Specification curves, tereco_PMC8318721



30 Figure S18. Specification curves, tscs_PMC13085461



31 Text S2. Novelty search protocol, 15 September 2026

31.1 Novelty search, 2026-09-15

Statement under review (manuscript, body.tex line 5, or paper.md):

“I am not aware of a study in physiotherapy, rehabilitation or exercise science that retrieved individual participant data from public repositories and recomputed the published primary analysis.”

Criterion for a counterexample: individual participant data obtained from a public repository or supplement **AND** the published primary analysis recomputed, in the field of physiotherapy, rehabilitation or exercise and sports science.

This check supplements the earlier novelty search of 2026-08-28 (documented in README.md, section “Neuheitsprüfung”). The neighbours excluded there (Jabouille 2025, Elghzali 2025, Murphy 2025, Dhanani 2022) are not reviewed again in full text here but taken as already known.

Verdict in advance: no counterexample found. The statement remains tenable. The closest neighbour is a registered report study in exercise science, running since January 2026 and not yet completed, that does aim to recompute the published primary analysis but explicitly works with **private** raw data shared confidentially by the authors rather than data from a public repository. Details below.

31.2 Methodology and deviations from the search plan

All requests were run via curl with -A "Mozilla/5.0" and a 1.5 second pause between calls. Two deviations from the search specification, both documented:

1. **medRxiv/bioRxiv**: the direct HTML search on www.medrxiv.org/search/... returns only a Cloudflare challenge page (“Just a moment...”) and no search result. As foreseen in the search plan itself, OpenAlex with `filter=type:preprint` was used instead.
2. **OSF Registrations**: <https://api.osf.io/v2/registrations/> returned HTTP 502 or connection timeouts throughout the entire session (<https://api.osf.io/v2/> itself responded with 200, only the registrations endpoint was affected, with several retries over roughly two minutes and no success). As a substitute, the SHARE search API (https://share.osf.io/api/v2/search/creativeworks/_search) was used, which also indexes OSF registrations and was reachable at the time of the search (HTTP 200). This is a documented compromise, not a full substitute for `filter[title]`, because SHARE uses a different relevance weighting and also indexes further sources (CrossRef, DataCite, PubMed Central).

31.3 Queries verbatim, with source and hit count

31.4 Europe PMC (<https://www.ebi.ac.uk/europepmc/webservices/rest/search,resultType=lite,pageSize=100>)

#	Query	Hits
Q1	("computational reproducibility" OR "analytic reproducibility" OR "analytical reproducibility" OR "reanalysis" OR "re-analysis" OR "reproduction") AND ("physiotherapy" OR "physical therapy" OR "rehabilitation" OR "exercise science" OR "sports science" OR "sport science") AND ("individual participant data" OR "individual patient data" OR "raw data" OR "open data" OR "shared data" OR "repository")	5002
Q2	("multiverse") AND (Feldbegriffe wie Q1) AND (Datenbegriffe wie Q1)	28
Q3	("specification curve") AND (Feldbegriffe) AND (Datenbegriffe)	5

#	Query	Hits
Q4	("vibration of effects") AND (Feldbegriffe) AND (Datenbegriffe)	5
Q5	("many analysts") AND (Feldbegriffe) AND (Datenbegriffe)	5
Q6	("robustness reanalysis") AND (Feldbegriffe) AND (Datenbegriffe)	0
Q7	"reproducibility" AND "physical therapy" AND "data" AND (PUB_TYPE:"Meta-Analysis" OR "meta-research" OR "metaresearch")	256
Q8	"many analysts" AND ("sport" OR "sports science" OR "exercise science")	10
Q9	("reanalysis" OR "re-analysis") AND ("open data") AND ("sports science" OR "exercise science" OR "sport science" OR "Journal of Sports Sciences" OR "Medicine and Science in Sports and Exercise")	24
Q10	"prevalence" AND "reproducible" AND ("sport science" OR "sports science" OR "exercise science")	928
Q11	("individual patient data" OR "individual participant data") AND ("reanalysis" OR "re-analysis" OR "recomputed" OR "recompute") AND ("physiotherapy" OR "physical therapy" OR "rehabilitation") AND ("repository" OR "Zenodo" OR "Dryad" OR "OSF" OR "figshare")	49
Q12	("reproduced the published" OR "recomputed the published" OR "recompute the published" OR "reanalyzed the original data" OR "reanalysed the original data") AND ("physiotherapy" OR "physical therapy" OR "rehabilitation" OR "exercise" OR "sports")	120
Q13	"computational reproducibility" AND ("physiotherapy" OR "physical therapy" OR "rehabilitation" OR "exercise science" OR "sports science" OR "sport science")	29
Q14	"analytic reproducibility" AND ("physiotherapy" OR "physical therapy" OR "rehabilitation" OR "exercise" OR "sports" OR "sport")	14
Q15	("downloaded" OR "retrieved") AND ("individual participant data" OR "individual patient data" OR "raw data") AND ("recomputed" OR "recompute" OR "reran the analysis" OR "re-ran the analysis") AND ("physiotherapy" OR "rehabilitation" OR "exercise" OR "sports" OR "physical therapy")	51
Q16	("Zenodo" OR "Dryad" OR "figshare" OR "Open Science Framework") AND ("reanalysis" OR "re-analysis" OR "recomputed") AND ("physiotherapy" OR "rehabilitation" OR "physical therapy" OR "exercise science" OR "sports science")	250

#	Query	Hits
Q17	"prevalence" AND "computational reproducibility" AND ("sport" OR "exercise" OR "physiotherapy" OR "physical therapy" OR "rehabilitation")	40
Q18	"Cochrane" AND ("individual patient data" OR "individual participant data") AND ("repository" OR "Zenodo" OR "Dryad" OR "Open Science Framework" OR "figshare") AND ("physiotherapy" OR "rehabilitation" OR "physical therapy")	256
Q19	TITLE:("Reanalysis of" OR "Re-analysis of" OR "A reanalysis of" OR "Reanalyzing") AND ("physiotherapy" OR "rehabilitation" OR "physical therapy" OR "exercise" OR "sports" OR "sport")	175

For hit counts above 100, only the first 100 (sorted by relevance) were loaded and reviewed. This is a lower bound on completeness, but Europe PMC sorts by relevance to the query, so hits from the target field should appear within the first 100 if they exist.

31.5 OpenAlex (https://api.openalex.org/works?search=<text>&per-page=50, teils &filter=from_publication_date:2015-01-01)

#	Search text	Hits
OA1	reanalysis individual participant data	18
OA2	physiotherapy rehabilitation repository	0
OA3	multiverse specification curve physiotherapy rehabilitation exercise trial	577
OA4	computational reproducibility sport science exercise physiotherapy	29 165
OA5	many analysts sports science	2073
OA6	reanalysis open data sports science journal	51
OA7	individual patient data repository recompute rehabilitation trial	5915
OA8	vibration of effects physiotherapy rehabilitation exercise	104
OA9	robustness reanalysis physiotherapy exercise rehabilitation	17
OA10	Mesquida many analysts sport	320 (not individually reviewed, author already covered via OA9/Sports Med neighbour)
OA11	Caldwell many analysts sport science	4
OA12	Vigotsky reanalysis sport science open data	439 (not individually reviewed)
OA13	Borg many analysts sport	97 (not individually reviewed)
OA14	Bosco many analysts sport science	170 (not individually reviewed)
OA15	Abt many analysts sport science	16
OA16	Warmenhoven many analysts biomechanics	16
PP1	Cochrane individual patient data reanalysis physiotherapy repository reanalysis individual participant data physiotherapy rehabilitation exercise (filter=type:preprint)	24

#	Search text	Hits
PP2	multiverse specification curve rehabilitation physical therapy trial (filter=type:preprint)	0

For OA3, OA4, OA5 and OA7, only the relevance-sorted top 15 to top 20 were reviewed, because free-text search without phrase binding spreads very broadly (for example, “vibration of effects” pulls in every study on vibration training in physiotherapy). For OA10, OA12, OA13 and OA14, individual review was omitted for reasons of time, because the associated authors (Caldwell, Borg, Bosco, Abt) are already covered through co-authorship with Murphy et al. 2025 and through Mesquida’s work list (OA9), where no IPD reanalysis appears.

In addition, targeted OpenAlex author queries were run: Matthieu Boisgontier (author.id:A5076503553, published since 2026-08-01: 4 works, all Kinarm validation plus one meta-analysis data deposit, no reanalysis), François Jabouille (author.id:A5084775748, 20 works, the two most recent are Kinarm validation studies and a French-language transparency checklist article from 2026-08-01, no reanalysis), Cristian Mesquida (author.id:A5082605089, 36 works, most recent are the software package metacheck, a type S/M error comment, and a power analysis scoping review, no IPD reanalysis), and Colby J. Vorland (author.id:A5074473858, 115 works, filter search=reanalysis: 16 hits, see candidate table).

31.6 medRxiv/bioRxiv

Direct access via www.medrxiv.org/search/... is blocked by Cloudflare (challenge page, no result extractable). Substitute: OpenAlex filter=type:preprint (see PP1, PP2 above).

31.7 OSF Registrations

[https://api.osf.io/v2/registrations/?filter\[title\]=<term>](https://api.osf.io/v2/registrations/?filter[title]=<term>) returned HTTP 502 or a timeout throughout, for all seven planned search terms (“reproducibility physiotherapy”, “reanalysis rehabilitation”, “multiverse exercise”, “reanalysis physiotherapy”, “multiverse physiotherapy”, “individual participant data physiotherapy”, “computational reproducibility sport”), even after several retries spread over two minutes. Substitute: SHARE search API.

#	SHARE query (q=)	Total hits
S1	reproducibility physiotherapy (Freitext, ungebunden)	50 851 (too broad to evaluate, see S2/S3)
S2	title:("reanalysis" OR "re-analysis" OR "multiverse" OR "individual participant data" OR "individual patient data") AND title:(physiotherapy OR rehabilitation OR "exercise science" OR "sports science" OR "sport science" OR "physical therapy")	21
S3	title:("multiverse" OR "specification curve" OR "computational reproducibility" OR "vibration of effects") AND title:(physiotherapy OR rehabilitation OR "exercise" OR "sports" OR "sport" OR "physical therapy")	2

S2 and S3 are queries restricted to the title (Elasticsearch query_string syntax), because pure free-text search (S1) was no longer meaningfully reviewable by hand, given the enormous SHARE register enriched with many external sources (ClinicalTrials.gov, DataCite, CrossRef).

31.8 Candidates reviewed, with verdict

Title	Authors	Year	Journal/Source	DOI	Verdict
Finding Predictive Factors of Stabilization Exercise Adherence in RCTs on Low Back Pain: An Individual Data Reanalysis Using Machine Learning Techniques	Pfeifer AC, Schröder-Pfeifer P, Schiltenswolf M, et al.	2025	Arch Phys Med Rehabil	10.1016/j.apmr.2024.12.015	Not a counterexample. A “preplanned reanalysis” by the same MiSpEx network that collected the original data, with no data obtained from a public repository. The research question concerns ML predictors of adherence, not the published primary analysis.
Dose-response relationship and effect modifier of stabilisation exercises in nonspecific low back pain: a project-wide individual patient data re-analysis on 1483 intervention participants	Niederer D, Pfeifer AC, Engel T, et al.	2023	Pain	10.1097/j.pain.0000000000002801	Not a counterexample. Same MiSpEx network, project-internal IPD not obtained from a public repository. A dose-response question rather than recomputation of a published primary analysis.
Re-analysis of data from a cluster RCT entitled “health literacy and exercise-focused interventions on clinical measurements in Chinese diabetes patients”	Jamshidi-Naeini Y, Golzarri-Arroyo L, Vorland CJ, Brown AW, Allison DB	2022	eClinicalMedicine	10.1016/j.eclinm.2022.101686	Not a counterexample, but methodologically the closest to our approach. The authors first explicitly reproduce the published figures, then recompute with a corrected model (LMM instead of GEE). The data source, however, is direct, confidential sharing by the original authors (“collegially shared”), not a public repository, and the field is diabetes care and health literacy, not physiotherapy, rehabilitation or exercise science in the narrower sense.
Contrary to the Conclusions Stated in the Paper, Only Dry Fat-Free Mass Was Different between Groups upon Reanalysis. Comment on “Intermittent Energy Restriction. . .”	Peos J, Brown AW, Vorland CJ, Allison DB, Sainsbury A	2020	J Funct Morphol Kinesiol	10.3390/jfmk5040085	Not a counterexample. Individual data come from the online supplement of the original publication (hence publicly accessible), field is exercise science and strength training. However, no recomputation of the published primary analysis, but an alternative, corrected model (ANCOVA/ITT instead of a completers-only DINS comparison) to refute the conclusion. As a letter/comment, not an independent study contribution and no multiverse.
Individually randomized trial mislabeled as a cluster-randomized trial. Comment on “Effectiveness of wearable technology to optimize youth soccer players’ off-training behaviour. . .”	Vorland CJ, et al.	2023	Sci Med Football	10.1080/24733938.2023.2190998	Not a counterexample. A short comment on a randomisation/analysis error, with no discernible retrieval of individual data from a repository and no complete recomputation of the primary analysis.

Title	Authors	Year	Journal/Source	DOI	Verdict
Replication concerns in sports and exercise science: a narrative review of selected methodological issues in the field	Mesquida C, Murphy J, Lakens D, Warne J	2022	R Soc Open Sci	10.1098/rsos.220946	Not a counterexample. A narrative review, no original reanalysis.
Estimating the Replicability of Sports and Exercise Science Research (inkl. Korrektur 2025-09-11)	Murphy J, Caldwell A, Mesquida C, et al.	2025	Sports Med	10.1007/s40279-025-02201-w	Already documented as a neighbour (README, as of 2026-08-28). A replication with new data, not an IPD reanalysis from a repository. The September 2025 correction does not change this.
Reproducible candidate kinematic-electromyographic waveform markers of post-stroke gait from public multimodal waveform exports	Calabrò RS, Calderone A, Sottile F, et al.	2026	Front Med Technol	10.3389/fmedt.2026.1863908	Not a counterexample. A “secondary analysis” of a public gait dataset, but one that produces new markers/metrics rather than recomputing the published primary analysis of the original dataset.
Computational Reproducibility in Sports Science: A Registered Report Reanalyzing Private Raw Data	Nolte S	Registered 2026-01-14	OSF Registries	10.17605/OSF.IO/VCWP8	Not a counterexample, but the most important new neighbour. The aim is precisely to recompute the published primary analyses of 50 articles in the Journal of Sports Sciences, assessing correctness and methodological vagueness. The data source, however, is explicitly private : authors are asked to share their raw data confidentially, precisely because “relying only on published data may yield a distorted sample”. No published results yet (Europe PMC search for the title: 0 hits, as of 2026-09-15).
Data and Code Availability in Sports Science: A Registered Report	Nolte S, Memmert D, Rein R	Preprint 2026-09-08	OSF Preprints	10.31222/osf.io/et5fw_v1	Not a counterexample. A pure availability audit of all original articles in Q1 exercise science journals over the past ten years (like Jabouille/Elghzali), no reanalysis. Notable as a scooping signal: the same group (Cologne) is thereby building “a resource with all articles that have shared data and/or code for reuse”, whose obvious follow-on application would be a public-repository reanalysis.

Title	Authors	Year	Journal/Source	DOI	Verdict
Prevalence and predictors of data and code sharing in the medical and health sciences: systematic review with meta-analysis of individual participant data	(OpenAlex hit, not field specific)	2023	–	–	Not a counterexample. “Individual participant data” here refers to the individual studies evaluated in the meta-analysis itself (the proportion with data/code sharing per study as the “observation”), not to clinical individual-level data. The topic is the prevalence of data/code sharing generally in medicine, not a physiotherapy-related field.
CaReMATCH / ExTraMATCH II / Precision rehabilitation for aphasia (several IPD meta-analysis protocols)	various	2018-2023	OSF/cardiac rehab/stroke J	various	Not a counterexample. Classic prospective IPD meta-analyses across several studies, with individual data collected directly from the participating study teams (not drawn from a public repository). The target quantity is a pooled meta-analytic effect across studies, not the recomputation of a single published primary analysis.

31.9 Update on scooping risk (supplement to README, as of 2026-08-28)

The Boisgontier/Jabouille line has shown no new direction since 28 August. The most recent works are a Kinarm validation study (registered report, 2026-08-26/09-11) and a French-language transparency checklist article in *Kinésithérapie, la Revue* (2026-08-01, “Les indicateurs de la science ouverte comme estimation du risque de pratiques de recherche douteuses en kinésithérapie”). Both confirm the earlier assessment: meta-research at the article level, no methodological shift to individual data.

New and more relevant is **Simon Nolte** (German Sport University Cologne, with Daniel Memmert and Robert Rein). Since January 2026 he has been pursuing two parallel, complementary projects: a field-wide availability audit of data and code in exercise science journals (preprint 2026-09-08, one week before this search date) and a registered-report reanalysis of the published primary analyses of 50 articles in the *Journal of Sports Sciences* (registered 2026-01-14). The second project is content-wise the closest to this manuscript, but it deliberately uses raw data shared privately and confidentially by the authors rather than data from a public repository, and it covers exercise science journal-wide rather than physiotherapy/rehabilitation studies with deposited repository data. No multiverse, no specification curve. Should Nolte in future link his availability-audit resource (“resource with all articles that have shared data and/or code”) to a reanalysis, that would be a direct competitor. At the time of this search, that step does not exist.

31.10 Conclusion

Across 19 Europe PMC queries, 17 OpenAlex queries (2 of them with `type:preprint`) and 3 SHARE/OSF queries (after the official OSF API failed), no published or registered contribution was found that (a) obtains individual data from a public repository or supplement, (b) thereby recomputes the published primary analysis of a physiotherapy, rehabilitation or exercise/sports science study, and (c) does so as an independent study contribution. The statement in the manuscript remains tenable after this second, extended search.

The closest neighbours, in descending order of proximity, are: Nolte (2026, registered intent to reanalyse, but private rather than public data), the Jamshidi-Naeini/Allison group (2022, first reproduces the original figures, then a corrected reanalysis, but outside the field and with data shared directly rather than deposited publicly), the Vorland/Allison/Peos “comment” series (occasionally uses supplement individual data in exercise science journals, but as a short error correction without recomputing the published figure and without a multiverse), and the MiSpEx network’s own reanalyses (Pfeifer 2025, Niederer 2023: project-internal IPD, a different research question). None of these satisfies both criteria simultaneously within the target field.

Limitation: the OSF registrations API was unreachable on the day of the search. The SHARE substitute covers the same registration pool, but with a different relevance weighting. A further direct check via `filter[title]` is recommended once the API is working again, but given the overlap of the SHARE hits with the Europe PMC/OpenAlex results, this is unlikely to change the overall verdict.