

TABLE V Regimen in Recommendations for Prophylactic Antibiotics Systematically Given to Patients with Open Fractures of Extremities*										
Recommended Drug†	Injury Severity Not Specified or for All Levels	Gustilo Type			Extremity‡		Soil, Marine, or Severely Contaminated Wounds	Gunshot Velocity		
		I and II	II and III	III	Upper	Lower		Not Specified	Low	High
Gram-positive coverage§	13 (18.1%)	34 (68.0%)				9 (26.5%)	9 (22.0%)	1 (50.0%)	3 (50.0%)	6 (42.9%)
Gram-positive and anaerobic coverage#		2 (4.0%)								1 (7.1%)
Gram-positive and MRSA coverage**	1 (1.4%)						1 (2.4%)			
Gram-negative coverage	1 (1.4%)	2 (4.0%)		5 (10.9%)		1 (2.9%)				
Broad coverage††	14 (19.4%)	10 (20.0%)	6 (85.7%)	35 (76.1%)		9 (26.5%)	26 (63.4%)	1 (50.0%)	1 (16.7%)	5 (35.7%)
Broad and anaerobic coverage	3 (4.2%)	1 (2.0%)	1 (14.3%)	3 (6.5%)		1 (2.9%)	4 (9.8%)			1 (7.1%)
Broad and MRSA coverage				1 (2.2%)			1 (2.4%)			
Broad, MRSA, and anaerobic coverage	5 (6.9%)					3 (8.8%)				
Drug name not specified	35 (48.6%)	1 (2.0%)		2 (4.3%)	4 (100%)	11 (32.4%)			2 (33.3%)	1 (7.1%)
Total	72 (100%)	50 (100%)	7 (100%)	46 (100%)	4 (100%)	34 (100%)	41 (100%)	2 (100%)	6 (100%)	14 (100%)