

TABLE III Regimen in Practice for Prophylactic Antibiotics Systematically Given to Patients with Open Fractures of Extremities*									
Antibiotic Regimen in Practice†	Injury Severity Not Specified or for All Levels	Gustilo Type			Extremity‡		Soil, Marine, or Severely Contaminated Wounds	Gunshot Velocity	
		I and II	II and III	III	Upper	Lower		Low	High
Gram-positive coverage§	2 (16.7%)	3 (17.6%)		1 (5.0%)	2 (40.0%)	7 (18.4%)			
Gram-positive and anaerobic coverage#	1 (8.3%)	4 (23.5%)		2 (10.0%)			1 (50.0%)		
Gram-positive and MRSA coverage**							1 (50.0%)		1 (33.3%)
Gram-negative coverage††		3 (17.6%)		3 (15.0%)					
Gram-negative and anaerobic coverage						1 (2.6%)			
Broad coverage‡‡	1 (8.3%)	7 (41.2%)	2 (100%)	7 (35.0%)		12 (31.6%)			1 (33.3%)
Broad and anaerobic coverage				3 (15.0%)	1 (20.0%)	2 (5.3%)			
Broad and MRSA coverage	1 (8.3%)								
Broad, MRSA, and anaerobic coverage						2 (5.3%)		1 (100%)	
Drug name not specified	7 (58.3%)			4 (20.0%)	2 (40.0%)	14 (36.8%)			1 (33.3%)
Total	12 (100%)	17 (100%)	2 (100%)	20 (100%)	5 (100%)	38 (100%)	2 (100%)	1 (100%)	3 (100%)