

Cartesian Products Avoiding Patterns

by

Jacob Denson

BSc. Computing Science, University of Alberta, 2017

A THESIS SUBMITTED IN PARTIAL FULFILLMENT
OF THE REQUIREMENTS FOR THE DEGREE OF

Master of Science

in

THE FACULTY OF SCIENCE

(Mathematics)

The University of British Columbia

(Vancouver)

April 2019

© Jacob Denson, 2019

The following individuals certify that they have read, and recommend to the Faculty of Graduate and Postdoctoral Studies for acceptance, the thesis entitled:

Cartesian Products Avoiding Patterns

submitted by **Jacob Denson** in partial fulfillment of the requirements for the degree of **Master of Science** in **Mathematics**.

Examining Committee:

Malabika Pramanik, Mathematics

Supervisor

Joshua Zahl, Mathematics

Supervisor

Abstract

The pattern avoidance problem seeks to construct a set $X \subset \mathbf{R}^d$ with large dimension that avoids a prescribed pattern such as three term arithmetic progressions, or more general patterns such as avoiding points x_1, \dots, x_n such that $f(x_1, \dots, x_n) = 0$ (three term arithmetic progressions are specified by the pattern $x_1 - 2x_2 + x_3 = 0$). Previous work on the subject has considered patterns described by polynomials, or by functions f satisfying certain regularity conditions. We consider the case of ‘rough patterns’. There are several problems that fit into the framework of rough pattern avoidance. As a first application, if $Y \subset [0, 1]$ is a set with Minkowski dimension α , we construct a set $X \subset [0, 1]$ with Hausdorff dimension $1 - \alpha$ so that $X + X$ is disjoint from Y . As a second application, given a set Y of dimension close to one, we can construct a subset $X \subset Y$ of dimension $1/2$ that avoids isosceles triangles.

Lay Summary

The lay or public summary explains the key goals and contributions of the research/scholarly work in terms that can be understood by the general public. It must not exceed 150 words in length.

Preface

At University of British Columbia (UBC), a preface may be required. Be sure to check the Graduate and Postdoctoral Studies (GPS) guidelines as they may have specific content to be included.

Contents

Abstract	iii
Lay Summary	v
Preface	vii
List of Figures	xi
Glossary	xiii
Acknowledgments	xv
1 Introduction	1
2 Background	3
2.1 Fractal Dimension	5
2.2 Hyperdyadic Covers	17
2.3 Branching Processes	18
2.4 Equidistribution Results	18
2.5 Results about Hypergraphs	21
2.6 Hyperdyadic Covers	22
2.7 Dimensions of Cantor Like Sets	25
2.8 Basic Covering	26
2.9 An Epsilon of Room	26
2.10 Bounds on Uniform Constructions	27
3 Related Work	29
3.1 Rusza: Difference Sets Without Squares	29
3.2 Keleti: A Translate Avoiding Set	34

3.3	Fraser/Pramanik: Extending Keleti Translation to Smooth Configurations	37
3.4	A Set Avoiding All Functions With A Common Derivative	40
4	Avoiding Rough Sets	43
4.1	Frequently Used Notation and Terminology	45
4.2	Avoidance at Discrete Scales	46
4.3	Fractal Discretization	50
4.4	Dimension Bounds	53
4.5	Applications	55
4.6	Relation to Literature, and Future Work	63
5	Extensions to Low Rank Configurations	67
5.1	Boosting the Dimension of Pattern Avoiding Sets by Low Rank Coordinate Changes	67
5.2	Extension to Well Approximable Numbers	70
5.3	Equidistribution and Real Valued Matrices	71
5.4	Applications of Low Rank Coordinate Changes	72
5.5	Idea: Generalizing This Problem to low rank smooth functions . .	74
5.6	Idea: Algebraic Number Fields	75
5.7	A Scheme for Avoiding Configurations	75
5.8	Square Free Sets	76
6	Constructing Squarefree Sets	77
6.1	Ideas For New Work	77
6.2	Squarefree Sets Using Modulus Techniques	79
6.3	Idea; Delaying Swaps	82
6.4	Squarefree Subsets Using Interval Dissection Methods	82
6.5	Finding Many Startpoints of Small Modulus	84
6.6	A Better Approach	85
7	Conclusions	87
	Bibliography	89

List of Figures

Glossary

This glossary uses the handy `acroynym` package to automatically maintain the glossary. It uses the package's `printonlyused` option to include only those acronyms explicitly referenced in the \LaTeX source.

GPS Graduate and Postdoctoral Studies

Acknowledgments

Thank those people who helped you.

Don't forget your parents or loved ones.

You may wish to acknowledge your funding sources.

Chapter 1

Introduction

In this thesis, we study a simple question from the perspective of geometric measure theory:

How large can Euclidean sets be not containing geometric patterns?.

The patterns manifest as certain configurations of point tuples, often affine invariant. For instance, we might like to find large subsets X of \mathbf{R}^n that contain no three collinear points, or do not contain all of the vertices of any isosceles triangle. Aside from purely geometric interest, these problems provide useful scenarios in which to test the methods of ergodic theory and harmonic analysis.

Geometric measure theory becomes important because sets which avoid the patterns we consider are highly irregular. Any open subset of \mathbf{R}^n must contain a copy of the vertices of any suitably small isosceles triangle, and the Lebesgue density theorem shows that this remains true if we consider any subset of \mathbf{R}^n of positive measure. Any set avoiding any of the patterns we study in this thesis must necessarily have Lebesgue measure zero. This means we cannot use the Lebesgue measure as a measure of the ‘maximal size’ of a set avoiding patterns. A ‘second-order’ measure of a set’s size has to be introduced, and this is satisfied by the fractional dimension of a set, first introduced by Felix Hausdorff in 1918. We rephrase our problem as determining the maximum Hausdorff dimension a set can have avoiding patterns.

At the time of this writing, many fundamental questions about the relation between the geometric arrangement of a set and its fractional dimension remain open. It might be expected that sets with sufficiently large Hausdorff dimension contain patterns, but this is not always true. For example, Theorem 6.8 of [12] shows that any set $X \subset \mathbf{R}^d$ with Hausdorff dimension exceeding one must contain

three colinear points. On the other hand, [10] constructs a set $X \subset \mathbf{R}^d$ with full Hausdorff dimension such that no four points in X form the vertices of a parallelogram. On the other hand, Theorem 6.8 of [12] shows that any set $X \subset \mathbf{R}^d$ with Hausdorff dimension exceeding one must contain three colinear points. For most geometric configurations, it remains unknown at what threshold patterns are guaranteed, or whether such a threshold exists at all. So finding sets with large Hausdorff dimension avoiding patterns is an important topic to study in the study of contemporary geometric measure theory.

Thus we derive new methods for constructing sets avoiding patterns. Furthermore, rather than studying particular instances of the configuration avoidance problem, we choose to study general methods for finding large subsets of space avoiding any pattern. In particular, we expand on a number of general *pattern dissection methods* which have proven useful in the area, originally developed by Keleti but also studied notably by Mathé, and Pramanik/Fraser.

Our novel contribution to the construction is that in an arbitrary configuration problem, random choices can overcome the lack of presence of structure. We use this to expand the utility of interval dissection methods to a set of *fractal avoidance problems*, that previous methods were completely unavailable to address. Such problems include finding large X such that the angles formed by any three distinct points of X avoid a specified set Y of angles, where Y has a fixed Hausdorff dimension. For instance, Y could be the set of all angles expressible as $q + x$ radians, where q is rational, and x lies in the Cantor set.

Chapter 2

Background

This thesis discusses methods to form large sets avoiding patterns. First, we give a precise definition by what we mean by a pattern, and what it means to avoid a pattern. We consider an ambient set \mathbf{A} . It's *n-point configuration space* is the set of distinct tuples of n points in \mathbf{A} , i.e.

$$\mathcal{C}^n(\mathbf{A}) = \{(x_1, \dots, x_n) \in X^n : x_i \neq x_j \text{ if } i \neq j\}.$$

The general *configuration space* of \mathbf{A} is $\mathcal{C}(\mathbf{A}) = \bigcup_{n=1}^{\infty} \mathcal{C}^n(\mathbf{A})$. A *pattern*, or *configuration*, on \mathbf{A} is a subset of $\mathcal{C}(X)$, and we say a subset Y of X *avoids* a configuration \mathcal{C} if $\mathcal{C}(Y)$ is disjoint from \mathcal{C} . We say \mathcal{C} is an *n point configuration* if it is a subset of $\mathcal{C}^n(X)$.

Example (Isocoles Triangle Configuration). *Consider the problem of finding a set avoiding the vertices of an isocoles triangle in the plane. Set*

$$\mathcal{C} = \{(x_1, x_2, x_3) \in \mathcal{C}^3(\mathbf{R}^2) : |x_1 - x_2| = |x_1 - x_3|\}.$$

Then \mathcal{C} is a 3-point configuration, and a set $X \subset \mathbf{R}^2$ avoids \mathcal{C} if and only if it contains no vertices of an isocoles triangle. Notice that $|x_1 - x_2| = |x_1 - x_3|$ holds if and only if $|x_1 - x_2|^2 = |x_1 - x_3|^2$, which is an algebraic equation in the coordinates of x_1, x_2 , and x_3 . Thus \mathcal{C} is an algebraic hypersurface in \mathbf{R}^6 .

Example (General Position Configuration). *Suppose we wish to find a subset of \mathbf{R}^d such that every collection of $k+1$ points in the set, for $k < d$, lies in 'general position', i.e. they do not lie in a k dimensional hyperplane. Set*

$$\mathcal{C}^{k+1} = \{(x_0, x_1, \dots, x_k) \in \mathcal{C}^{k+1}(\mathbf{R}^d) : x_1 - x_0, \dots, x_k - x_0 \text{ are linearly dependant}\}.$$

and then consider the configuration $\mathcal{C} = \bigcup_{k=2}^d \mathcal{C}^k$. A set X avoids \mathcal{C} if and only if all of its points lie in general position. Notice that

$$\mathcal{C}^{k+1} = \bigcup \{ \text{span}(y_1, \dots, y_k) \times \{y\} : y = (y_1, \dots, y_k) \in \mathcal{C}^k(\mathbf{R}^d) \} \cap \mathcal{C}^{k+1}(\mathbf{R}^d).$$

so each \mathcal{C}^{k+1} is essentially a union of k dimensional hyperplanes.

Even though our problem formulation assumes configurations are formed by distinct sets of points, one can still formulate avoidance problems involving repeated points in our framework, because an instance of a configuration involving n points which may contain repetitions can be seen as an instance of a configuration involving fewer than n distinct points.

Example (Sum Set Configuration). Let G be an abelian group, and fix $Y \subset G$. Set

$$\mathcal{C}^1 = \{g \in \mathcal{C}^1(G) : 2g \in Y\} \quad \text{and} \quad \mathcal{C}^2 = \{(g_1, g_2) \in \mathcal{C}^2(G) : g_1 + g_2 \in Y\}.$$

Then set $\mathcal{C} = \mathcal{C}^1 \cup \mathcal{C}^2$. A set $X \subset G$ avoids \mathcal{C} if and only if $(X + X) \cap Y = \emptyset$.

Our main focus in this thesis is on the *pattern avoidance problem*: Given a configuration \mathcal{C} on \mathbf{A} , there are various way of measuring how large can $X \subset \mathbf{A}$ be avoiding \mathcal{C} :

- If \mathbf{A} is finite, the goal is to find X with large cardinality.
- If \mathbf{A} is a discrete limit of finite sets \mathbf{A}_n , the goal is to find X such that $X \cap \mathbf{A}_n$ has large cardinality asymptotically in n .
- If $\mathbf{A} = \mathbf{R}^d$, but \mathcal{C} is a sufficiently discrete configuration, then a satisfactory goal is to find X with large Lebesgue measure avoiding \mathcal{C} .

In this thesis, inspired by methods in the past three settings, we establish methods for avoiding non-discrete configurations \mathcal{C} , i.e. we study configurations \mathcal{C} for which $X^m \cap \mathcal{C}^m$ are dense in X^m . The next section shows that Lebesgue measure completely fails to measure the degree of success for a solution to the pattern avoidance problem in this setting, but provides an alternate measurement which does succeed, and which we use as a metric in the pattern avoidance problems we consider.

2.1 Fractal Dimension

The Lebesgue measure is not the correct measurement for how large a pattern-avoiding set for non-discrete patterns. This is because for most of these patterns, every pattern-avoiding set has measure zero. One intuition as to why this is true is that a set with positive Lebesgue measure behaves in many respects like an open set, and an open set certainly intersects a dense set somewhere. The rigorous instance of this phenomenon we use is the Lebesgue density theorem. Its proof takes us too far afield into differentiation theory, so we merely state the result without proof.

Theorem 1 (Lebesgue Density Theorem). *Let $E \subset \mathbf{R}^d$ have positive Lebesgue measure. Then for almost every point $x \in E$,*

$$\lim_{r \rightarrow 0} \frac{|E \cap B_r(x)|}{|B_r(x)|} = 1.$$

Under mild non-discreteness conditions, which are certainly satisfied by the example configurations given in the last section, no set with positive Lebesgue measure can avoid a configuration.

Theorem 2. *Let \mathcal{C} be an n -point configuration on \mathbf{R}^d such that*

- (A) Translation Invariance: *For any $b \in \mathbf{R}^d$, $\mathcal{C} + b \subset \mathcal{C}$.*
- (B) Non-Discreteness: *For any $\varepsilon > 0$, there is (c_1, \dots, c_n) in the configuration \mathcal{C} such that $\text{diam}\{c_1, \dots, c_n\} \leq \varepsilon$.*

Then no set with positive Lebesgue measure avoids \mathcal{C} .

Proof. Let $X \subset \mathbf{R}^d$ have positive Lebesgue measure. Applying the Lebesgue density theorem, we find a point $x_0 \in X$ such that

$$\lim_{r \rightarrow 0} \frac{|X \cap B_r(x_0)|}{|B_r(x_0)|} = 1.$$

Give $\varepsilon > 0$, we fix r_0 such that $|X \cap B_{r_0}(x_0)| \geq (1 - \varepsilon)|B_{r_0}(x)|$. Let $C = (c_1, \dots, c_n) \in \mathcal{C}$ be an instance of the configuration such that $\text{diam}\{c_1, \dots, c_n\} \leq \varepsilon r_0$. For each $p \in B_{r_0}(x_0)$, let $C(p) = (c_1(p), \dots, c_n(p)) \in \mathcal{C}$, where $c_i(p) = p + (c_i - c_1)$. Then a

union bound gives

$$\begin{aligned}
\left| \bigcup_{i=1}^n \{p \in B_{r_0/2}(x_0) : c_i(p) \notin X\} \right| &\leq \sum_{i=1}^n |B_{r_0/2}(x_0) \cap (X + (c_1 - c_i))^c| \\
&= \sum_{i=1}^n |B_{r_0/2}(x_0)| - |B_{r_0/2}(x_0 + c_i - c_1) \cap X| \\
&\leq n(|B_{r_0/2}(x_0)| - |B_{r_0(1/2-\varepsilon)}(x_0) \cap X|) \\
&\leq n[(r_0/2)^d - (1-\varepsilon)(1/2-\varepsilon)^d r_0^d] \\
&\leq \varepsilon n(2d+1)(r_0/2)^d.
\end{aligned}$$

If $\varepsilon < 1/n(2d+1)$, we conclude that

$$\begin{aligned}
\left| \bigcap_{i=1}^n \{p \in B_{r_0/2}(x_0) : c_i(p) \in X\} \right| &= |B_{r_0/2}(x_0)| - \left| \bigcup_{i=1}^n \{p \in B_{r_0/2}(x_0) : c_i(p) \notin X\} \right| \\
&\geq (r_0/2)^d - \varepsilon n(2d+1)(r_0/2)^d > 0.
\end{aligned}$$

In particular, the intersection is non-empty, and so there exists p such that $C(p) \in \mathcal{C}(X)$, and so X does not avoid \mathcal{C} . \square

Fortunately, we have a second order notion of size, which is able to distinguish between sets of measure zero, known as *fractional dimension*. Intuitively, the fractional dimension provides a measure of the local density of a set, and so we view a set which is more dense as ‘larger’, for the purposes of a pattern avoidance problem. There are two definitions of fractional dimension we use in this thesis: Minkowski dimension and Hausdorff dimension. The main difference between the two is that Minkowski dimension measures relative density at a single scale, whereas Hausdorff dimension measures relative density at various scales.

We begin by discussing the Minkowski dimension, which is the easiest of the two dimensions to define. Given a length l , we let

$$\mathcal{B}(l, \mathbf{R}^d) = \{[a_1, a_1 + l) \times \cdots \times [a_d, a_d + l) : a_k \in l \cdot \mathbf{Z}\}.$$

Furthermore, given an arbitrary set $E \subset \mathbf{R}^d$, we let

$$\mathcal{B}(l, E) = \{I \in \mathcal{B}(l, \mathbf{R}^d) : I \cap E \neq \emptyset\}.$$

If E is a bounded set in \mathbf{R}^d , then for any length l , $\mathcal{B}(l, E)$ is a finite collection of

intervals, and we define the *lower* and *upper* Minkowski dimension as

$$\underline{\dim}_{\mathbf{M}}(E) = \liminf_{l \rightarrow 0} \left(\frac{\#\mathcal{B}(l, E)}{\log(1/l)} \right) \quad \overline{\dim}_{\mathbf{M}}(E) = \limsup_{l \rightarrow 0} \left(\frac{\#\mathcal{B}(l, E)}{\log(1/l)} \right).$$

If $\overline{\dim}_{\mathbf{M}}(E) = \underline{\dim}_{\mathbf{M}}(E)$, then we refer to this common quantity as the *Minkowski dimension* of E , denote $\dim_{\mathbf{M}}(E)$. This means that if $\dim_{\mathbf{M}}(E) = \alpha$, then as $\delta \rightarrow 0$, $|E_\delta| = \delta^{d-\alpha+o(1)}$. In particular, every set with positive Lebesgue measure has Minkowski dimension d , so Minkowski dimension is really only interesting for sets with Lebesgue measure zero. We can extend the lower and upper Minkowski dimension to unbounded sets E by considering the supremum and infimum of the dimensions of all bounded subsets of E .

Example. *Let*

$$C = \left\{ \sum_{i=1}^{\infty} a_i/4^i : a_i \in \{0, 3\} \right\}$$

If $1/4^{N+1} \leq \delta \leq 1/4^N$, then

$$\left\{ \sum_{i=1}^{\infty} a_i/4^i : a_1, \dots, a_{N+1} \in \{0, 3\} \right\} \subset C_\delta \subset \left\{ \sum_{i=1}^{\infty} a_i/4^i : a_1, \dots, a_N \in \{0, 3\} \right\}$$

The former set has volume $2^{N+1}/4^{N+1} = 1/2^{N+1} \geq \delta^{1/2}$, whereas the latter set has volume $2^N/4^N = 1/2^N \leq (2\delta)^{1/2}$. Thus $\log|C_\delta| = \log \delta/2 + O(1)$, and so

$$\dim_{\mathbf{M}}(C) = \lim_{\delta \rightarrow 0} \left(1 - \frac{\log|C_\delta|}{\log \delta} \right) = 1 - 1/2 = 1/2$$

So C has Minkowski dimension $1/2$.

There are a few things to notice about this calculation. First, we performed an upper and lower bound on powers of $1/4^k$, and then used this to obtain bounds at all scales. This is a general phenomenon for measuring Minkowski dimension. If we fix M , and consider $1/M^{k+1} \leq \delta \leq 1/M^k$, then as $k \rightarrow \infty$, we find $\log \delta \sim \log(1/M^k)$, and combined with the fact that $C_{1/M^{k+1}} \subset C_\delta \subset C_{1/M^k}$, we find

$$\frac{|C_{1/M^k}|}{\log(1/M^k)} \sim \frac{|C_{1/M^k}|}{\log(\delta)} \leq \frac{|C_\delta|}{\log(\delta)} \leq \frac{|C_{1/M^{k+1}}|}{\log(\delta)} \sim \frac{|C_{1/M^{k+1}}|}{\log(1/M^{k+1})}$$

Taking $k \rightarrow \infty$, we conclude that for any set E and any integer M ,

$$\underline{\dim}_{\mathbf{M}}(E) = \liminf_{n \rightarrow \infty} \frac{|C_{1/M^k}|}{\log(1/M^k)} \quad \text{and} \quad \overline{\dim}_{\mathbf{M}}(E) = \limsup_{n \rightarrow \infty} \frac{|C_{1/M^k}|}{\log(1/M^k)}$$

Thus Minkowski dimension need only be measured at a set of lengths of *dyadic type*. The second point is that we understood the dimension of C via a *Cantor-type* decomposition. The set C can be understood as the limit of a family of sets which are simple unions of intervals. If we set

$$C_k = \left\{ \sum_{i=1}^{\infty} a_i / 4^i : a_1, \dots, a_{N+1} \in \{0, 3\} \right\}.$$

Then $\{C_k\}$ is a nested family of sets, with $C = \lim C_k$. If for an index k , we set

$$\mathcal{B}^d(l, \mathbf{R}^d) = \{[a_1, a_1 + l) \times \dots \times [a_d, a_d + l) : a_k \in l \cdot \mathbf{Z}\}.$$

Then C_k is a union of 2^k cubes in $\mathcal{B}^d(1/4^k, \mathbf{R}^d)$, and

$$\frac{\log(2^k)}{\log(4^k)} = \log_4(2) = 1/2.$$

For a set E , take $\mathcal{B}^d(l, E)$ to be the set of all cubes in $\mathcal{B}^d(l, \mathbf{R}^d)$ intersecting E , i.e.

$$\mathcal{B}^d(l, E) = \{I \in \mathcal{B}^d(l, \mathbf{R}^d) : I \cap E \neq \emptyset\}.$$

It is intuitive that $\bigcup \mathcal{B}^d(l, E)$ is essentially the same as the l thickening E_l , which leads to the following Lemma.

Lemma 1. *If E is a bounded set in \mathbf{R}^d and M is an integer, then*

$$\underline{\dim}_{\mathbf{M}}(E) = \liminf_{l \rightarrow 0} \frac{\#\mathcal{B}^d(l, E)}{\log(1/l)} \quad \text{and} \quad \overline{\dim}_{\mathbf{M}}(E) = \limsup_{l \rightarrow 0} \frac{\#\mathcal{B}^d(l, E)}{\log(1/l)},$$

where the limit is taken over lengths $l = 1/M^k$ for $k \geq 0$.

Proof. Let $l = 1/M^k$. For each cube $I \in \mathcal{B}^d(l, \mathbf{R}^d)$, the l thickening I_l is contained in 3^d cubes in $\mathcal{B}^d(l, \mathbf{R}^d)$. Conversely, if $I \in \mathcal{B}^d(l, E)$, then $I \subset E_{d^{1/2}l}$, so $I \subset E_{M^{k_0}l}$ for $M^{k_0} \geq d^{1/2}$. Thus

$$|E_l| \leq 3^d \#\mathcal{B}^d(l, E)l^d \quad \text{and} \quad |E_{M^{k_0}l}| \geq \#\mathcal{B}^d(l, E)l^d.$$

So as $k \rightarrow \infty$,

$$d - \frac{\log |E_l|}{\log l} = \frac{\log(|E_l|l^{-d})}{\log(1/l)} \leq \frac{\log(3^d \# \mathcal{B}^d(l, E))}{\log(1/l)} = \frac{\# \mathcal{B}^d(l, E)}{\log(1/l)} + o_d(1)$$

and

$$d - \frac{\log |E_{M^{k_0}l}|}{\log(M^{k_0}l)} = \frac{\log(|E_{M^{k_0}l}|(d^{1/2}l)^{-d})}{\log(1/l)} \geq o_d(1) + \frac{\# \mathcal{B}^d(l, E)}{\log(1/l)}.$$

Taking the actual limit as $k \rightarrow \infty$ completes the proof. \square

Our main methods for pattern avoidance use ‘Cantor-type’ constructions. And Lemma 1 will be crucial either for calculating the Minkowski dimension of these constructions. It is especially useful when trying to restrict attention to a finite set of scales, because we can view $\bigcup \mathcal{B}^d(l, E)$ as a ‘discretization’ of a set E at the scale l .

Example. We will often consider sets whose dimension behaves differently at various scales. This often occurs when performing multi-scale constructions where the scales in the construction decay inverse superexponentially. A toy example of this phenomenon can be obtained by modifying the last example slightly so that the construction of the Cantor set behaves differently at various scales. We fix an increasing sequence of integers $\{N_k\}$, with $N_0 = 0$, and consider

$$C = \left\{ \sum_{i=1}^{\infty} a_i / 4^i : a_i \in \{0, 3\} \text{ if there is } k \geq 0 \text{ such that } N_{2k} \leq i \leq N_{2k+1} \right\}.$$

Then $C = \lim C_n$, where

$$C_n = \left\{ \sum_{i=1}^{\infty} a_i / 4^i : a_i \in \{0, 3\} \text{ if there is } k \leq n \text{ such that } N_{2k} \leq i \leq N_{2k+1} \right\}.$$

Counting the number of choices of the a_i given the constraints in the definition of C_n , we find C_n is the union of

$$\prod_{k=0}^{n-1} 2^{N_{2k+1}-N_{2k}} \prod_{k=0}^n 4^{N_{k2+2}-N_{2k+1}} = 2^{-N_1+N_2-N_3+\dots-N_{2n-1}+2N_{2n}} \geq 2^{2N_{2n}-N_{2n-1}}.$$

sidelength l_n cubes, where $l_n = 1/4^{N_{2n}}$. Each of these cubes intersects C , so

$$\frac{\log \# \mathcal{B}^1(l_n, C)}{\log(1/l_n)} \geq \frac{2N_{2n} - N_{2n-1}}{2N_{2n}} = 1 - \frac{N_{2n-1}}{2N_{2n}}.$$

Provided that $N_{2n-1}/N_{2n} = o(1)$, which occurs if the values N_k increase superexponentially, i.e. if $N_k = 2^{k^2}$, we conclude that $\overline{\dim}_{\mathbf{M}}(C) = 1$. On the other hand, C_n is also the union of

$$\prod_{k=0}^n 2^{N_{2k+1} - N_{2k}} \prod_{k=0}^n 4^{N_{2k+2} - N_{2k+1}} = 2^{-N_1 + N_2 - \dots + N_{2n} + N_{2n+1}} \leq 2^{N_{2n} + N_{2n+1}}$$

sidelength r_n cubes, where $r_n = 1/4^{N_{2n+1}}$. Thus

$$\frac{\log \# \mathcal{B}^1(r_n, C)}{\log(1/r_n)} \leq \frac{N_{2n} + N_{2n+1}}{2N_{2n+1}} = 1/2 + \frac{N_{2n}}{2N_{2n+1}}.$$

Again, if $N_{2n}/N_{2n+1} = o(1)$, then $\underline{\dim}_{\mathbf{M}}(C) \leq 1/2$. One can fairly easily check that the values r_n are the ‘worst case’ scales, so that $\underline{\dim}_{\mathbf{M}}(C) = 1/2$. Thus the set C ‘looks’ half dimension between l_n and r_n , for each n , but ‘looks’ full dimensional between the scales r_n and l_{n+1} . This actually impacts the Minkowski dimension of C provided that the gaps between l_n and r_n are made inverse superexponential.

Hausdorff dimension is a version of fractal dimension which is more stable under analytical operations, but is less easy to measure at a single scale. It is obtained by finding a canonical ‘ s dimensional measure’ H^s on \mathbf{R}^d for $s \in [0, \infty)$, which for $s = 1$, measures the ‘length’ of a set, for $s = 2$, measure the ‘area’, and so on and so forth. We then set the dimension of E to be the supremum of the values s such that $H^s(E) < \infty$. For a subset E of Euclidean space, we define the *Hausdorff content*

$$H_\delta^s(E) = \inf \left\{ \sum_{k=1}^{\infty} l_k^s : E \subset \bigcup_{k=1}^{\infty} I_k, I_k \in \mathcal{B}^d(l_k, \mathbf{R}^d), \text{ and } l_k \leq \delta \right\}.$$

We then define $H^s(E) = \lim_{\delta \rightarrow 0} H_\delta^s(E)$. It is easy to see H^s is an exterior measure, and $H^s(E \cup F) = H^s(E) + H^s(F)$ if $d(E, F) > 0$. So H^s is actually a metric exterior measure, and the Caratheodory extension theorem shows H^s is actually a Borel measure.

Example. Let $s = 0$. Then $H_\delta^0(E)$ is the number of δ balls it takes to cover E ,

which tends to ∞ as $\delta \rightarrow 0$ unless E is finite, and in the finite case, $H_\delta^0(E) \rightarrow \#E$. Thus H^0 is just the counting measure.

Example. Let $s = d$. If E has Lebesgue measure zero, then for any $\varepsilon > 0$, there exists a sequence of balls $\{B(x_k, r_k)\}$ covering E with

$$\sum_{k=1}^{\infty} r_k^d < \varepsilon^d.$$

Then we know $r_k < \varepsilon$, so $H_\varepsilon^s(E) < \varepsilon^d$. Letting $\varepsilon \rightarrow 0$, we conclude $H^d(E) = 0$. Thus H^d is absolutely continuous with respect to the Lebesgue measure. The measure H^d is translation invariant, so H^d is actually a constant multiple of the Lebesgue measure.

Lemma 2. If $t < s$ and $H^t(E) < \infty$, $H^s(E) = 0$, and if $H^s(E) = \infty$, $H^t(E) = \infty$.

Proof. If, for any cover of E by balls $B(x_k, r_k)$, $\sum r_k^t \leq A$, and $r_k \leq \delta$, then $\sum r_k^s \leq \sum r_k^{s-t} r_k^t \leq \delta^{s-t} A$. Thus $H_\delta^s(E) \leq \delta^{s-t} A$, and taking $\delta \rightarrow 0$, we conclude $H^s(E) = 0$. The latter point is just proved by taking contrapositives. \square

Example. Since $[-N, N]^d$ has finite Lebesgue measure for each N , if $s > d$, then by Lemma 2, $H^s[-N, N]^d = 0$, and so by countable additivity, $H^s(\mathbf{R}^d) = 0$. Thus H^s is a trivial measure if $s > d$.

Given any Borel set E , the last example, combined with Lemma 2, implies there is a unique value $s_0 \in [0, d]$ such that $H^s(E) = 0$ for $s > s_0$, and $H^s(E) = \infty$ for $0 \leq s < s_0$, though it is possible for $H^{s_0}(E)$ to take any value in $[0, \infty]$, though Lemma 2 implies that s_0 is the only value for which $H^{s_0}(E) \in (0, \infty)$. We refer to s_0 as the *Hausdorff dimension* of E , denoted $\dim_{\mathbf{H}}(E)$.

Theorem 3. For any bounded set E , $\dim_{\mathbf{H}}(E) \leq \underline{\dim}_{\mathbf{M}}(E) \leq \overline{\dim}_{\mathbf{M}}(E)$.

Proof. We consider the simple bound $H_l^s(E) \leq \#\mathcal{B}^d(l, E) \cdot l^s$. Taking $l \rightarrow 0$, we conclude that $H^s(E) \leq \liminf_{l \rightarrow 0} \#\mathcal{B}^d(l, E) \cdot l^s$. Fix $\varepsilon > 0$. If $s \geq \underline{\dim}_{\mathbf{M}}(E) + 2\varepsilon$, consider a sequence of lengths $\{l_k\}$ converging to zero with

$$\frac{\log(\#\mathcal{B}^d(l_k, E))}{\log(1/l_k)} \leq \underline{\dim}_{\mathbf{M}} + \varepsilon \leq s - \varepsilon.$$

Thus $\#\mathcal{B}^d(l_k, E) \leq (1/l_k)^{s-\varepsilon}$, and so $H^s(E) \leq \lim \#\mathcal{B}^d(l_k, E) l_k^s = \lim l_k^\varepsilon = 0$. Taking $\varepsilon \rightarrow 0$ proves the claim. \square

A simple intuition behind the two approaches to fractal dimension we have considered here is that Minkowski dimension measures the efficiency of covers of a set at a fixed scale, whereas Hausdorff dimension measures the efficiency of covers of a set at a simultaneous set of infinitely many scales, which explains both why in certain cases the Hausdorff dimension is smaller than the Minkowski dimension, and also why the Hausdorff dimension is more stable under analytical operations. For instance, for any sequence $\{E_k\}$, $\dim_{\mathbf{H}}(\bigcup_{k=1}^{\infty} E_k) = \sup \dim_{\mathbf{H}}(E_k)$. This need not be true for the Minkowski dimension; a single point has Minkowski dimension zero, but the rational numbers, which are a countable union of points, have Minkowski dimension one. But the Hausdorff dimension still behaves in many respects like the Minkowski dimension. In particular, we can restrict our attention to dyadic-type lengths.

Lemma 3. *For a fixed M , let*

$$H_{\delta}^{s,M}(E) = \inf \left\{ \sum_{k=1}^{\infty} l_k^s : E \subset \bigcup_{k=1}^{\infty} I_k, I_k \in \mathcal{B}^d(1/M^{k_i}, \mathbf{R}^d) \text{ for some } k_i, \text{ and } l_k \leq \delta \right\},$$

and let $H^{s,M}(E) = \lim_{\delta \rightarrow 0} H_{\delta}^{s,M}(E)$. Then $H^s(E) \leq H^{s,M}(E) \leq (3M)^d H^s(E)$.

Proof. Let $E \subset \bigcup_{k=1}^{\infty} I_k$, where $I_k \in \mathcal{B}^d(l_k, \mathbf{R}^d)$, and $l_k \leq \delta$. If l'_k is the smallest multiple of $1/M^k$ greater than l_k , then $l_k \leq l'_k \leq M l_k \leq M \delta$, and $\#\mathcal{B}^d(l'_k, I_k) \leq 3^d$. We now consider the infinite family of cubes $\bigcup_{k=1}^{\infty} \mathcal{B}^d(l'_k, I_k)$. Since

$$\sum_{k=1}^{\infty} l_k^s \leq \sum_{k=1}^{\infty} \#\mathcal{B}^d(l'_k, I_k) \cdot l'_k \leq (3M)^d \sum_{k=1}^{\infty} l_k^s$$

and the cover was arbitrary, we conclude that $H_{\delta}^s(E) \leq H_{M\delta}^{s,M}(E) \leq (3M)^d H_{\delta}^s(E)$. Taking $\delta \rightarrow 0$, we conclude that $H^s(E) \leq H^{s,M}(E) \leq (3M)^d H^s(E)$. \square

It is often easy to upper bound Hausdorff dimension, as in Theorem 3. But it is often non-trivial to lower bound the Hausdorff dimension of a given set. A key technique to this process is *Frostman's lemma*, which says that a set has large Hausdorff dimension if and only if it supports a probability measure which is suitably sparse. We say a measure μ is a *Frostman measure* of dimension s if it is non-zero, compactly supported, and for any length l , if I is a cube with side-length l , then $\mu(I) \lesssim l^s$. The proof of Frostman's lemma will indicate an important principle, known as the *mass distribution principle*.

Lemma 4 (Mass Distribution Principle). *Fix M , and let $\mathcal{B} = \bigcup_{k=1}^{\infty} \mathcal{B}^d(1/M^k, \mathbf{R}^d)$. Suppose μ is a function from \mathcal{B} to $[0, \infty)$ such that for any $I \in \mathcal{B}^d(1/M^k, \mathbf{R}^d)$,*

$$\sum \{ \mu(J) : J \in \mathcal{B}^d(1/M^{k+1}, I) \} = \mu(I)$$

Furthermore, assume that

$$\sum \{ \mu(I) : I \in \mathcal{B}^d(1, I) \} < \infty$$

Then μ extends to a unique finite Borel measure on \mathbf{R}^d .

Proof. For each k , let E_k be the operator mapping Borel measures to functions on \mathbf{R}^d , defined by

$$E_k(\mu) = \sum \{ \mu(I) \cdot \chi_I : I \in \mathcal{B}^d(1/M^k, \mathbf{R}^d) \}$$

Define a sequence of Borel measures μ_k by

$$\mu_k(E) = \sum \{ \mu(I) |E \cap I| : I \in \mathcal{B}^d(1/M^k, \mathbf{R}^d) \}$$

The main condition of the theorem can be summarized by saying that $E_j(\mu_k) = \mu_j$ if $j \leq k$. An important thing to notice about the operators E_k is that they are continuous from the weak topology to the pointwise convergence topology. Indeed, if $\nu_i \rightarrow \nu$ weakly, then $\nu_i(I) \rightarrow \nu(I)$ for each fixed $I \in \mathcal{B}^d(1/M^k, \mathbf{R}^d)$. Since χ_I and $\chi_{I'}$ have disjoint support if $I \neq I'$, $E_k(\nu_i) \rightarrow E_k(\nu)$ pointwise. Since the measures μ_k are finite, with total variation $\|\mu_k\|$ bounded independantly of k , the weak compactness of the unit ball implies that there is a subsequence μ_{k_i} converging weakly to some measure μ' . But then by continuity,

$$E_j(\mu') = \lim_{i \rightarrow \infty} E_j(\mu_{k_i}) = \mu_j$$

which implies that $\mu' = \mu$, wherever the two are both defined. \square

In the next lemma, it will help to notice that the definition of the Frostman measure is also robust to working over dyadic-type lengths. Suppose we can establish a result $\mu(I) \lesssim 1/M^{ks}$ for all $I \in \mathcal{B}^d(1/M^k, \mathbf{R}^d)$ and all indices k . Given any length l , there is a value of k such that $1/M^{k+1} \leq l \leq 1/M^k$, so $1/M^k \leq Ml$. For any $I \in \mathcal{B}^d(l, \mathbf{R}^d)$, $\# \mathcal{B}^d(1/M^k, I) \leq 3^d$, so

$$\mu(I) \leq \mu\left(\bigcup \mathcal{B}^d(1/M^k, I)\right) \lesssim 3^d / M^{ks} \leq 3^d / (Ml)^s \lesssim_M 1/l^s$$

Thus μ is a Frostman measure.

Lemma 5. *let μ^+ be a function from \mathcal{B} to $[0, \infty)$ such that for any $I \in \mathcal{B}^d(1/M^k, \mathbf{R}^d)$,*

$$\sum \{\mu^+(J) : J \in \mathcal{B}^d(1/M^{k+1}, I)\} \leq \mu^+(I)$$

Assume there exists $c > 0$ such that for all k ,

$$\sum \{\mu^+(I) : I \in \mathcal{B}^d(1/M^k, I)\} \geq c$$

and

$$\sum \{\mu^+(I) : I \in \mathcal{B}^d(1, I)\} < \infty$$

Then there exists a non-zero Borel measure μ such that $\mu(I) \leq \mu^+(I)$ for $I \in \mathcal{B}$.

Proof. As in the last lemma, define the operators E_k and the measures μ_k . By weak compactness, a subsequence of these measures converge weakly to some measure μ , and $E_k(\mu) = \lim E_k(\mu_{j_k}) \leq \mu_k$. The measure μ is nonzero, since $\|\mu_{j_k}\| \geq c$ for each k , and so $\|\mu\| \geq c$. \square

Lemma 6 (Frostman's Lemma). *If E is Borel, $H^s(E) > 0$ if and only if there exists an s dimensional Frostman measure supported on E .*

Proof. Suppose that μ is s dimensional and supported on E . If $H^s(F) = 0$, then for $\varepsilon > 0$ there is a sequence of cubes $\{I_k\}$ and lengths $\{l_k\}$ with $I_k \in \mathcal{B}^d(l_k)$ and $\sum_{k=1}^{\infty} l_k^s \leq \varepsilon$. But then

$$\mu(F) \leq \mu\left(\bigcup_{k=1}^{\infty} I_k\right) \leq \sum_{k=1}^{\infty} \mu(I_k) \lesssim \sum_{k=1}^{\infty} l_k^s \leq \varepsilon.$$

Taking $\varepsilon \rightarrow 0$, we conclude $\mu(F) = 0$, so μ is absolutely continuous with respect to H^s . Thus there exists a Radon Nikodym derivative $d\mu/dH^s \in L^1(\mathbf{R}^d, H^s)$ such that

$$\mu(F) = \int_F \frac{d\mu}{dH^s} dH^s.$$

In particular,

$$\mu(E) = \int_E \frac{d\mu}{dH^s} dH^s = 1,$$

so we must have $H^s(E) > 0$. Conversely, suppose $H^s(E) > 0$. Then by translating, we may assume that $H^s(E \cap [0, 1]^d) > 0$, and so without loss of generality we may

assume that $E \subset [0, 1]^d$. Fix M , and for each $I \in \mathcal{B}^d(1/M^k, \mathbf{R}^d)$, define

$$\mu^+(I) = H_{1/M^k}^{s,M}(E \cap I)$$

Then $\mu^+(I) \leq 1/M^{ks}$, and μ^+ is subadditive. We use it to recursively define a measure μ to which we can apply Lemma 4, such that $\mu(I) \leq \mu^+(I)$ for each $I \in \mathcal{B}^d(1/M^k)$. We initially define μ by setting $\mu([0, 1]^d) = \mu^+([0, 1]^d)$. Given $I \in \mathcal{B}^d(1/M^k, [0, 1]^d)$, we enumerate all the children $J_1, \dots, J_M \in \mathcal{B}^d(1/M^{k+1}, I)$. We then consider any values of A_1, \dots, A_M such that

$$A_1 + \dots + A_M = \mu(I) \quad \text{and} \quad A_i \leq \mu^+(J_i)$$

This is feasible to do because $\sum_{i=1}^M \mu^+(J_i) \geq \mu^+(I) \geq \mu(I)$. We then define $\mu(J_i) = A_i$. The recursive constraint is satisfied, so μ is well defined. The mass distribution principle then implies that μ extends to a full measure, which satisfies $\mu(I) \leq \mu^+(I) \leq 1/M^{ks}$ for each $I \in \mathcal{B}^d(1/M^k, \mathbf{R}^d)$, so if we rescale μ so it has total mass one, we find it is a Frostman measure with dimension s . \square

Thus the problem of lower bounding Hausdorff dimension reduces to the problem of upper bounding the mass of certain measures constructed on sets. Because we consider Cantor-type constructions, and we define measures μ by the mass distribution principle, it is most natural to establish bounds $\mu(I) \lesssim_\varepsilon l_k^{s-\varepsilon}$ when $I \in \mathcal{B}_{l_k}^d$, for a sequence of scales l_k which decrease rapidly. Using certain tighter estimates than required, these bounds are sufficient to guarantee that μ is $s - \varepsilon$ dimensional for all $\varepsilon > 0$.

Lemma 7. *Let E be a set, and μ a Borel probability measure supported on E . Suppose that for any ε , there exists a constant c_ε such that if $\mathcal{B}^d(1/M^k, E) \leq c_\varepsilon M^{k(s-\varepsilon)}$, then $\mu(E) \lesssim 1/k^2$. then E has Hausdorff dimension s .*

Proof. Suppose $H^{s-\varepsilon}(E) = 0$. Then for any N there exists a cover of E by cubes $\{I_k\}$, with lengths $\{l_k\}$ such that $I_k \in \mathcal{B}^d(l_k, \mathbf{R}^d)$, $l_k \leq 1/M^N$ for all k , and $\sum l_k^{s-\varepsilon} \leq c_\varepsilon/M^d$. For each m , let $A_m = \#\{k : 1/M^{m+1} \leq l_k \leq 1/M^m\}$. Then

$$\sum_{m=N}^{\infty} A_m M^{-(m+1)(s-\varepsilon)} \leq \sum_{k=1}^{\infty} l_k^{s-\varepsilon} \leq c_\varepsilon/M^d$$

Thus $A_m \leq c_\varepsilon M^{(m+1)(s-\varepsilon)}$. This means that if E_m is formed from the union of all intervals I_k with $1/M^{m+1} \leq l_k \leq 1/M^m$, then $\#\mathcal{B}_s^d(E_m) \leq c_\varepsilon M^{(m+1)(s-\varepsilon)}$. Thus

$\mu(E_m) \lesssim 1/k^2$, so

$$\mu(E) \leq \sum_{m=N}^{\infty} \mu(E_m) \lesssim \sum_{m=N}^{\infty} 1/k^2$$

as $N \rightarrow \infty$, we conclude $\mu(E) = 0$, which is impossible. Thus $H^{s-\varepsilon}(E) > 0$ for all ε , so $\dim_{\mathbf{H}}(E) \geq s$. \square

The hypothesis of Lemma 7 is certainly satisfied if $\mu(I) \lesssim_{\varepsilon} l_k^{s-\varepsilon}/k^2$ for each $I \in \mathcal{B}^d(l_k)$, where $l_k = 1/M^k$. Thus establishing a Frostman-type bound at a sequence of dyadic type scales is enough to obtain a dimensional result for E . The advantage of this proof is that we can continue the argument to give results when the sequence of scales decreases much faster than scales of dyadic type.

Theorem 4. *Let E be a set, and μ a Borel probability measure supported on E . Suppose that for any ε , there exists a constant c_{ε} such that if $\mathcal{B}^d(1/M^k, E) \leq c_{\varepsilon} M^{k(s-\varepsilon)}$, then $\mu(E) \lesssim 1/k^2$. then E has Hausdorff dimension s .*

Proof. As before, if $H^{s-\varepsilon}(E) = 0$, consider a covering by $\{I_k\}$ with parameters $\{l_k\}$. Fix $\alpha > 1$, and consider

$$A_m = \#\{k : 1/M^{\alpha^{m+1}} \leq l_k \leq 1/M^{\alpha^m}\}$$

Then

$$\sum A_m / M^{(s-\varepsilon)\alpha^{m+1}} \leq \sum l_k^{s-\varepsilon} \leq c_{\varepsilon}$$

so $A_m \leq c_{\varepsilon} M^{(s-\varepsilon)\alpha^{m+1}}$. Thus if we define E_m , then $\mu(E_m) \lesssim 1/k^2$. \square

The requirement of this lemma is satisfied if we are able to prove $\mu(I) \lesssim_{\varepsilon} l_k^{s-\varepsilon}/k^2$ where $l_k = M^{-\alpha^k}$. These are *hyperdyadic* numbers.

Proof. Suppose that $H^{s-\varepsilon}(E) = 0$. Then for any M and c_{ε} , E is covered by cubes $\{I_k\}$ with sidelengths $\{r_k\}$ such that $I_k \in \mathcal{B}^d(r_k, \mathbf{R}^d)$, $r_k \leq l_M$ for each k , and $\sum r_k^{s-\varepsilon} \leq c_{\varepsilon}$. For each k , let $A_m = \#\{k : l_{m+1} \leq r_k \leq l_m\}$. Then

$$\sum_{m=M}^{\infty} A_m l_{m+1}^{s-\varepsilon} \leq \sum_{k=1}^{\infty} r_k^{s-\varepsilon} \leq c_{\varepsilon}$$

so $A_m \leq c_{\varepsilon} / l_{m+1}^{s-\varepsilon}$. But if we can establish an estimate $\mu(I) \lesssim_{\varepsilon} B_m l_m^{s-\varepsilon}$, then

$$\mu(E) \lesssim_{\varepsilon} \sum_{m=M}^{\infty} A_m (B_m l_m^{s-\varepsilon}) \leq \sum_{m=M}^{\infty} c_{\varepsilon} B_m (l_m / l_{m+1})^{s-\varepsilon}$$

□

2.2 Hyperdyadic Covers

Theorem 5. *If $\inf_{\delta>0} H_\delta^s(E) = 0$, then $\dim_{\mathbf{H}}(E) \leq s$.*

Proof. Suppose that $H_\delta^s(E) \leq \varepsilon$. Then there is a sequence of cubes $\{I_k\}$ with parameters $\{l_k\}$ such that $I_k \in \mathcal{B}^d(l_k)$, $l_k \leq \delta$ for all k , and $\sum_{k=1}^\infty l_k^s \leq 2\varepsilon$. Then $l_k \leq (2\varepsilon)^{1/s}$ for all k , so we can actually take $\delta \rightarrow 0$? □

Theorem 6. *If X is strongly covered by $\{X_k\}$, and there is δ such that*

$$\sum_{k=1}^\infty H_\delta^s(X_k) < \infty$$

then $\dim_{\mathbf{H}}(X) \leq s$.

Proof. By subadditivity, for any N ,

$$H_\delta^s(X) \leq \sum_{k=N}^\infty H_\delta^s(X_k)$$

and as $N \rightarrow \infty$, we conclude $H_\delta^s(X) = 0$. □

For a fixed $0 < \varepsilon \ll 1$, a *hyperdyadic number* is a number of the form $h_k = 2^{-\lfloor (1+\varepsilon)^k \rfloor}$ for some $k \geq 0$. A set E is δ *discretized* if it is the union of balls, each with radius between $c_\varepsilon \delta^{1+C\varepsilon}$ and $C_\varepsilon \delta^{1-C\varepsilon}$. A set E is a $(\delta, s)_d$ set if it is bounded, δ discretized, and for all $\delta \leq r \leq 1$, $|E \cap B(x, r)| \leq C_\varepsilon \delta^{n-s-C\varepsilon} r^s$.

Lemma 8. *Let $0 < s < d$, and let E be a compact subset of \mathbf{R}^n .*

- *If $\dim(E) \leq s$, for each k , we can associate a $(h_k, s)_d$ set E_k such that E is strongly covered by the E_k .*
- *If C is sufficiently large, and there is a $(h_k, s-C\varepsilon)_n$ set E_k strongly covering E , then $\dim(E) \leq s$.*

Proof. We first prove the latter claim, assuming without loss of generality that E is contained in the unit ball. Suppose E is strongly covered by the E_k . If E_k is a $(h_k, s-C\varepsilon)_d$ set, then

$$|E_k| \leq C_\varepsilon \delta^{n-(s-C\varepsilon)-C\varepsilon} = C_\varepsilon \delta^{n-s}.$$

$$H_{C_\varepsilon h_k^{1-C_\varepsilon}}^s$$

□

2.3 Branching Processes

2.4 Equidistribution Results

The classical Weyl equidistribution theorem says that if α is an irrational number, then the decimal parts of the numbers $n\alpha$ are equidistributed in $\mathbf{T} = \mathbf{R}/\mathbf{Z}$, in the sense that for any continuous function $f : \mathbf{T} \rightarrow \mathbf{R}$,

$$\frac{1}{N} \sum_{n \leq N} f(n\alpha) \rightarrow \int_{\mathbf{T}} f(x) dx$$

This is equivalent to prove that for any interval $[a, b] \in [0, 1)$,

$$\frac{\#\{1 \leq n \leq N : x_n \in [a, b]\}}{N} \rightarrow b - a$$

as $N \rightarrow \infty$. By approximating a continuous function f by a Fourier series, to prove this is true for a particular sequence, it suffices to prove it for $f(x) = e^{2\pi i n x}$, for each nonzero integer n . Certain techniques we are developing in the theory of cantor decompositions require a higher dimensional variant of such a result, so this section details some information which might help us in the future. We will encounter sequences that are not equidistributed over the entire space, so if G is any closed subgroup of \mathbf{T} , we say a sequence x_n is equidistributed over G if $x_n \in G$ for all n , and for any continuous function $f : G \rightarrow \mathbf{R}$,

$$\frac{1}{N} \sum_{n \leq N} f(n\alpha) \rightarrow \int_G f(x) dx$$

or alternatively, if for any closed set K in G ,

$$\frac{\#\{1 \leq n \leq N : x_n \in K\}}{N} \rightarrow |K|$$

where $|K|$ is taken with respect to the Haar probability measure on G .

Theorem 7. A sequence x_1, x_2, \dots is equidistributed in \mathbf{T}^d if and only if for every nonzero $\xi \in \mathbf{Z}^n$,

$$\frac{1}{N} \sum_{n \leq N} e(\xi \cdot x_n) \rightarrow 0$$

Proof. We prove that the exponential sum condition implies the general result, noting that the other direction is clear. Clearly the exponential sum condition implies the result for all functions f which are trigonometric polynomials. But then, by the Stone Weirstrass theorem and basic Abelian Harmonic analysis, the multivariate trigonometric polynomials are dense in $C(\mathbf{T}^d)$, and we may apply a standard limiting argument. \square

Example. If $x_n = n\alpha + \beta$, then

$$\frac{1}{N} \sum_{n \leq N} e(\xi \cdot x_n) = \frac{e(\xi \cdot \beta)}{N} \sum_{n \leq N} e(n\xi \cdot \alpha) = \begin{cases} \frac{1}{N} \frac{e(\xi \cdot \beta)(e((n+1)\xi \cdot \alpha) - 1)}{e(\xi \cdot \alpha) - 1} & : \xi \cdot \alpha \notin \mathbf{Z} \\ e(\xi \cdot \beta) & : \xi \cdot \alpha \in \mathbf{Z} \end{cases}$$

Weyl's exponential sum theorem implies that x_n is equidistributed on \mathbf{T}^n precisely when $\xi \cdot \alpha \notin \mathbf{Z}$ for all $\xi \in \mathbf{Z}^n$. What's more, it is simple to see from this that x_n is still equidistributed for any subsequence whose indices form an arithmetic progression.

Thus in one dimension, an arithmetic sequence is either equidistributed over the entire torus, or over a discrete set of points forming a *discrete subgroup* of the torus. We can think of this discrete subgroup as a zero dimension torus, which leads us to suspect that in higher dimensions, an arithmetic sequence is always equidistributed, but not necessarily over the whole torus, but instead over a lower dimensional subtorus.

Theorem 8 (Ratner). If $x_n = n\alpha + \beta$, then we can write $\alpha = \alpha_0 + \alpha_1$, where the sequence $n\alpha_0 + \beta$ is periodic in \mathbf{T}^d , and $n\alpha_1 + \beta$ is equidistributed over a subtorus of \mathbf{T}^d . In particular, if $\beta = 0$, then Ratner's theorem says that x_n has an evenly spaced subsequence equidistributed over a subtorus of the space containing the origin.

Proof. We induct on the dimension. For $d = 1$, the theorem is obvious, since either α is rational, and therefore $n\alpha + \beta$ is periodic, or α is irrational, and $n\alpha + \beta$ is equidistributed on \mathbf{T} . So now let us consider a sequence on the torus \mathbf{T}^{d+1} . If α is irrational, then x_n is equidistributed, and the theorem is obvious. Otherwise, there exists $\xi \in \mathbf{Z}^n$ such that $\xi \cdot \alpha \in \mathbf{Z}$. We may write $\alpha = \alpha_0 + \alpha_1$, where $\alpha_0 \in$

\mathbf{Q}^d , and $\xi \cdot \alpha_1 = 0$. The sequence $n\alpha_0 + \beta$ is periodic, whereas $n\alpha_1$ takes values in the subtorus T of points x with $\xi \cdot x = 0$. Since $\xi \neq 0$, T is a d dimensional compact subgroup of \mathbf{T}^{d+1} isomorphic to \mathbf{T}^d . To see this, we assume for simplicity that $\xi_d \neq 0$. Then ξ^\perp is generated by the basis of d vectors $v_n = \xi_d e_n - \xi_n e_d$, for $1 \leq n \leq d$. Thus we get a homomorphism between \mathbf{R}^d and T given by the map $x \mapsto \sum x_n v_n$. If $\sum x_n v_n \in \mathbf{Z}$, so that $\sum x_n v_n = 0$ on T , then this means that $x_n \in \mathbf{Z}/\xi_d$ for each n , implying that the kernel of this homomorphism is discrete. Lattice theory implies that we can write the kernel as $\oplus \mathbf{Z}\langle w_n \rangle$, for d generating vectors w_1, \dots, w_d . But then the map $f(x) = \sum x_n w_n$ gives an isomorphism between \mathbf{T}^d and T . If we set $\beta = f^{-1}(\alpha_0)$, then $f^{-1}(n\alpha_0) = n\beta$, and so by induction, we can write $\beta = \beta_0 + \beta_1$, where $n\beta_0$ is periodic, and $n\beta_1$ is equidistributed over a subtorus of \mathbf{T}^d . But then $nf(\beta_0) = f(n\beta_0)$ is periodic on T , and $nf(\beta_1) = f(n\beta_1)$ is equidistributed over a subtorus of T . Since the sum of two periodic sequences is periodic, $n\alpha_0 + nf(\beta_0)$ is periodic, and $nf(\beta_1)$ is equidistributed over a subtorus. We have $\alpha_0 + f(\beta_0) + f(\beta_1) = \alpha_0 + f(\beta) = \alpha_0 + \alpha_1$, completing the proof. \square

Corollary. *Any linear sequence in \mathbf{T}^d is equidistributed in a finite union of cosets of a subtorus of \mathbf{T}^d .*

In general, ergodic theory results do not give rates on how long it takes for a sequence to equidistribute over a set. This is not a problem in the constructions we perform, since the rates that our intervals shrink can be arbitrarily fast. However, it is important to note that the convergence rates are uniform across all intervals.

Theorem 9. *If x_n is equidistributed over a torus \mathbf{T}^d , and*

$$A_N = \sup_I \left| \frac{\#\{1 \leq n \leq N : x_n \in I\}}{N} - |I| \right|$$

where I ranges over all boxes in \mathbf{T}^d , then $A_N \rightarrow 0$ as $N \rightarrow \infty$.

Proof. For notational simplicity, we let

$$\#(I, N) = \#\{1 \leq n \leq N : x_n \in I\}$$

For each n , we can partition \mathbf{T}^d into finitely many disjoint cubes $\{I_n\}$ with side-lengths $1/n$. Since there are only finitely many such cubes, there is N_n such that for $M \geq N_n$, $|\#(I_n, M)/M - |I_n|| \leq 1/n$. Now given any box J , we can find sets J_1 and J_2 , each unions of the cubes I_n , with $J_1 \subset J \subset J_2$ and $|J - J_1|, |J_2 - J| \lesssim_d 1/n$. Thus

$$|J| - \frac{2}{n} \leq |J_1| - \frac{1}{n} \leq \frac{\#(J_1, M)}{M} \leq \frac{\#(J, M)}{M} \leq \frac{\#(J_2, M)}{M} \leq |J_2| + \frac{1}{n} \leq |J| + \frac{2}{n}$$

which completes the proof, since N_n is independent of J . □

2.5 Results about Hypergraphs

Lemma 9 (Turán). *For any k uniform hypergraph $H = (V, E)$ with $|E| \leq |V|^\alpha$, V contains an independant set of size $\Omega(|V|^{(k-\alpha)/(k-1)})$.*

Proof. We create an independant set I by the following procedure. First, select a subset S of vertices, including each independently with probability p . Delete a single vertex from each edge in each hypergraph entirely contained in S , obtaining an independant set I . We find that each edge in V is entirely included in S with probability p^k , and S has expected size $p|V|$, so $\mathbf{E}|I| = p|V| - p^k|E|$. If $|E| = |V|^\alpha$ for $\alpha \geq 1$, then setting $p = (1/2)|V|^{(1-\alpha)/(k-1)}$ induces a set I with size

$$|V|^{(k-\alpha)/(k-1)}(1/2 - 1/2^k)$$

We create an independant set I by the following procedure. First, select a subset S of vertices, including each vertex independently with probability p . Delete a single vertex from each edge in each hypergraph which is entirely contained in S . Then I is an independant set with respect to each hypergraph, and we shall show that for an appropriate choice of p , $\mathbf{E}|I| \geq h$.

Trivially, we find $\mathbf{E}|S| = p|V|$. For any $i \geq 2$, the expected number of edges of H_i falling entirely in S is

$$p^i|E_i| \leq \frac{p^i|V|^i}{c_k h^{i-1}}$$

therefore

$$\mathbf{E}|I| = p|V| - \sum_{i=2}^k \frac{p^i|V|^i}{c_k h^{i-1}}$$

Setting $p = 2h/|V|$ and $c_k = 2^{k+1}$ gives

$$\mathbf{E}|I| = h \left(2 - \sum_{i=2}^k \frac{1}{2^{k+1-i}} \right) > h$$

which completes the proof. □

2.6 Hyperdyadic Covers

Recall the definition of the Hausdorff measure $H^\alpha(E) = \lim_{\delta \rightarrow 0} H_\delta^\alpha(E)$, where $H_\delta^\alpha(E)$ is the greatest lower bound of $\sum r_n^\alpha$, over all choices of covers of E by cubes I_1, I_2, \dots , where I_n has sidelengths r_n . We then define the Hausdorff dimension of E to be the least upper bound of the scalars α such that $H^\alpha(E) = 0$, or alternatively, the greatest lower bound of α such that $H^\alpha(E) = \infty$.

To determine the Hausdorff dimension of E , it suffices to consider only dyadic cubes in the cover of E . Define $H^\alpha(E) = \lim_{\delta \rightarrow 0} H_{D,\varepsilon}^\alpha(E)$, where $H_{D,\varepsilon}^\alpha(E)$ is the greatest lower bound of $\sum r_n^\alpha$ over *dyadic* covers I_1, I_2, \dots , with $I_n \in \mathcal{B}(r_n)$. Then H_D^α is comparable with H^α .

Theorem 10. *For any set E , $H^\alpha(E) \leq H_D^\alpha(E) \leq 2^{d+\alpha} H^\alpha(E)$.*

Proof. Given any not necessarily dyadic cover I_1, I_2, \dots , we can replace each sidelength r_n cube I_n with at most 2^d dyadic cubes with radius at most $2r_n$, which gives $H_{D,\varepsilon}^\alpha(E) \leq 2^{1+\alpha} H_\varepsilon^\alpha(E)$, and taking the limit as $\varepsilon \rightarrow 0$ then gives the required upper bound for H_D^α . \square

If we are restricting ourselves to cubes lying at a series of discrete scales, it seems as if the dyadic sequence is about as fast as we can use so that the resultant Hausdorff measure is comparable to the usual Hausdorff measure. Nonetheless, using a weak type bound we can get results for a faster decreasing family of scales. This is necessary for our calculations. We fix a positive δ , and consider a sequence of **hyperdyadic scales** $H_N = 2^{-\lfloor (1+\delta)^N \rfloor}$. A **hyperdyadic cube** is then a cube in $\mathcal{B}(H_N)$ for some N .

Proof. For any sidelength L cube, we can cover the cube by at most 2^d hyperdyadic cubes with sidelength at most $2L^{1-\delta} \geq 2L^{(1+\delta)^{-1}}$. This is because

$$2H_{N+1}^{(1+\delta)^{-1}} = 2^{1-(1+\delta)^{-1}\lfloor (1+\delta)^{N+1} \rfloor} \geq 2^{1-(1+\delta)^N} \geq 2^{\lfloor (1+\delta)^N \rfloor} = H_N$$

If E has Hausdorff dimension α , for every ε and N we can find a collection of dyadic cubes I_1, I_2, \dots covering E with I_k sidelength $L_k \leq H_N$, and $\sum L_{N,i}^{\alpha+\varepsilon} \lesssim_\varepsilon 1$. A weak type bound implies the number of cubes I_k with $H_{N+1} \leq L_k \leq H_N$ is $O_\varepsilon(1/H_{N+1}^{\alpha+\varepsilon})$. But

$$1/H_{N+1}^{\alpha+\varepsilon} \leq (H_N/H_{N+1})^{\alpha+\varepsilon} 1/H_N^{\alpha+\varepsilon} \lesssim 1/H_N^{\alpha+\varepsilon+\delta}$$

and so the cover of E by hyperdyadic cubes contains $O_\varepsilon(1/H_N^{\alpha+\varepsilon+\delta})$ length H_N cubes for each N .

If we swap each cube $I_{N,i}$ with 2^d hyperdyadic cubes of length at most $2L^{1-\delta}$, we obtain

$$\sum 2^d (2L_{N,i}^{1-\delta})^{\alpha+\varepsilon} = 2^{d+\alpha+\varepsilon} \sum L_{N,i}^{(1-\delta)(\alpha+\varepsilon)} \lesssim_\varepsilon 1$$

Thus $H_{HD}^{(1-\delta)\alpha+\varepsilon}(E) \lesssim_\varepsilon 1$.

We can swap each cube I_i with 2^d hyperdyadic cubes of length at most $2L^{(1+\delta)^{-1}}$, without effecting the estimate too much.

Then for every hyperdyadic number H_N , we can find a collection of cubes $I_{N,1}, I_{N,2}, \dots$ covering E with $I_{N,i}$ sidelength $r_{N,i} \leq H_N$, and $\sum r_{N,i}^{\alpha+\varepsilon} \lesssim_\varepsilon 1$. Covering each cube by 2^d cubes with hyperdyadic sidelengths, which magnifies $r_{N,i}$ by at most

$$2 \cdot 2^{(1+\delta)^{N+1} - (1+\delta)^N} = 2 \cdot 2^{\delta(1+\delta)^N} \lesssim 2 \cdot r_{N,i}^{-\delta}$$

We conclude that

$$2^{d+\alpha+\varepsilon} 2^{(\alpha+\varepsilon)\delta(1+\delta)^N} C_\varepsilon$$

□

We assume δ and ε are some fixed parameters. If $A(\varepsilon, \delta)$ and $B(\varepsilon, \delta)$ are two quantities depending on ε and δ , we write $A \leq B$ mean $A \lesssim_\varepsilon \delta^{-C\varepsilon} B$ for some C , and for every ε . We let $A \approx B$ mean $A \leq B$ and $B \leq A$ hold simultaneously. We say a union of balls is δ discretized if it is the union of balls with radius $\approx \delta$. Thus there exists C_ε and C such that for each ball B_r of radius r , $|r - \delta| \leq C_\varepsilon \delta^{1-C\varepsilon}$. Thus

$$\delta(1 - C_\varepsilon \delta^{-C\varepsilon}) \leq r \leq \delta(1 + C_\varepsilon \delta^{-C\varepsilon})$$

In particular, the dyadic scales $2^{-\lfloor (1+\varepsilon)^k \rfloor}$ are allowed in a discretization of a hyperdyadic scale $2^{-(1+\varepsilon)^k}$, since we can choose C_ε and C such that

$$1 - C_\varepsilon 2^{C(1+\varepsilon)^k \varepsilon} \leq 1 \leq 2^{(1+\varepsilon)^k - \lfloor (1+\varepsilon)^k \rfloor} \leq 2 \leq 1 + C_\varepsilon 2^{(1+\varepsilon)^k C\varepsilon}$$

Theorem 11. *Let E be a compact subset of \mathbf{R}^n . If $0 < \alpha < n$, and $\dim(E) \leq \alpha$, then for each hyperdyadic number δ , we can associate a δ discretized set X_δ with $|X_\delta \cap B(x, r)| \leq \delta^n (r/\delta)^\alpha$ for all $\delta \leq r \leq 1$ and $x \in \mathbf{R}^n$, and every element of E is contained in infinitely many of the X_δ .*

Proof. Fix E . For every hyperdyadic δ , we can find a cover of E by balls $B(x_{\delta n}, r_{\delta n})$

such that $r_{\delta n} < \delta$, and

$$\sum_n r_{\delta n}^{\alpha+C\varepsilon} \lesssim 1 \quad (2.1)$$

Choose $m_{\delta n}$ such that $2^{-(1+\varepsilon)^{m_{\delta n}+1}} \leq r_{\delta n} \leq 2^{-(1+\varepsilon)^{m_{\delta n}}}$. We calculate

$$\begin{aligned} \frac{2^{-(\alpha+C'\varepsilon)(1+\varepsilon)^{m_{\delta n}}}}{r_{\delta n}^{\alpha+C\varepsilon}} &\leq \frac{2^{-(\alpha+C'\varepsilon)(1+\varepsilon)^{m_{\delta n}}}}{2^{-(\alpha+C\varepsilon)(1+\varepsilon)^{m_{\delta n}+1}}} \\ &= \left(2^{\varepsilon(1+\varepsilon)^{m_{\delta n}}}\right)^{\alpha+(C(1+\varepsilon)-C')} \end{aligned}$$

Provided that $C' > \alpha + C(1+\varepsilon)$, the quantity on the left is ≤ 1 , which is independent of ε provided that ε is bounded from above, and so we conclude

$$\sum_n \left(2^{-(1+\varepsilon)^{m_{\delta n}}}\right)^{\alpha+C'\varepsilon} \leq \sum_n r_{\delta n}^{\alpha+C\varepsilon}$$

Thus we may assume by changing the value of C that the quantities $r_{\delta n}$ are hyperdyadic from the outset. This means that at each hyperdyadic scale δ , the number of hyperdyadic balls at the scale δ in each cover is $\lesssim (1/\delta)^{\alpha+C\varepsilon}$. STOP IS THIS ALL WE NEED, THEN COME BACK TO THE PROOF.

For a pair of hyperdyadic numbers δ and γ we set

$$Y_{\delta\gamma} = \bigcup_{r_{\delta n}=\gamma} B(x_{\delta n}, r_{\delta n})$$

Every element of X is in infinitely many of the $Y_{\delta n}$. For each δ and γ , we let $Q_{\delta\gamma}$ be the collection of hyperdyadic cubes with sidelength at least γ covering $Y_{\delta\gamma}$ and minimizing $\sum_{Q \in Q_{\delta\gamma}} l(Q)^\alpha$. From condition (1.1) we obtain that $Y_{\delta\gamma}$ can be covered by at most $r^{-\alpha-\varepsilon}$ sidelength r cubes, so

$$\sum_{Q \in Q_{\delta\gamma}} l(Q)^\alpha \leq Cr^{-\varepsilon}$$

and so $l(Q) \leq Cr^{-\varepsilon/\alpha}$ for all $Q \in Q_{\delta\gamma}$. From the construction of $Q_{\delta\gamma}$, we see that the Q are all disjoint, and for any hyperdyadic cube I ,

$$\sum_{\substack{Q \in Q_{\delta\gamma} \\ Q \subset I}} l(Q)^\alpha \leq l(I)^\alpha$$

since otherwise we could replace such elements of Q in $Q_{\delta\gamma}$ by I itself. \square

2.7 Dimensions of Cantor Like Sets

This section provides tools to calculate the Hausdorff dimension of Cantor like sets. We analyze the dimension of compact sets $X \subset [0, 1]^d$ by viewing the set as a limit of a nested family of discretized sets X_N , each the union of cubes of a fixed length. For such a construction, we can naturally associate a sequence of Borel probability measures μ_N supported on X_N , weakly converging to a measure μ supported on X . Our main tools will establish estimates about μ through estimates on μ_N , which then allows us to apply Frostman's lemma.

To begin with, we introduce some notation. For a given length L , we let $\mathcal{B}(L)$ denote the collection of all cubes whose corners lie on the lattice $(L\mathbb{Z})^d$. Given a set X , $\mathcal{B}(X, L)$ shall denote all cubes in $\mathcal{B}(L)$ intersecting X . For a given cube I , we let $L(I) = |I|^{1/d}$ denote the sidelength of the cube. It will be natural to consider a decreasing sequence of lengths L_N with $L_{N+1} \mid L_N$ for all N . The divisor condition is natural so that the cover $\mathcal{B}(X, L_{N+1})$ is a refinement of the cover $\mathcal{B}(X, L_N)$. Because of this, it is obvious that $X_{N+1} \subset X_N$, and that they converge to X as $N \rightarrow \infty$.

Since the initial set X_0 is a union of intervals, we can form a probability measure μ_0 supported on it by placing a uniform mass over all points in X_0 . We perform an inductive definition of the remaining μ_N . To form μ_{N+1} from μ_N , we consider each $I \in \mathcal{B}(X, L_N)$, and distribute the mass $\mu_N(I)$ uniformly over the intervals in $\mathcal{B}(X, L_{N+1})$ which intersect I . It is easy to see that the distribution functions of the μ_N converge pointwise, so the measures μ_N must converge weakly to some measure μ supported on X . Most importantly, it satisfies $\mu(I) = \mu_N(I)$ for each $I \in \mathcal{B}(L_N)$. It is easy to obtain estimates on the measures μ_N from combinatorial arguments, especially when X is constructed in the Cantor set style, as a limit of sets of the form X_N . This appendix discusses how we can use estimates on μ from these estimates, so that we may apply Frostman's lemma to get Hausdorff dimension bounds for X .

Remark. *I am still not sure how 'natural' μ is to the problem. If $\mu(I) \lesssim |I|^\alpha$ is not satisfied for all I , does it follow that $\dim_{\mathbf{H}}(X) \leq \alpha$? It seems this probability measure is strongly related to the structure of the Cantor set so this isn't entirely implausible.*

2.8 Basic Covering

It is easy to establish estimates of the form $\mu(I) = \mu_N(I) \lesssim L_N^\alpha$ for $I \in \mathcal{B}(L_N)$, independent of N by counting arguments. To obtain general bounds, we must apply covering arguments. For any cube I with $L_{N+1} \leq L(I) \leq L_N$, there are two obvious options. We can cover I by $O(1)$ cubes in $\mathcal{B}(L_N)$, obtaining $\mu(I) \lesssim L_N^\alpha = (L_N/L(I))^\alpha L(I)^\alpha$. Alternatively, we cover I by $O((L(I)/L_{N+1})^d)$ cubes in $\mathcal{B}(L_{N+1})$, so $\mu(I) \lesssim (L(I)/L_{N+1})^d L_{N+1}^\alpha = (L(I)/L_{N+1})^{d-\alpha} L(I)^\alpha$. By considering both covering techniques simultaneously, optimizing for any particular interval length $|I|$, we obtain the tighter bound

$$\begin{aligned} \mu(I) &\lesssim \min\left((L_N/L(I))^\alpha, (L(I)/L_{N+1})^{d-\alpha}\right) L(I)^\alpha \\ &\leq (L_N/L_{N+1})^{\alpha(d-\alpha)/d} L(I)^\alpha \end{aligned}$$

Thus the general estimate $\mu(I) \lesssim |I|^\alpha$ is obtained if $L_N/L_{N+1} = O(1)$, which means L_N can decrease at most exponentially. Unfortunately, this rarely occurs even in the most basic constructions.

2.9 An Epsilon of Room

We get slightly more useful results by giving ourselves an epsilon of room. Factoring in an extra $|I|^\delta$ into the minimization, we find

$$\begin{aligned} \mu(I) &\lesssim \min\left((L_N/L(I))^\alpha, (L(I)/L_{N+1})^{d-\alpha}\right) L(I)^\delta L(I)^{\alpha-\delta} \\ &\leq (L_N/L_{N+1})^{\alpha(d-\alpha)/d} (L_N^\alpha L_{N+1}^{d-\alpha})^{\delta/d} L(I)^{\alpha-\delta} \\ &\leq \left[(L_N/L_{N+1})^{\alpha(d-\alpha)/d} L_N^\delta\right] L(I)^{\alpha-\delta} \end{aligned}$$

Thus if $L_N/L_{N+1} = O_\varepsilon(L_N^{-\varepsilon})$ for every $\varepsilon > 0$, then we obtain $\mu(I) \lesssim_\varepsilon |I|^{\alpha-\varepsilon}$ for each $\varepsilon > 0$, which is good enough to conclude $\dim_{\mathbf{H}}(X) \geq \alpha$. We should expect this bound to be much more versatile than the bound in the last section. The ratios L_N/L_{N+1} are obtained at a single scale, whereas the lengths L_N are obtained from compounding lengths over many, many scales. As such, they should have enough kick to overwhelm the ratio even when ε is arbitrarily small. In particular, this method applies if $L_{N+1} = \Omega(L_N^\alpha)$ for some universal parameter $\alpha \geq 1$. So a polynomial rate of decay is now allowed.

Example. If $L_N = e^{-r_N}$, then $L_N/L_{N+1} = e^{r_{N+1}-r_N}$, and $L_N^{-\varepsilon} = e^{\varepsilon r_{N+1}}$. Thus it suffices to show $r_{N+1} - r_N - \varepsilon r_{N+1} \leq 0$ for suitably large N depending on ε . If r_N

is differentiable in N , accelerating as N increases, and $r'_{N+1} - \epsilon r_N \leq 0$, the mean value theorem implies this bound. Thus, in particular, we may apply the bound if $L_N = \exp(-x^M)$, for any fixed $M > 0$.

Example. Kaleti's method constructs a full dimensional set avoiding translates of itself using the Cantor set method. One can choose $L_N = 1/N! \cdot 10^N$, in which case $L_N/L_{N+1} = 10N$, which is eventually bounded by $L_{N+1}^{-\epsilon} \geq 8^{\epsilon N}$ for any $\epsilon > 0$. The discrete bounds for $I \in \mathcal{B}(L_N)$ are easily established in this case, enabling us to conclude Kaleti's set is full dimensional.

It is, of course, easy to construct superexponential examples of L_N which increase fast enough that we cannot use this technique. For instance, if $L_N = e^{-N!}$, or $L_N = e^{-e^N}$, then these results cannot be applied. In the next section, we provide a method which works when the lengths L_N have an *arbitrary* rate of decay, given additional *uniform* selection assumptions on the Cantor set construction.

2.10 Bounds on Uniform Constructions

Our final method for interpolating requires extra knowledge of the dissection process, but enables us to choose the L_N arbitrarily rapidly, which means the mass of μ can be distributed at lacunary lengths. The idea behind this is that there is an additional sequence of lengths R_N with $L_N \leq R_N \leq L_{N-1}$. The difference between R_N and L_{N-1} is allowed to be arbitrary, but the decay rate between L_N and R_N is polynomial, which enables us to use the covering methods of the previous section. In addition, we rely on a 'uniform mass bound' between R_N and L_N to cover the remaining classes of intervals. Because we can take R_N arbitrarily large relative to L_N , this renders any constants that occur in the construction to become immediately negligible. For two quantities A and B , we will let $A \lesssim_N B$ stand for an inequality with a hidden constant depending only on parameters with index smaller than N , i.e. $A \leq C(L_1, \dots, L_N, R_1, \dots, R_N)B$ for some constant $C(L_1, \dots, L_N, R_1, \dots, R_N)$.

Theorem 12. *If we have discrete bounds $\mu(I) \lesssim_{N-1} L(I)^\alpha$ for $I \in \mathcal{B}(L_N)$, we have a change of scale bound $L_N/R_N \gtrsim_{N-1, \epsilon} R_N^\epsilon$ for all $\epsilon > 0$, and we have a uniform distribution bound $\mu(J) \lesssim_{N-1} (R_{N+1}/L_N)^d \mu(I)$ for $J \subset I$ with $J \in \mathcal{B}(R_{N+1})$ and $I \in \mathcal{B}(L_N)$, then for suitably fast decaying L_N and R_N , $\dim_{\mathbf{H}}(X) \geq \alpha$.*

Proof. If I satisfies $R_{N+1} \leq L(I) \leq L_N$, we can cover I by elements of $\mathcal{B}(R_{N+1})$,

then applying the uniform distribution bound to obtain, for small ε ,

$$\mu(I) \lesssim_{N-1} (L(I)/R_{N+1})^d (R_{N+1}/L_N)^d L_N^\alpha = L(I)^d / L_N^{d-\alpha} \leq L(I)^{\alpha-\varepsilon} L_N^\varepsilon$$

Since the hidden constants only depend on constants before L_N and R_N , we can pick L_N suitably large to give a parameter independent bound. For each N , we pick L_N suitably large such that the bound $\mu(I) \leq |I|^{\alpha-\varepsilon}$ holds for all $\varepsilon \geq 1/N$ when $R_{N+1} \leq |I| \leq L_N$. By doing this, we obtain that $\mu(I) \lesssim_\varepsilon |I|^{\alpha-\varepsilon}$ for all intervals I where there is N such that $R_{N+1} \leq |I| \leq L_N$.

On the other hand, the uniform distribution bound implies $\mu(I) \lesssim_{N-1} |I|^\alpha$ for $I \in \mathcal{B}(R_N)$. If there is N such that $L_N \leq |I| \leq R_N$, this means we apply the epsilon of room trick in the last section to find that for all $\delta > 0$,

$$\mu(I) \lesssim_{N-1} (R_N/L_N)^{\alpha(1-\alpha)/d} R_N^\delta |I|^{\alpha-\delta}$$

Thus if $R_N/L_N \lesssim_{N-1,\varepsilon} R_N^{-\varepsilon}$ for all ε , we can take R_N suitably large to annihilate the constant depending on previous constants in the bound above. \square

Example. In Fraser and Pramanik's paper, for $\alpha = 1/(d-1)$, they choose $R_N = A_N L_{N+1}^\alpha$ for some constant A_N , and choose L_N suitably rapidly decaying to establish the inequalities $\mu_\beta(I) \lesssim |I|^\alpha$ for $|I| = L_N$. The decomposition algorithm naturally leads to the uniform distribution inequality $\mu_\beta(J) \lesssim (R_N/L_N) \mu_\beta(I)$, which shows that we can choose L_N giving an α dimensional set.

Chapter 3

Related Work

3.1 Ruzsa: Difference Sets Without Squares

In this section, we describe the work of Ruzsa on the discrete squarefree difference problem, which provides inspiration for our speculated results for the squarefree subset problem in the continuous setting. If X and Y are subsets of integers, we shall let $X \pm Y = \{x \pm y : x \in X, y \in Y, x \pm y > 0\}$ denote the sums and difference of the set. The *differences* of a set X are elements of $X - X$, and so the squarefree difference set problem asks to consider how large a subset of the integers can be, whose differences do not contain the square of any positive integer. We let $D(N)$ denote the maximum number of integers which can be selected from $[1, N]$ whose differences do not contain a square.

Example. *The set $X = \{1, 3, 6, 8\}$ is squarefree, because $X - X = \{2, 3, 5, 7\}$, and none of these elements are perfect squares. On the other hand, $\{1, 3, 5\}$ is not a squarefree subset, because $5 - 1 = 4$ is a perfect square.*

There are a few tricks to constructing large subsets of integers avoiding squares. If p is prime, then $p\mathbf{Z} \cap [1, p^2)$ avoids squares, because the difference of two numbers must be divisible by p , but not by p^2 . If $N = p^2$, this gives a set with $N^{1/2}$ elements. However, we can do just as well without using any properties of the set of squares except for their sparsity, by greedily applying a sieve. We start by writing out a large list of integers $1, 2, 3, 4, \dots, N$. Then, while we still have numbers to pick, we greedily select the smallest number x_* we haven't crossed out of the list, add it to our set X of squarefree numbers, and then cross out all integers y such that $y - x_*$ is a positive square. Thus we cross out $x_*, x_* + 1, x_* + 4$, and so on, all

the way up to $x_* + m^2$, where m is the largest integer with $x_* + m^2 \leq N$. This implies $m \leq \sqrt{N - x_*} \leq \sqrt{N - 1}$, hence we cross out at most $\sqrt{N - 1} + 1$ integers whenever we add a new element x_* to X . When the algorithm terminates, all integers must be crossed out, and if the algorithm runs n iterations, a union bound gives that we cross out at most $n[\sqrt{N - 1} + 1]$ integers, hence $n[\sqrt{N - 1} + 1] \geq N$. It follows that the set X we end up with contains $\Omega(N^{1/2})$ elements. What's more, this algorithm generates an increasing family of squarefree subsets of the integers as n increases, so we may take the union of these subsets over all N to find an infinite squarefree subset X with $|X \cap [1, N]| = \Omega(\sqrt{N})$ for all N .

In 1978, Sárközy proved an upper bound on the size of squarefree subsets of the integers, showing $D(N) = O(N(\log N)^{-1/3+\varepsilon})$ for every $\varepsilon > 0$. In particular, this proves a conjecture of Lovász that every infinite squarefree subset has density zero, because if X is any infinite squarefree subset, then $|X \cap [1, N]| = o(N)$. Sárközy even conjectured that $D(N) = O(N^{1/2+\varepsilon})$ for all $\varepsilon > 0$. Thus the sieve technique is essentially optimal, an incredibly pessimistic point of view, since the Sieve method doesn't depend on any properties of the set of perfect squares. Ruzsa's results shows we should be more optimistic, taking advantage of the digit expansion of numbers to obtain infinite squarefree subsets X with $|X \cap [1, N]| = \Omega(N^{0.73})$. The method reduces the problem to a finitary problem of maximizing squarefree subsets modulo a squarefree integer m .

Theorem 13. *If m is a squarefree integer, then*

$$D(N) \geq \frac{n^{\gamma_m}}{m} = \Omega_m(n^{\gamma_m})$$

where

$$\gamma_m = \frac{1}{2} + \frac{\log_m |R^*|}{m}$$

and R^* denotes the maximal subset of $[1, m]$ whose differences contain no squares modulo m . Setting $m = 65$ gives

$$\gamma_m = \frac{1}{2} \left(1 + \frac{\log 7}{\log 65} \right) = 0.733077 \dots$$

and therefore $D(N) = \Omega(n^{0.7})$. For $m = 2$, we find $D(N) \geq \sqrt{N}/2$, which is only slightly worse than the sieve result.

Remark. *Let us look at the analysis of the sieve method backwards. Rather than fixing N and trying to find optimal solutions of $[1, N]$, let's fix a particular strategy*

(to start with, the sieve strategy), and think of varying N and seeing how the size of the solution given by the strategy on $[1, N]$ increases over time. In our analysis, the size of a solution is directly related to the number of iterations the strategy can produce before it runs out of integers to add to a solution set. Because we apply a union bound in our analysis, the cost of each particular new iteration is the same as the cost of the other iterations. If the cost of each iteration was independent of N , we could increase the solution size by increasing N by a fixed constant, leading to family of solutions which increases on the order of N . However, as we increase N , the cost of each iteration increases on the order of \sqrt{N} , leading to us only being able to perform $N/\sqrt{N} = \sqrt{N}$ iterations for a fixed N . Rusza's method applies the properties of the perfect squares to perform a similar method of expansion. At an exponential cost, Rusza's method increases the solution size exponentially. The advantage of exponentials is that, since Rusza's is based on a particular parameter, a squarefree integer m , we can vary m to improve the iteration numbers more naturally.

The idea of Rusza's construction is to break the problem into exponentially large intervals, upon which we can solve the problem modulo an integer. More enerally, Rusza constructs a set whose differences are free of d 'th powers.

Theorem 14. *Let $R \subset [1, m]$ be a subset of integers such that no difference is a power of d modulo m , where m is a squarefree integer. Construct the set*

$$A = \left\{ \sum_{k=0}^n r_k m^k : 0 \leq n < \infty, r_k \in \begin{cases} R & d \text{ divides } N \\ [1, m] & \text{otherwise} \end{cases} \right\}$$

Then A is squarefree.

Proof. Suppose that we can write $\sum (r_k - r'_k) m^k = N^d$. Let s to be the smallest index with $r_s \neq r'_s$. Then $(r_s - r'_s) m^s + M m^{s+1} = N^d$ where M is some positive integer. If $s = ds_0$, then $(N/m^{s_0})^d = (r_s - r'_s) + Mm$, and this contradicts the fact that $r_s - r'_s$ cannot be a d 'th power modulo m . On the other hand, we know m^s divides N^d , but m^{s+1} does not. This is impossible if s is not divisible by d , because primes in N^d occur in multiples of d , and m is squarefree. \square

For any n , we find

$$A \cap [1, m^n - 1] = \left\{ \sum_{k=0}^{n-1} r_k m^k : r_k \in [1, m], r_k \in R \text{ when } d \text{ divides } k \right\}$$

which therefore has cardinality

$$|R|^{1+\lfloor \frac{n-1}{d} \rfloor} m^{n-1-\lfloor \frac{n-1}{d} \rfloor} = m^n \left(\frac{|R|}{m} \right)^{1+\lfloor \frac{n-1}{d} \rfloor} \geq m^n \left(\frac{|R|}{m} \right)^{n/d} = m^{n\gamma(m,d)}$$

where $\gamma(m,d) = 1 - 1/d + \log_m |R|/d$. Therefore, for $m^{n+1} - 1 \geq k \geq m^n - 1$

$$A \cap [1, k] \geq A \cap [1, m^n] \geq m^{n\gamma(m,d)} = \frac{m^{(n+1)\gamma(m,d)}}{m} \geq \frac{k^{\gamma(m,d)}}{m}$$

This completes Rusza's construction. Thus we have proved a more general result than was required.

Theorem 15. *For every d and squarefree integer m , we can construct a set X whose differences contain no d th powers and*

$$|X \cap [1, n]| \geq \frac{n^{\gamma(d,m)}}{m} = \Omega(n^{\gamma(d,m)})$$

where $\gamma(d,m) = 1 - 1/d + \log_m |R^*|/d$, and R^* is the largest subset of $[1, m]$ containing no d 'th powers modulo m .

For $m = 65$, the group $\mathbf{Z}_{65}^* \cong \mathbf{Z}_5^* \times \mathbf{Z}_{13}^*$ has a set of squarefree residues of the form $\{(0,0), (0,2), (1,8), (2,1), (2,3), (3,9), (4,7)\}$, which gives the required value for γ_{65} . In 2016, Mikhail Gabdullin proved that if m is squarefree, then in \mathbf{Z}_m , any set R such that $R - R$ is squarefree has $|R| \leq me^{-c \log m / \log \log m}$, where n denotes the number of odd prime divisors of m , so that

$$\gamma(d,m) \leq 1 - 1/d + \frac{\log(me^{-c \log m / \log \log m})}{m}$$

Rusza believes that we cannot choose m to construct squarefree subsets of the integers growing better than $\Omega(n^{3/4})$, and he claims to have proved this assuming m is squarefree and consists only of primes congruent to 1 modulo 4. Looking at some sophisticated papers in number theory (Though I forgot to write down the particular references), it seems that using modern estimates this is quite easy to prove. Thus expanding on Rusza's result in the discrete case requires a new strategy, or perhaps Rusza's result is the best possible.

Let $D(N,d)$ denote the largest subset of $[1, N]$ containing no d th powers of positive integer. The last part of Rusza's paper is devoted to lower bounding the polynomial growth of $D(N,d)$ asymptotically.

Theorem 16. *If p is the least prime congruent to one modulo $2d$, then*

$$\limsup_{N \rightarrow \infty} \frac{\log D(N, d)}{\log N} \geq 1 - \frac{1}{d} + \frac{\log_p d}{d}$$

Proof. The set X we constructed in the last theorem shows that for any m ,

$$\frac{\log D(N, d)}{\log n} \geq \gamma(d, m) - \frac{\log m}{\log n} = 1 - \frac{1}{d} + \frac{\log_m |R^*|}{d} - \frac{\log m}{\log n}$$

Hence

$$\limsup_{N \rightarrow \infty} \frac{\log D(N, d)}{\log n} \geq 1 - \frac{1}{d} + \frac{\log_m |R^*|}{d}$$

The claim is then completed by the following lemma. □

Lemma 10. *If p is a prime congruent to 1 modulo $2d$, then we can construct a set $R \subset [1, p]$ whose differences do not contain a d th power modulo p with $|R| \geq d$.*

Proof. Let $Q \subset [1, p]$ be the set of powers $1^k, 2^k, \dots, p^k$ modulo p . We have

$$|Q| = \frac{p-1}{k} + 1$$

This follows because the nonzero elements of Q are the images of the group homomorphism $x \mapsto x^k$ from \mathbf{Z}_p^* to itself. Since \mathbf{Z}_p^* is cyclic, the equation $x^k = 1$ has the same number of solutions as the equation $kx = 0$ modulo $p-1$, and since $p \equiv 1$ modulo $2k$, there are exactly k solutions to this equation. The sieve method yields a k th power modulo p free subset of size greater than or equal to

$$p/q = \frac{p}{1 + \frac{p-1}{k}} = \frac{pk}{p+k-1} \rightarrow k$$

as $p \rightarrow \infty$, which is greater than $k-1$ for large enough p . This shows the theorem is essentially trivial for large primes. However, for smaller primes a more robust analysis is required. We shall construct a sequence $b_1, \dots, b_k \in \mathbf{Z}_p$ such that $b_i - b_j \notin Q$ for any i, j and $|B_j + Q| \leq 1 + j(q-1)$. Given b_1, \dots, b_j , let b_{j+1} be any element of $(B_j + Q + Q) - (B_j + Q)$. Since $b_{j+1} \notin B_j + Q$, $b_{j+1} - b_i \notin Q$ for any i . Since $b_{j+1} \in B_j + Q + Q$, the sets $B_j + Q$ and $b_{j+1} + Q$ are not disjoint (we have used

$Q = -Q$, which is implied when $p \equiv 1 \pmod{2k}$, and so

$$\begin{aligned} |B_{j+1} + Q| &= |(B_j + Q) \cup (b_{j+1} + Q)| \\ &\leq |B_j + Q| + |b_{j+1} + Q| - 1 \\ &\leq 1 + j(q-1) + q - 1 \\ &= 1 + (j+1)(q-1) \end{aligned}$$

This procedure ends when $B_j + Q + Q = B_j + Q$, and this can only happen if $B_j + Q = \mathbf{Z}_p$, because we can obtain all integers by adding elements of Q recursively, so $1 + j(q-1) \geq p$, and thus $j \geq k$. \square

Corollary. *In the special case of avoiding squarefree numbers, we find*

$$\limsup \frac{\log D(N)}{\log N} \geq \frac{1}{2} + \frac{\log_5 2}{2} = 0.71533\dots$$

which is only slightly worse than the bound we obtain with $m = 65$.

Rusza's leaves the ultimate question of whether one can calculate

$$\alpha = \lim_{N \rightarrow \infty} \log D(N) / \log N$$

or even whether it exists at all. The consequence of this would essentially solve the squarefree integers problem, since it gives the exact growth of $D(N)$ in terms of a monomial. Because of how conclusive this problem is, we should not expect to find a nontrivial way to calculate this constant.

3.2 Keleti: A Translate Avoiding Set

Keleti's two page paper constructs a full dimensional subset X of $[0, 1]$ such that X intersects $t + X$ in at most one place for each nonzero real number t . Malabika has adapted this technique to construct high dimensional subsets avoiding non-trivial solutions to differentiable functions. In this section, and in the sequel, we shall find it is most convenient to avoid certain configurations by expressing them in terms of an equation, whose properties we can then exploit. One feature of translation avoidance is that the problem is specified in terms of a linear equation.

Lemma 11. *A set X avoids translates if and only if there do not exist values $x_1 < x_2 \leq x_3 < x_4$ in X with $x_2 - x_1 = x_4 - x_3$.*

Proof. Suppose $t + X \cap X$ contains two points $a < b$. Without loss of generality, we may assume that $t > 0$. If $a \leq b - t$, then the equation $a - (a - t) = t = b - (b - t)$ satisfies the constraints, since $a - t < a \leq b - t < b$ are all elements of X . We also have $(b - t) - (a - t) = b - a$ which satisfies the constraints if $a - t < b - t \leq a < b$. This covers all possible cases. Conversely, if there are $x_1 < x_2 \leq x_3 < x_4$ in X with $x_2 - x_1 = t = x_4 - x_3$, then $X + t$ contains $x_2 = x_1 + (x_2 - x_1)$ and $x_4 = x_3 + (x_4 - x_3)$. \square

The basic, but fundamental idea to Keleti's technique is to introduce memory into Cantor set constructions. Keleti constructs a nested family of discrete sets $X_0 \supset X_1 \supset \dots$ converging to X , with each X_N a union of disjoint intervals in $\mathcal{B}(L_N)$, for a decreasing sequence of lengths L_N converging to zero, to be chosen later, but with $10L_{N+1} \mid L_N$. We initialize $X_0 = [0, 1]$, and $L_0 = 1$. Furthermore, we consider a queue of intervals, initially just containing $[0, 1]$. To construct X_1, X_2, \dots , Keleti iteratively performs the following procedure:

Algorithm 1 Construction of the Sets X_N

Set $N = 0$

Repeat

Take off an interval I from the front of the queue

For all $J \in \mathcal{B}(L_N)$ contained in X_N :

Order the intervals in $\mathcal{B}(L_{N+1})$ contained in J as J_0, J_1, \dots, J_M

If $J \subset I$, add all intervals J_i to X_{N+1} with $i \equiv 0$ modulo 10

Else add all J_i with $i \equiv 5$ modulo 10

Add all intervals in $\mathcal{B}(L_{N+1})$ to the end of the queue

Increase N by 1

After each iteration of the algorithm, we obtain a new set X_{N+1} , and so leaving the algorithm to repeat infinitely produces a sequence of sets X_1, X_2, \dots converging to a set X . We claim that with the appropriate choice of parameters, X is a translate avoiding set.

If X is not translate avoiding, there is $x_1 < x_2 \leq x_3 < x_4$ with $x_2 - x_1 = x_4 - x_3$. Since $L_N \rightarrow 0$, there is N such that x_1 is contained in an interval $I \in \mathcal{B}(L_N)$ that x_2, x_3, x_4 are not contained in. At stage N of the algorithm, the interval I is added to the end of the queue, and at a much later stage M , the interval I is retrieved. Find the startpoints $x_1^\circ, x_2^\circ, x_3^\circ, x_4^\circ \in L_M \mathbb{Z}$ to the intervals in $\mathcal{B}(L_M)$ containing x_1, x_2, x_3 , and x_4 . Then we can find n and m such that $x_4^\circ - x_3^\circ = (10n)L_M$, and $x_2^\circ - x_1^\circ =$

$(10m+5)L_M$. In particular, this means that $|(x_4^\circ - x_3^\circ) - (x_2^\circ - x_1^\circ)| \geq 5L_M$. But

$$\begin{aligned} |(x_4^\circ - x_3^\circ) - (x_2^\circ - x_1^\circ)| &= |[(x_4^\circ - x_3^\circ) - (x_2^\circ - x_1^\circ)] - [(x_4 - x_3) - (x_2 - x_1)]| \\ &\leq |x_1^\circ - x_1| + \dots + |x_4^\circ - x_4| \leq 4L_M \end{aligned}$$

which gives a contradiction.

The algorithm shows that X_N contains $L_{N-1}/10L_N$ times the number of intervals that X_{N-1} has, but they are at a length L_N rather than L_{N-1} . This means that in total, X_N contains $1/10^N L_N$ intervals, of length L_N . Since $L_N/10^N L_N = o(1)$, this shows our set will have Lebesgue measure zero irrespective of our parameters. However, if L_N decays suitably fast, then we might have $L_N^{1-\varepsilon}/10^N L_N \gtrsim_\varepsilon 1$ for all $\varepsilon > 0$, which would imply that X has positive $1 - \varepsilon$ dimensional Hausdorff measure for all ε , so X still has Hausdorff dimension one. For this to be true, L_N must decay superexponentially, i.e. the inequalities above are equivalent to $L_N \lesssim_B 1/B^N$ for all choices of B . Choosing an arbitrarily fast decaying sequence, such as $L_N = 1/N! \cdot 10^N$ or $L_N = 1/10^{10^N}$, suffices to obtain a Hausdorff dimension one set.

Lemma 12. *If L_N decays superexponentially, X has Hausdorff dimension one.*

Proof. We use the mass distribution principle, as used in our note on calculating Hausdorff dimensions. It is easy to establish the bounds $\mu_N(I) \lesssim_\varepsilon L(I)^{1-\varepsilon}$ for $I \in \mathcal{B}(L_N)$, and since we can choose L_N suitably slowly decreasing to use the epsilon of room technique, this gives the result. Alternatively, we can use the uniform distribution bounds with $L_N = R_N$, since if $J \in \mathcal{B}(L_{N+1})$, $I \in \mathcal{B}(L_N)$, $\mu(J) = 1/10^{N+1} L_{N+1}$, $\mu(I) = 1/10^N L_N$, and so $\mu(J) \lesssim (L_{N+1}/L_N)\mu(I)$. This gives the result if L_N grows too fast. \square

Remark. *Keleti briefly remarks that by replacing the 10 in the algorithm with a slowly increasing set of numbers, one can obtain a Hausdorff dimension one set which is linearly independent over the rational numbers. To see why this works, the condition of linear independence would fail if $a_1 x_1 + \dots + a_M x_M = 0$, where $x_1 < x_2 < \dots < x_M$, and a_1, \dots, a_M are integers with no common factor. One can again reduce this by picking intervals with indices congruent to a certain large modulus.*

3.3 Fraser/Pramanik: Extending Keleti Translation to Smooth Configurations

Inspired by Keleti's result, Pramanik and Fraser obtained a generalization of the queue method which allows one to find sets avoiding solutions to *any* smooth function satisfying suitably mild regularity conditions. To do this, rather than making a linear shift in one of the intervals we avoid as in Keleti's approach, one must use the smoothness properties of the function to find large segments of an interval avoiding solutions to another interval.

Theorem 17. *Suppose that $f : \mathbf{R}^{d+1} \rightarrow \mathbf{R}$ is a C^1 function, and there are sets $T_0, \dots, T_d \subset [0, 1]$, with each T_n a union of almost disjoint closed intervals of length $1/M$ such that $A \leq |\partial_0 f|$ and $|\nabla f| \leq B$ on $T_0 \times \dots \times T_d$. Then there exists a rational constant C and arbitrarily large integers $N \in M\mathbf{Z}$ for which there exist subsets $S_n \subset T_n$ such that*

- (i) $f(x) \neq 0$ for $x \in S_0 \times \dots \times S_d$.
- (ii) For $n \neq 0$, if we divide each interval T_n into length $1/N$ intervals, then S_n contains an interval of length C/N^d of each of these intervals.
- (iii) If T_0 is split into length $1/N$ intervals, then for a fraction $1 - 1/M$ of such intervals, S_0 is a union of length C/N^d intervals with total length C/N .

Proof. We begin by dividing the sets T_1, \dots, T_d into length $1/N$ intervals, and let S_n be defined by including a length C_0/N^d segment, for some constant C_0 to be chosen later. Then once we fix C_0 , the S_n will satisfy property (ii) of the theorem. We define

$$\mathbf{A} = \{a \in \mathbf{R}^{d-1} : a_n \text{ is a startpoint of a length } 1/N \text{ interval in } T_n\}$$

Then $|\mathbf{A}| \leq N^d$, since each interval T_n is contained in $[0, 1]$, and therefore can only contain at most N almost disjoint intervals of length $1/N$. Hence if we define the set of 'bad points' in T_0 as

$$\mathbf{B} = \{x \in T_0 : \text{there is } a \in \mathbf{A} \text{ such that } f(x, a) = 0\}$$

Then $|\mathbf{B}| \leq MN^d$. This is because for each fixed a , the function $x \mapsto f(x, a)$ is either strictly increasing or decreasing over each interval in the decomposition of T_0 , or which there are at most M because $T_0 \subset [0, 1]$. If we split T_0 into length $1/N$

intervals, and choose a subcollection of such intervals I such that $|I \cap \mathbf{B}| \leq M^3 N^{d-1}$, then we throw away at most $MN^d / M^3 N^{d-1} = N/M^2$ intervals, and so we keep $(N/M)(1 - 1/M)$ intervals, which is $1 - 1/M$ of the total number of intervals in the decomposition of T_0 . The lemma we prove after this theorem implies that there exists a constant C_1 such that if $x \in S_n$, and $f(y, x) = 0$, then $d(y, \mathbf{B}) \leq C_0 C_1 / N^d$. If we split each interval I with $|I \cap \mathbf{B}| \leq M^3 N^{d-1}$ into $4M^3 N^{d-1}$ length $1/4M^3 N^d$ intervals, and we choose C_0 such that $C_0 C_1 < 1/4M^3$, then the set S_0 obtained by discarding each interval that contains or is adjacent to an interval containing an element of \mathbf{B} satisfies $d(S_0, \mathbf{B}) > C_0 C_1 / N^d$, and therefore there does not exist any $x_n \in S_n$ and $y \in S_0$ such that $f(y, x) = 0$. S_0 satisfies property (iii) of the theorem since for the interval I we are considering, we keep at least $M^3 N^{d-1}$ length $1/4M^3 N^d$ intervals, which in total has length at least $1/4N$. \square

Remark. *The length $1/N$ portion of each interval guaranteed by (iii) is unnecessary to the Hausdorff dimension bound, since the slightly better bounds obtained on scales where an interval is dissected as a $1/N$ are decimated when we eventually divide the further subintervals into $1/N^{d-1}$ intervals. The importance of (iii) is that it implies that the set we will construct has full Minkowski dimension. The reason for this is that Minkowski dimension lacks the ability to look at varying dissection depths at once, and since, at any particular depth, there exists a length $1/N$ dissection, the process appears to Minkowski to be full dimensional, even though at later scales this $1/N$ dissection is dissected into $1/N^{d-1}$ intervals.*

Lemma 13. *Given the f, T_0, \dots, T_d , there exists a constant C_1 depending on these quantities, such that for any C_0 , and $x \in S_1 \times \dots \times S_{d-1}$, if $f(y, x) = 0$, then $d(y, \mathbf{B}) \leq C_0 C_1 / N^{d-1}$.*

Proof. Since $T_0 \times \dots \times T_d$ breaks into finitely many cubes with sidelengths $1/M$, it suffices to prove the theorem for a particular cube J in this decomposition, where we assume the zeroset of f intersects J . If $J = I \times J'$, where I is an interval, we let U be the set of all $x \in J'$ for which there is y in the interior of I such that $f(y, x) = 0$. Then U is open. The implicit function theorem implies that there exists a C^1 function $g : U \rightarrow I$ such that $f(x, y) = 0$ if and only if $y = g(x)$. Then the function $h(x) = f(x, g(x))$ vanishes uniformly, so

$$0 = \partial_n h(x) = (\partial_n f)(g(x), x) + (\partial_0 f)(g(x), x) \partial_n g(x)$$

Hence for $x \in U$,

$$|(\nabla g)(x)| = \frac{|(\nabla f)(x)|}{|(\partial_d f)(x, g(x))|} \leq \frac{B}{A}$$

If N is chosen large enough, then for every $x \in U \cap (S_1 \times \cdots \times S_d)$ there is $a \in \mathbf{A} \cap U$ in the same connected component of U as x with $|x - a| \lesssim C_0/N^{d-1}$, and this means that

$$|g(x) - g(a)| \leq \|\nabla g\|_\infty |x - a| \lesssim \frac{BC_0}{AN^{d-1}}$$

and $g(a) \in \mathbf{B}$, completing the proof. \square

How do we use this lemma to construct a set avoiding solutions to f ? We form an infinite queue which will eventually filter out all the possible zeroes of the equation. Divide the interval $[0, 1]$ into d intervals, and consider all orderings of $d - 1$ subsets of these intervals, and add them to the queue. Now on each iteration N of the algorithm, we have a set $X_N \subset [0, 1]$. We take a particular sequence of intervals T_1, \dots, T_d from the queue, and then use the lemma above to dissect the $X_N \cap T_n$, which are unions of intervals, into sets avoiding solutions to the equation, and describe the remaining points as X_{N+1} . We then add all possible orderings of d intervals created into the end of the queue, and rinse and repeat. The set $X = \lim X_n$ then avoids all solutions to the equation with distinct inputs.

What remains is to bound the Hausdorff dimension of X by constructing a probability measure supported on X with suitable decay. To construct our probability measure, we begin with a uniform measure on the interval, and then, whenever our interval is refined, we uniformly distribute the volume on that particular interval uniformly over the new refinement. Let μ denote the weak limit of this sequence of probability distributions. At each step n of the process, we let $1/M_n$ denote the size of the intervals at the beginning of the n 'th subdivision, $1/N_n$ denote the size of the split intervals in the lemma, and C_n the n 'th constant. We have the relation $1/M_{n+1} = C_n/N_n^{d-1}$. If K is a length $1/M_{N+1}$ interval, J a length $1/N_N$ interval, and I a length $1/M_N$ interval with $K \subset J \subset I$ and all receiving some mass in μ . To calculate a bound on their mass, we consider the decompositions considered in the algorithm:

- If J is subdivided in the non-specialized manner, then every length $1/N_N$ interval receives the same mass, which is allocated to a single length $1/M_{N+1}$ interval it contains. Thus $\mu(K) = \mu(J) \leq (M_N/N_N)\mu(I)$.
- In the second case, at least a fraction $1 - 1/M_N$ of the length $1/N_N$ intervals are assigned mass, so $\mu(J) = (M_N/N_N)(1 - 1/M_N)^{-1} \leq (2M_N/N_N)\mu(I)$, and more than C_N/N_N of each length $1/N_N$ interval is maintained, so

$$\mu(K) = \frac{N_N}{C_N M_{N+1}} \mu(J) \leq \frac{2M_N}{C_N M_{N+1}} \mu(I)$$

Thus in both cases, we have $\mu(J) \lesssim (N_N/M_N)\mu(I)$, $\mu(K) \lesssim_N |K|$, and $N_N = M_{N+1}^{1/(d-1)}/C_N \lesssim_N M_{N+1}^{1/(d-1)}$. From this, we conclude using the results of the appendix that there exists a family of rapidly decaying parameters which gives a $1/(d-1)$ dimensional set.

Remark. *The set X constructed is precisely a $1/(d-1)$ dimensional set. Recall that $X = \lim X_n$, where X_n is a union of a certain number of length $1/M_n$ intervals I_1, \dots, I_N . For each n , the interval I_i is inevitably subdivided at a stage J_i into length $C_{J_i} N_{J_i}^{1-d}$ intervals for each length $1/N_{J_i}$ interval that I_i contains. Thus*

$$H_{1/M_n}^\alpha(X) \leq \sum_{i=1}^N \frac{N_{m_i}}{M_n} (C_{m_i} N_{m_i}^{1-d})^\alpha = \frac{1}{M_n} \sum_{i=1}^N C_{m_i}^\alpha N_{m_i}^{1-\alpha(d-1)}$$

We may assume that $C_{m_i} \leq 1$, so if $\alpha > 1/(d-1)$, using the fact that $N \leq M_n$, since X_n is contained in $[0, 1]$, we obtain

$$H_{1/M_n}^\alpha(X) \leq \frac{1}{M_n} \sum_{i=1}^N N_{m_i}^{1-\alpha(d-1)} \leq N_{\max(m_i)}^{1-\alpha(d-1)} \leq 1$$

Thus, taking $n \rightarrow \infty$, we conclude $H^\alpha(X) \leq 1 < \infty$, so as $\alpha \downarrow 1/(d-1)$, we conclude that X has Hausdorff dimension bounded above by $1/(d-1)$.

3.4 A Set Avoiding All Functions With A Common Derivative

In the latter part's of their paper, Pramanik and Fraser apply an iterative technique to construct, for each α with $\sum \alpha_n = 0$ and $K > 0$, a set E of positive Hausdorff dimension avoiding solutions to any function $f: \mathbf{R}^d \rightarrow \mathbf{R}$ satisfying with $(\partial_n f)(0) = \alpha_n$,

$$|f(x) - \sum \alpha_n x_n| \leq K \sum_{n \neq 1} (x_n - x_1)^2$$

The set of such f is an uncountable family, which makes this situation interesting. The technique to create such a set relies on another iterative procedure.

Lemma 14. *Let $I \not\subseteq [1, d]$ be a strict subset of indices, and $\delta_0 > 0$. Then there exists $\varepsilon > 0$ such that for any $\lambda > 0$ and two disjoint intervals J_1 and J_2 , with J_1 occuring*

before J_2 , and if we set

$$[a_n, b_n] = \begin{cases} J_1 & n \in I \\ J_2 & n \notin I \end{cases}$$

then for $\delta < \delta_0$, either for all $x_n \in [a_n, a_n + \varepsilon\lambda]$ or for all $x_n \in [b_n - \varepsilon\lambda, b_n]$,

$$|\sum \alpha_n x_n| \geq \delta\lambda$$

Proof. If $C^* = \sum |\alpha_n|$, then for $|x_n - a_n| \leq \varepsilon\lambda$,

$$|\sum \alpha_n (x_n - a_n)| \leq C^* \varepsilon\lambda$$

Thus if $|\sum \alpha_n a_n| > (\delta + \varepsilon C^*)\lambda$, then $|\sum \alpha_n x_n| \geq \delta\lambda$. If this does not occur □

Chapter 4

Avoiding Rough Sets

A major question in modern geometric measure theory is whether sets with sufficiently large Hausdorff dimension necessarily contain copies of certain patterns. For example, Theorem 6.8 of [12] shows that each set $X \subset \mathbf{R}^d$ with Hausdorff dimension exceeding one must contain three colinear points. On the other hand, the main result of [10] constructs a set $X \subset \mathbf{R}^2$ with full Hausdorff dimension such that no four points in X form the vertices of a parallelogram. At the time of this writing, many fundamental questions about the relation between the geometric patterns of a set and its fractional dimension remain open. The threshold dimension above which patterns are guaranteed can vary for different patterns, or not even exist at all. In this paper, we provide lower bounds for the threshold by solving the pattern avoidance problem; constructing sets with large dimension avoiding patterns.

One natural way to approach the pattern avoidance problem is to find general methods for constructing pattern avoiding sets by exploiting a particular geometric feature of the pattern. In [11], Máthé shows that, given a pattern specified by a countable union of rational coefficient bounded degree polynomials f_1, f_2, \dots in nd variables, one can construct a set $X \subset \mathbf{R}^d$ such that for every collection of distinct points x_1, \dots, x_n , and each index k , $f_k(x_1, \dots, x_n) \neq 0$. In [6], Fraser and the second author consider a related problem where the polynomials f_k are replaced with full rank C^1 functions.

In this paper, we also find general methods for constructing pattern avoid sets. But rather than avoiding the zeroes of a function, we fix $Z \subset \mathbf{R}^{dn}$, and construct sets X such that for any distinct $x_1, \dots, x_n \in X$, $(x_1, \dots, x_n) \notin Z$. We then say X avoids the pattern specified by Z . For instance, if $Z = \{(x_1, x_2, x_3) \in (\mathbf{R}^d)^3 : (x_1, x_2, x_3) \text{ are colinear}\}$, then $X \subset \mathbf{R}^d$ avoids the pattern specified by Z precisely when X does not contain three colinear points. Our problem generalizes the prob-

lem statements considered in [11] and [6] if we set $Z = \bigcup_{k=1}^{\infty} f_k^{-1}(0)$. From this perspective, Máthé constructs sets avoiding a set Z formed from a countable union of algebraic varieties, while Fraser and the second author construct sets avoiding a set Z formed from the countable union of C^1 manifolds.

The advantage of the formulation of pattern avoidance we consider is that it is now natural to consider ‘rough’ sets $Z \subset \mathbf{R}^{dn}$ which are not naturally specified as the zero set of a function. In particular, in this paper we consider Z formed from the countable union of sets, each with lower Minkowski dimension bounded above by some α . In contrast with previous work, no further assumptions are made on Z . The dimension of the avoiding set X we will eventually construct depends only on the codimension $nd - \alpha$ of the set Z , and the number of variables n .

Theorem 18. *Let $Z \subset \mathbf{R}^{nd}$ be a countable union of compact sets, each with lower Minkowski dimension at most α , with $d \leq \alpha < dn$. Then there exists a set $X \subset [0, 1]^d$ with Hausdorff dimension at least $(nd - \alpha)/(n - 1)$ such that whenever $x_1, \dots, x_n \in X$ are distinct, then $(x_1, \dots, x_n) \notin Z$.*

Remark. *When $\alpha < d$, the pattern avoidance problem is trivial, since $X = [0, 1]^d - \pi(Z)$ is full dimensional and solves the pattern avoidance problem, where $\pi(x_1, \dots, x_n) = x_1$ is a projection map from \mathbf{R}^{dn} to \mathbf{R}^d . The case $\alpha = dn$ is trivial as well, since we can set $X = \emptyset$.*

When Z is a countable union of smooth manifolds, Theorem 18 generalizes Theorem 1.1 and 1.2 from [6]. Section 4.6 is devoted to a comparison of our methods with [6], as well as other general pattern avoidance methods. But our result is more interesting when it can be applied to truly ‘rough’ sets. One surprising application to our method is obtained by considering a ‘rough’ set Y in addition to a set Z , and finding a set $X \subset Y$ of large dimension avoiding Z . Previous methods in the literature fundamentally exploit the ability to select points on the entirety of Euclidean space, and so it remains unlikely that we can obtain any result of this form using their methods, unless Y is suitably flat, i.e. a smooth surface. To contrast this, our result even applies for certain sets Y which are of Cantor type. We discuss applications of our result in Section 5.

Theorem 18 is proved using a Cantor-type construction, a common theme in the surrounding literature, described explicitly in Section 4.3. For a sequence of decreasing lengths $\{l_n\}$ with $l_n \rightarrow 0$, the construction specifies a selection mechanism for a nested family of sets $X_n \rightarrow X$, with X_n a union of sidelength l_n cubes

and avoiding Z at scales close to l_n . Our *key* contribution to this process is using the probabilistic method to guarantee the existence of efficient selections at each scale, described in Section 2. This selection also assigns mass uniformly on intermediate scales, akin to [6], ensuring that the selection procedure at a single scale is the sole reason for the Hausdorff dimension we calculate in Section 4. Furthermore, the randomization allows us to avoid the complicated queueing techniques in [9] and [6], which has the additional benefit that we can confront the entire behaviour of Z at scales close to l_n simultaneously during the n th step of the construction.

4.1 Frequently Used Notation and Terminology

Our argument heavily depends upon discretizing sets into unions of cubes. Throughout our argument, we use the following notation:

- (A) A *dyadic length* is a length l equal to 2^{-k} for some non-negative integer k .
- (B) Given a length $l > 0$, we let \mathcal{B}_l^d denote the family of all half open cubes in \mathbf{R}^d with sidelength l and corners on the lattice $(l \cdot \mathbf{Z})^d$. That is,

$$\mathcal{B}_l^d = \{[a_1, a_1 + l] \times \cdots \times [a_d, a_d + l] : a_k \in l \cdot \mathbf{Z}\}.$$

If $E \subset \mathbf{R}^d$, $\mathcal{B}_l^d(E)$ is the family of cubes in \mathcal{B}_l^d intersecting E , i.e.

$$\mathcal{B}_l^d(E) = \{I \in \mathcal{B}_l^d : I \cap E \neq \emptyset\}.$$

- (C) The *lower* and *upper Minkowski dimension* of a compact set $Z \subset \mathbf{R}^d$ are defined as

$$\underline{\dim}_{\mathbf{M}}(Z) = \liminf_{l \rightarrow 0} \frac{\log(\#\mathcal{B}_l^d(Z))}{\log(1/l)} \quad \text{and} \quad \overline{\dim}_{\mathbf{M}}(Z) = \limsup_{l \rightarrow 0} \frac{\log(\#\mathcal{B}_l^d(Z))}{\log(1/l)}.$$

If $\underline{\dim}_{\mathbf{M}}(Z) = \overline{\dim}_{\mathbf{M}}(Z)$, then we define $\dim_{\mathbf{M}}(Z)$ equal to the common value.

- (D) If $0 \leq \alpha$ and $\delta > 0$, we define the dyadic Hausdorff content of a set $E \subset \mathbf{R}^d$ as

$$H_\delta^\alpha(E) = \inf \left\{ \sum_{k=1}^m l_k^\alpha : E \subset \bigcup_{k=1}^m I_k \text{ and } I_k \in \mathcal{B}_{l_k}^d, l_k \leq \delta \text{ for all } k \right\}.$$

The α -dimensional dyadic Hausdorff measure H^α on \mathbf{R}^d is $H^\alpha(E) = \lim_{\delta \rightarrow 0} H_\delta^\alpha(E)$, and the *Hausdorff dimension* of a set E is $\dim_{\mathbf{H}}(E) = \inf\{\alpha \geq 0 : H^\alpha(E) = 0\}$.

- (E) Given $I \in \mathcal{B}_l^{dn}$, we can decompose I as $I_1 \times \cdots \times I_n$ for unique cubes $I_1, \dots, I_n \in \mathcal{B}_l^d$. We say I is *strongly non-diagonal* if the cubes I_1, \dots, I_n are distinct. Strongly non-diagonal cubes will play an important role in Section 4.2, when we solve a discrete version of Theorem 18.
- (F) Adopting the terminology of [8], we say a collection of sets $\{U_k\}$ is a *strong cover* of a set E if $E \subset \limsup U_k$, which means every element of E is contained in infinitely many of the sets U_k . This idea will be useful in Section 4.3.
- (G) A *Frostman measure* of dimension α is a non-zero compactly supported probability measure μ on \mathbf{R}^d such that for every cube I of sidelength l , $\mu(I) \lesssim l^\alpha$. Note that a measure μ satisfies this inequality for every cube I if and only if it satisfies the inequality for cubes whose sidelengths are dyadic lengths. *Frostman's lemma* says that

$$\dim_{\mathbf{H}}(E) = \sup \left\{ \alpha : \begin{array}{l} \text{there is an } \alpha \text{ dimensional Frostman} \\ \text{measure supported on } E \end{array} \right\}.$$

4.2 Avoidance at Discrete Scales

In this section we describe a method for avoiding Z at a single scale. We apply this technique in Section 4.3 at many scales to construct a set X avoiding Z at all scales. This single scale avoidance technique is the core part of our construction, and the efficiency with which we can avoid Z at a single scale has direct consequences on the Hausdorff dimension of the set X obtained in Theorem 18.

At a single scale, we solve a discretized version of the problem, where all sets are unions of cubes at two dyadic lengths $l \geq s$. In the discrete setting, Z is replaced by a discretization version of itself, i.e. a union of cubes in \mathcal{B}_s^{dn} , denoted by Z_s . Given a set E , which is a union of cubes in \mathcal{B}_l^d , our goal is to construct a union of cubes in \mathcal{B}_s^d , denoted F , such that $F \subset E$ and F^n is disjoint from the strongly non-diagonal cubes of Z_s (see Definition (E)).

In order to ensure the final set X obtained in Theorem 18 has large Hausdorff dimension regardless of the rapid decay of scales used in the construction of X ,

it is crucial that F is spread uniformly over E . We achieve this by decomposing E into sub-cubes in \mathcal{B}_r^d for some intermediate scale $r \in [s, l]$, and distributing F evenly among as many of these intermediate sub-cubes as possible. Assuming a mild regularity condition on the volume of Z_s , this is possible.

Lemma 15. *Fix two dyadic lengths $l \geq s$. Let E be a nonempty union of cubes in \mathcal{B}_l^d , and let Z_s be a union of cubes in \mathcal{B}_s^d such that $|Z_s| \leq l^{dn}/2$. Then there exists a dyadic length $r \in [s, l]$ such that*

$$A_l |Z_s|^{1/d(n-1)} \leq r \leq \max(s, 2A_l |Z_s|^{1/d(n-1)}) \quad \text{where} \quad A_l = (2^{1/d}/l)^{1/(n-1)}. \quad (4.1)$$

For this scale, there exists a set $F \subset E$, which is a nonempty union of cubes in \mathcal{B}_s^d , satisfying the following three properties:

- (A) Avoidance: For any distinct $J_1, \dots, J_n \in \mathcal{B}_s^d(F)$, $J_1 \times \dots \times J_n \notin \mathcal{B}_s^{dn}(Z_s)$.
- (B) Non-Concentration: For any $I \in \mathcal{B}_r^d(E)$, there is at most one $J \in \mathcal{B}_s^d(F)$ with $J \subset I$.
- (C) Large Size: For any $I \in \mathcal{B}_l^d(E)$, $\#\mathcal{B}_s^d(F \cap I) \geq \#\mathcal{B}_r^d(I)/2 = (l/r)^d/2$.

In other words, F avoids strongly non-diagonal cubes in Z_s , and contains a single sidelength s portion of more than half of the sidelength r cubes contained in any sidelength l cube in E .

Proof. Let r be the smallest dyadic length larger than s and $A_l |Z_s|^{1/d(n-1)}$, so that (4.1) is satisfied. The assumption that $|Z_s| \leq l^{dn}/2$ implies

$$A_l |Z_s|^{1/d(n-1)} \leq A_l l^{n/(n-1)} / 2^{1/d(n-1)} = l.$$

Thus we have guaranteed that $r \in [s, l]$. For each $I \in \mathcal{B}_r^d(E)$, let J_I be a random element of $\mathcal{B}_s^d(I)$ chosen uniformly at random, independantly from the other random variables $J_{I'}$ with $I \neq I'$. Define a random set

$$U = \bigcup \{J_I : I \in \mathcal{B}_r^d(E)\},$$

and for each U , set

$$\mathcal{K}(U) = \{K \in \mathcal{B}_s^{dn}(Z_s) : K \in U^n, K \text{ strongly non-diagonal}\}$$

Then set

$$F_U = U - \{\pi(K) : K \in \mathcal{K}(U), K \text{ is strongly diagonal}\}, \quad (4.2)$$

where $\pi : \mathbf{R}^{dn} \rightarrow \mathbf{R}^d$ maps $K_1 \times \dots \times K_n$ to K_1 for each $K_i \in \mathbf{R}^d$. Given any strongly non-diagonal cube $J_1 \times \dots \times J_n \in \mathcal{B}_s^{dn}(Z_s)$, either $J_1 \times \dots \times J_n \notin \mathcal{B}_s^{dn}(U^n)$, or $J_1 \times \dots \times J_n \in \mathcal{B}_s^{dn}(U^n)$. If the former occurs then $J_1 \times \dots \times J_n \notin \mathcal{B}_s^{dn}(F_U^n)$ since $F_U \subset U$, while if the latter occurs then $K \in \mathcal{K}(U)$, so $J_1 \notin \mathcal{B}_s^d(F_U)$. In either case, $J_1 \times \dots \times J_n \notin \mathcal{B}_s^{dn}(F_U^n)$, so F_U satisfies Property (A). By construction, U contains at most one subcube $J \in \mathcal{B}_s^{dn}$ for each $I \in \mathcal{B}_l^{dn}(E)$. Since $F_U \subset U$, F_U satisfies Property (B). These properties are satisfied for any instance of U , but it is not true that Property (C) holds for each F_U . The remainder of this proof is devoted to showing this property holds for F_U with non-zero probability, guaranteeing the existence of the required set F satisfying the properties of the lemma.

For each cube $J \in \mathcal{B}_s^d(E)$, there is a unique ‘parent’ cube $I \in \mathcal{B}_r^d(E)$ such that $J \subset I$. Since I contains $(r/s)^d$ elements of $\mathcal{B}_s^d(E)$, and J_I is chosen uniformly at random from $\mathcal{B}_s^d(I)$,

$$\mathbf{P}(J \subset U) = \mathbf{P}(J_I = J) = (s/r)^d.$$

The cubes J_I are chosen independantly, so if J_1, \dots, J_k are distinct cubes in $\mathcal{B}_s^d(E)$, then the last calculation combined with Property (B) shows that

$$\mathbf{P}(J_1, \dots, J_k \in U) = \begin{cases} (s/r)^{dk} & : \text{if } J_1, \dots, J_k \text{ have distinct parents} \\ 0 & : \text{otherwise} \end{cases}$$

Let $K = J_1 \times \dots \times J_n \in \mathcal{B}_s^{dn}(Z_s)$ be a strongly non-diagonal cube. Then cubes J_1, \dots, J_n are distinct, so the calculation we just performed implies

$$\mathbf{P}(K \subset U^n) = \mathbf{P}(J_1, \dots, J_n \in U) \leq (s/r)^{dn}.$$

Together with linearity of expectation, and (4.1), if K ranges over the strongly non-diagonal cubes of $\mathcal{B}_s^{dn}(Z_s)$, we find

$$\begin{aligned} \mathbf{E}(\#\mathcal{K}(U)) &= \sum_K \mathbf{P}(K \subset U^n) \leq \#\mathcal{B}_s^{dn}(Z_s) \cdot (s/r)^{dn} = |Z_s| r^{-dn} \\ &= \left[|Z_s| r^{-d(n-1)} \right] r^{-d} \leq \left[|Z_s| (A_l |Z_s|^{1/d(n-1)})^{-d(n-1)} \right] r^{-d} = \left[l^d / 2 \right] r^{-d} = (l/r)^d / 2. \end{aligned}$$

Thus there exists at least one (non-random) set U_0 such that

$$\#\mathcal{K}(U_0) \leq \mathbf{E}(\#\mathcal{K}(U)) \leq (l/r)^d / 2. \quad (4.3)$$

This means that the resultant set F_{U_0} is obtained by removing at most $(l/r)^d / 2$ cubes in \mathcal{B}_s^d from U_0 . Since U_0 contains $(l/r)^d$ cubes in $\mathcal{B}_s^d(I)$ for each $I \in \mathcal{B}_l^d(E)$,

F_{U_0} contains at least $(l/r)^d/2$ cubes in $\mathcal{B}_s^d(I)$ for each $I \in \mathcal{B}_l^d(E)$, so F_{U_0} satisfies Property (C). Setting $F = F_{U_0}$ completes the proof. \square

Remark. While the existence of the set F in Lemma 15 was obtained by probabilistic techniques, we emphasize that its existence is a purely deterministic statement. One can find a candidate F constructively by checking all of the finitely many possible choice of U to find one particular choice U_0 which satisfies (4.3), and then defining F by (4.2). Thus the set we obtain in Theorem 18 exists by purely constructive means.

Our inability to select almost every cube in Lemma 15 means that repeated applications of the result will lead to a loss in Hausdorff dimension. In fact, in the worst case, applying the lemma causes us to lose as much Hausdorff dimension as is permitted by Theorem 1. Note that the value r specified in Theorem 15 is always bounded below by $|Z_s|$. We will later see that if Z is the countable union of sets with lower Minkowski dimension α , the scale s discretization Z_s satisfies $|Z_s| \leq s^{dn-\alpha-\varepsilon}$, for some small positive ε converging to zero as $s \rightarrow 0$. But we can certainly find sets Z with lower Minkowski dimension α satisfying $|Z_s| \geq s^{dn-\alpha}$ for all s . In this situation, (4.1) shows

$$r \geq A_l |Z_s|^{1/d(n-1)} \geq A_l s^{(dn-\alpha)/d(n-1)} \geq s^{(dn-\alpha)/d(n-1)} \quad (4.4)$$

If we combine this inequality with Property (B) of Lemma 15, we conclude

$$\frac{\log \# \mathcal{B}_s^d(F)}{\log(1/s)} \leq \frac{\log \# \mathcal{B}_r^d(E)}{\log(1/s)} = \frac{\log |E| r^{-d}}{\log(1/s)} = \frac{d \log(1/r) - \log |E|^{-1}}{\log(1/s)} \leq \frac{dn - \alpha}{n - 1}.$$

Given a set $F = \bigcap F_k$, where $\{F_k\}$ are infinitely many sets obtained from an application of Lemma 15 at a sequence of scales $\{s_k\}$ with $s_k \rightarrow 0$ as $k \rightarrow \infty$, and where $|Z_{s_k}| \geq s_k^{dn-\alpha}$, then the last computation shows

$$\dim_{\mathbf{H}}(F) \leq \overline{\dim}_{\mathbf{M}}(F) \leq \lim_{s_k \rightarrow 0} \frac{\log \# \mathcal{B}_{s_k}^d(F_k)}{\log(1/s_k)} \leq \frac{dn - \alpha}{n - 1}.$$

This is why we can only lower the Hausdorff dimension of the set X obtained in Theorem 18 by $(dn - \alpha)/(n - 1)$. Furthermore, we see that we must be very careful to ensure applications of the discrete lemma are the only place in our proof where dimension is lost.

Remark. Lemma 15 is the core method in our avoidance technique. The remain-

ing argument is fairly modular. If, for a special case of Z , one can improve the result of Lemma 15 so that r is chosen on the order of $s^{\beta/d}$, then the remaining parts of our paper can be applied near verbatim to yield a set X with Hausdorff dimension β , as in Theorem 18. The last paragraph shows that when $|Z_s| \geq s^{dn-\alpha}$, which is certainly possible given the hypothesis of Theorem 15, the length r is chosen on the order of $s^{(dn-\alpha)/d(n-1)}$, as we saw in (4.4), which is why we obtain a Hausdorff dimension $(dn - \alpha)/(n - 1)$ set.

4.3 Fractal Discretization

In this section we will construct the set X by applying Lemma 15 at many scales. Since Z is a countable union of compact sets with Minkowski dimension at most α , there exists a strong cover (see Definition (F)) of Z by cubes restricted to a sequence of dyadic lengths $\{l_k\}$. We will select this strong cover so that the scales l_k converge to 0 very quickly.

Lemma 16. *Let $Z \subset \mathbf{R}^{dn}$ be a countable union of compact sets, each with lower Minkowski dimension at most α . Let $\{\varepsilon_k\}$ be a sequence of positive numbers and let $\{f_k\}$ be a sequence of functions such that $f_k: (0, \infty) \rightarrow (0, \infty)$. Then there exists a sequence of dyadic lengths $\{l_k\}$ and compact sets $\{Z_k\}$ such that*

- (A) *For each index $k \geq 2$, $l_k \leq f_{k-1}(l_{k-1})$.*
- (B) *For each index k , Z_k is a union of cubes in $\mathcal{B}_{l_k}^{dn}$.*
- (C) *Z is strongly covered by the sets $\{Z_k\}$.*
- (D) *For each index k , $\#\mathcal{B}_{l_k}^{dn}(Z_k) \leq (1/l_k)^{\alpha+\varepsilon_k}$.*

Proof. Let Z be the union of sets $\{Y_k\}$ with $\underline{\dim}_{\mathbf{M}}(Y_k) \leq \alpha$ for any k . Let m_1, m_2, \dots be a sequence of integers that repeats each integer infinitely often. For each positive integer k , since $\underline{\dim}_{\mathbf{M}}(Y_{m_k}) \leq \alpha$, definition (C) implies that there exists arbitrarily small lengths l which satisfy $\#\mathcal{B}_l^{dn}(Y_{m_k}) \leq 1/l^{\alpha+(\varepsilon_k/2)}$. Replacing l with a dyadic length at most twice the size of l , there are infinitely many *dyadic* scales l with

$$\#\mathcal{B}_l^{dn}(Y_{m_k}) \leq \frac{1}{(l/2)^{\alpha+\varepsilon_k}} \leq \frac{2^{dn}}{l^{\alpha+(\varepsilon_k/2)}} = \frac{(2^{dn}l^{\varepsilon_k/2})}{l^{\alpha+\varepsilon_k}}.$$

In particular, we may select a dyadic length l such that $2^{dn}l^{\varepsilon_k/2} \leq 1$, and, if $k \geq 2$, also satisfying $l \leq f_{k-1}(l_{k-1})$. The first constraint together with the last calculation

implies $\#\mathcal{B}_l^{dn}(Y_{m_k}) \leq 1/l^{\alpha+\varepsilon_k}$. We then set $l_k = l$, and define Z_k to be the union of all cubes in $\mathcal{B}_{l_k}^{dn}(Y_{m_k})$. \square

We now construct X by avoiding the various discretizations of Z at each scale. The aim is to find a nested decreasing family of discretized sets $\{X_k\}$ with $X = \bigcap X_k$. One condition guaranteeing that X avoids Z is that X_k^n is disjoint from *strongly non-diagonal* cubes in Z_k .

Lemma 17. *Let $Z \subset \mathbf{R}^{dn}$ and let $\{l_k\}$ be a sequence of lengths converging to zero. For each index k , let Z_k be a union of cubes in $\mathcal{B}_{l_k}^{dn}$, and suppose the sets $\{Z_k\}$ strongly cover Z . For each index k , let X_k be a union of cubes in $\mathcal{B}_{l_k}^d$. Suppose that for any k , X_k^n avoids strongly non-diagonal cubes in Z_k . If $X = \bigcap X_k$, then $(x_1, \dots, x_n) \notin Z$ for any distinct $x_1, \dots, x_n \in X$.*

Proof. Let $z \in Z$ be a point with distinct coordinates z_1, \dots, z_n . Set

$$\Delta = \{(w_1, \dots, w_n) \in \mathbf{R}^{dn} : \text{there exists } i \neq j \text{ such that } w_i = w_j\}.$$

Then $d(\Delta, z) > 0$, where d is the Hausdorff distance between Δ and z . Since $\{Z_k\}$ strongly covers Z , there is a subsequence $\{k_m\}$ such that $z \in Z_{k_m}$ for any index m . For suitably large m , the sidelength l_k cube I in Z_{k_m} containing z is disjoint from Δ . But this means I is strongly non-diagonal, and so $z \notin X_{k_m}^n$. In particular, z is not an element of X^n . \square

We are now ready to construct the set X in Theorem 18. Let $l_0 = 1$ and $X_0 = [0, 1)^d$. For each $k \geq 1$, define $\varepsilon_k = c_0/k$, where we view $c_0 = (dn - \alpha)/4$ as a irrelevant constant small enough that $dn - \alpha - 2\varepsilon_k > 0$ for any k . We set

$$f_k(x) = \min\left(x^{k^2}, (x^{dn}/2)^{1/(dn-\alpha-\varepsilon_{k+1})}, (1/2A_x)^{d(n-1)/\varepsilon_{k+1}}\right).$$

Apply Lemma 16 to Z with this choice of $\{\varepsilon_k\}$ and $\{f_k\}$; let $\{l_k\}$ be the resulting sequence of dyadic lengths and let $\{Z_k\}$ be the resulting strong cover of Z . Observe that the definition of f_k implies that for any index k ,

$$l_{k+1} \leq l_k^2, \quad l_{k+1}^{dn-\alpha-\varepsilon_{k+1}} \leq l_k^{dn}/2, \quad \text{and} \quad 2A_{l_k} l_{k+1}^{\varepsilon_{k+1}/d(n-1)} \leq 1. \quad (4.5)$$

For each index $k \geq 1$, define $l = l_k$ and $s = l_{k+1}$. Observe that X_k is a non-empty union of cubes in \mathcal{B}_l^d ; that Z_{k+1} is a union of cubes in \mathcal{B}_s^{dn} , and that (4.5) implies

$$|Z_{k+1}| \leq l_{k+1}^{dn-\alpha-\varepsilon_{k+1}} \leq l_k^{dn}/2 = l^{dn}/2.$$

Setting $Z_s = Z_{k+1}$, we are therefore justified in applying Lemma 15. This produces a dyadic length r , which we denote by r_{k+1} . Assuming $\alpha \geq d$, by (4.1) and (4.5), we find

$$\begin{aligned} r_{k+1} &\leq \max(l_{k+1}, 2A_{l_k}|Z_{k+1}|^{1/d(n-1)}) \leq \max(l_{k+1}, 2A_{l_k}l_{k+1}^{(dn-\alpha-\varepsilon_{k+1})/d(n-1)}) \\ &\leq \max(l_{k+1}, l_{k+1}^{(dn-\alpha-2\varepsilon_{k+1})/d(n-1)}) = l_{k+1}^{(dn-\alpha-2\varepsilon_{k+1})/d(n-1)}. \end{aligned} \quad (4.6)$$

For this choice of r , we obtain a set $F \subset X_k$ which is a union of cubes in $\mathcal{B}_s^d(E)$ satisfying Properties (A), (B), and (C) from the lemma, and we define $X_{k+1} = F$. Property (A) implies X_{k+1} avoids strongly non-diagonal cubes in Z_{k+1} , so if we define $X = \bigcap X_k$, then Lemma 17 implies $(x_1, \dots, x_n) \notin Z$ for any distinct $x_1, \dots, x_n \in X$.

Lemma 18.

$$\overline{\dim}_{\mathbf{M}}(X) \geq \frac{dn - \alpha}{n - 1}$$

Proof. Consider X as the limit of the sequence $\{X_k\}$. Since every cube in $\mathcal{B}_{l_{k+1}}^d(X_{k+1})$ intersects X , $\mathcal{B}_{l_{k+1}}^d(X_{k+1}) = \mathcal{B}_{l_{k+1}}^d(X)$. And so by Property (C) of Lemma 15, (4.5), and (4.6), we conclude

$$\begin{aligned} \frac{\log(\#\mathcal{B}_{l_{k+1}}^d(X))}{\log(1/l_{k+1})} &= \frac{\log(\#\mathcal{B}_{l_{k+1}}^d(X_{k+1}))}{\log(1/l_{k+1})} \geq \frac{\log((l_k/r_{k+1})^d \cdot \#\mathcal{B}_{l_k}^d(X_k))}{\log(1/l_{k+1})} \\ &= \frac{d \log(1/r_{k+1})}{\log(1/l_{k+1})} - \frac{d \log(1/l_k)}{\log(1/l_{k+1})} \\ &\geq \frac{d \log(1/l_{k+1}^{(dn-\alpha-2\varepsilon_{k+1})/d(n-1)})}{\log(1/l_{k+1})} - \frac{d \log(1/l_k)}{\log(1/l_k^{k^2})} \\ &= \frac{dn - \alpha - 2\varepsilon_{k+1}}{n - 1} - \frac{d}{k^2} \\ &= \frac{dn - \alpha}{n - 1} - o(1) \end{aligned}$$

Taking $k \rightarrow \infty$, we conclude $\overline{\dim}_{\mathbf{M}}(X) \geq (dn - \alpha)/(n - 1)$. \square

4.4 Dimension Bounds

To complete the proof of Theorem 18, we must show that $\dim_{\mathbf{H}}(X) \geq \beta$, where

$$\beta = \frac{dn - \alpha}{n - 1}.$$

We begin with a rough outline of our proof strategy. Recall that from the previous section, we have a decreasing sequence of lengths $\{l_k\}$. The most convenient way to examine the dimension of X at various scales is to use Frostman's lemma (see Definition (G)). We construct a probability measure μ supported on X such that for all $\varepsilon > 0$, for all dyadic lengths l , and for all $I \in \mathcal{B}_l^d$, $\mu(I) \lesssim_\varepsilon l^{\beta - \varepsilon}$. We begin by proving the bound $\mu(I) \lesssim l_k^{\beta - O(1/k)}$ when $I \in \mathcal{B}_{l_k}^d$, which we view as saying X looks like a set with dimension $\beta - O(1/k)$ at the lengths $\{l_k\}$. To obtain the complete dimension bound, it then suffices to interpolate to get an acceptable bound at all intermediate scales. In this construction, as in [6], the rapid decay of the lengths $\{l_k\}$ forced on us in the construction means interpolation poses a significant difficulty. We avoid this difficulty because of the uniform way that we have selected cubes in consecutive scales. This will imply that between the scales l_k and r_{k+1} , the mass of μ distributes with similar properties to the full dimensional Lebesgue measure, which makes interpolation easy.

We now define the measure μ , by first defining it as a premeasure on $\bigcup_{i=1}^{\infty} \mathcal{B}_{l_i}^d[0, 1)^d$. We initially set $\mu([0, 1)^d) = 1$. Given $I \in \mathcal{B}_{l_k}^d$, we find a parent cube $I' \in \mathcal{B}_{l_{k-1}}^d$ with $I \subset I'$. If $I \subset X_k$, then we set

$$\mu(I) = \frac{\mu(I')}{\#\mathcal{B}_{l_k}^d(X_k \cap I')}. \quad (4.7)$$

Otherwise, if $I \not\subset X_k$, we set $\mu(I) = 0$. Notice

$$\mu(I') = \sum_{I \in \mathcal{B}_{l_k}^d(X_k \cap I')} \frac{\mu(I)}{\#\mathcal{B}_{l_k}^d(X_k \cap I')} = \sum_{I \in \mathcal{B}_{l_k}^d(I')} \mu(I).$$

Thus mass is maintained at each stage of the construction, and then Proposition 1.7 of [5] implies μ is a premeasure, and thus extends to a measure on the entire Borel sigma algebra. The remainder of this section is devoted to showing that μ is a Frostman measure of dimension $\beta - \varepsilon$ for any $\varepsilon > 0$.

Lemma 19. *If $I \in \mathcal{B}_{l_k}^d$, then*

$$\mu(I) \leq 2^k \left[\frac{r_k \dots r_1}{l_{k-1} \dots l_1} \right]^d. \quad (4.8)$$

Proof. We prove the theorem inductively on k . For $k = 0$, the theorem is obvious, because $\mu(I) \leq 1$. For the purposes of induction, let $I \in \mathcal{B}_{l_k}^d$, together with a parent cube $I' \in \mathcal{B}_{l_{k-1}}^d$ with $I \subset I'$. If $\mu(I) > 0$, $I \subset X_k$, so $I' \subset X_{k-1}$. Because X_k was obtained from X_{k-1} via an application of Lemma 1, Property (C) of that lemma states that $\#\mathcal{B}_{l_k}^d(X_k \cap I) \geq (l_{k-1}/r_k)^d/2$, so together with the inductive hypothesis and (4.7), we conclude

$$\mu(I) = \frac{\mu(I')}{\#\mathcal{B}_{l_k}^d(X_k \cap I)} \leq \frac{\mu(I')}{(l_{k-1}/r_k)^d/2} \leq \frac{2^{k-1} \cdot \left[\frac{r_{k-1} \dots r_1}{l_{k-2} \dots l_1} \right]^d}{(l_{k-1}/r_k)^d/2} = 2^k \left[\frac{r_k \dots r_1}{l_{k-1} \dots l_1} \right]^d. \quad \square$$

Treating all parameters in (4.8) which depend on indices smaller than k as essentially constant, and using (4.6), we ‘conclude’ that

$$\mu(I) \lesssim r_k^d \lesssim l_k^{\beta-2\cdot\epsilon_k/(n-1)} = l_k^{\beta-O(1/k)}.$$

The bounds in (4.5) imply l_k decays very rapidly, which enables us to ignore quantities depending on previous indices, and obtain a true inequality.

Corollary. *There exists $c > 0$ such that for all $I \in \mathcal{B}_{l_k}^d$, $\mu(I) \lesssim l_k^{\beta-c/k}$.*

Proof. Given $\varepsilon > 0$, Lemma 19, Equation (4.6), the inequality $l_k \leq l_{k-1}^{(k-1)^2}$ from (4.5), and the fact that there exists a constant c_0 such that $\varepsilon_{k+1} = c_0/k$, we find

$$\begin{aligned} \mu(I) &\leq 2^k \left[\frac{r_k \dots r_1}{l_{k-1} \dots l_1} \right]^d \leq \left(\frac{2^k}{l_{k-1}^d \dots l_1^d} \right) l_k^{\beta-\varepsilon_k/(n-1)} = \left(\frac{2^k l_k^{2d/k}}{l_{k-1}^{d(k-1)}} \right) l_k^{\beta-\varepsilon_k/(n-1)-2d/k} \\ &\leq \left(2^k l_{k-1}^{(2d/k)(k-1)^2-d(k-1)} \right) l_k^{\beta-(c_0/(n-1)+2d)/k} = o\left(l_k^{\beta-(c_0/(n-1)+2d)/k} \right), \end{aligned}$$

so we can then set $c = c_0/(n-1) + 2d$. \square

Corollary 3 gives a clean expression of the β dimensional behaviour of μ at discrete scales. To obtain a Frostman measure bound at *all* scales, we need to apply a covering argument. This is where the uniform mass assignment technique

comes into play. Because μ behaves like a full dimensional set between the scales l_k and r_{k+1} , we won't be penalized for making the gap between l_k and r_{k+1} arbitrarily large. This is essential to our argument, because l_k decays faster than 2^{-k^m} for any $m > 0$.

Lemma 20. *If l is dyadic and $I \in \mathcal{B}_l^d$, then $\mu(I) \lesssim_k l^{\beta-c/k}$ for each integer k .*

Proof. We begin by assuming $l \leq l_k$. To bound $\mu(I)$, we apply a covering argument, which breaks into cases depending on the size of l in proportion to the scales l_k and r_k :

- If $r_{k+1} \leq l \leq l_k$, we can cover I by $(l/r_{k+1})^d$ cubes in $\mathcal{B}_{r_{k+1}}^d$. Because of Property (B) and (C) of Lemma 15, we know that the mass of each cube in $\mathcal{B}_{r_{k+1}}^d$ is bounded by at most $2(r_{k+1}/l_{k+1})^d$ times the mass of a cube in $\mathcal{B}_{l_k}^d$. Thus

$$\mu(I) \lesssim (l/r_{k+1})^d (2(r_{k+1}/l_{k+1})^d) l_k^{\beta-c/k} \leq 2l^d / l_k^{d-\beta+c/k} \leq 2l^{\beta-c/k},$$

where we used the fact that $d - \beta + c/k \geq 0$, so $l_k^{d-\beta+c/k} \geq l^{d-\beta+c/k}$.

- If $l_{k+1} \leq l \leq r_{k+1}$, we can cover I by a single cube in $\mathcal{B}_{r_{k+1}}^d$. Because of Property (B) of Lemma 15, each cube in $\mathcal{B}_{r_{k+1}}^d$ contains at most one cube of $\mathcal{B}_{l_{k+1}}^d(X_{k+1})$, so

$$\mu(I) \lesssim l_{k+1}^{\beta-c/k} \leq l^{\beta-c/k}.$$

- If $l \leq l_{k+1}$, there certainly exists m such that $l_{m+1} \leq l \leq l_m$, and one of the previous cases yields that $\mu(I) \lesssim l^{\beta-c/m} \leq l^{\beta-c/k}$.

If $l \geq l_k$, then $\mu(I) \leq 1 \lesssim_k l_k^{\beta-c/k} \leq l^{\beta-c/k}$, so $\mu(I) \lesssim_k l^{\beta-c/k}$ for arbitrary dyadic l . \square

Applying Frostman's lemma to Lemma 20 gives $\dim_{\mathbf{H}}(X) \geq \beta - c/k$ for each $k > 0$. Taking $k \rightarrow \infty$ proves the needed dimension bound. Since we have already shown in Section 4.3 that $(x_1, \dots, x_n) \notin Z$ for any distinct $x_1, \dots, x_n \in X$, this concludes the proof of Theorem 18.

4.5 Applications

As discussed in the introduction, Theorem 1 generalizes Theorems 1.1 and 1.2 from [6]. In this section, we present two applications of Theorem 18 in settings

where previous methods cannot obtain any results.

Theorem 19 (Sum-sets avoiding specified sets). *Let $Y \subset \mathbf{R}^d$ be a countable union of sets of Minkowski dimension at most α . Then there exists a set $X \subset \mathbf{R}^d$ with Hausdorff dimension at least $1 - \alpha$ such that $X + X$ is disjoint from Y .*

Proof. Define $Z = Z_1 \cup Z_2$, where

$$Z_1 = \{(x, y) : x + y \in Y\} \quad \text{and} \quad Z_2 = \{(x, y) : y \in Y/2\}.$$

Since Y is a countable union of sets of Minkowski dimension at most α , Z is a countable union of sets with lower Minkowski dimension at most $1 + \alpha$. Applying Theorem 18 with $d = 1$, $n = 2$, giving a set $X \subset \mathbf{R}^d$ with Hausdorff dimension $1 - \alpha$ avoiding Z . Since X avoids Z_1 , whenever $x, y \in X$ are distinct, $x + y \notin Y$. Since X avoids Z_2 , $X \cap (Y/2) = \emptyset$, and thus for any $x \in X$, $x + x \notin Y$. Thus $X + X$ is disjoint from Y . \square

Remark. *One weakness of our result is that as the number of variables n increases, the dimension of X tends to zero. If we try and make the n -fold sum $X + \dots + X$ disjoint from Y , current techniques only yield a set of dimension $(1 - \alpha)/(n - 1)$. We have ideas on how to improve our main result when Z is ‘flat’, in addition to being low dimension, which will enable us to remove the dependence of $\dim_{\mathbf{H}}(X)$ on n . In particular, we expect to be able to construct a set X of dimension $1 - \alpha$, such that X is disjoint from Y , and X is closed under addition, and multiplication by rational numbers. In particular, given a \mathbf{Q} subspace V of \mathbf{R}^d with dimension α , we can always find a ‘complementary’ \mathbf{Q} vector space W with complementary fractional dimension $d - \alpha$ such that $V \cap W = (0)$.*

One of the most interesting uses of our method is to construct subsets of fractals avoiding patterns. In [6], Fraser and the second author show that if γ is a C^2 curve with non-vanishing curvature, then there exists a set $E \subset \gamma$ of Hausdorff dimension $1/2$ that does not contain isocles triangles. Our method can extend this result from the case of curves to more general sets, which gives an example of the flexibility of our method. For simplicity, we stick to an analysis of planar sets.

Theorem 20 (Restricted sets avoiding isocles triangles). *Let $Y \subset \mathbf{R}^2$ and let $\pi : \mathbf{R}^2 \rightarrow \mathbf{R}$ be an orthogonal projection such that $\pi(Y)$ has non-empty interior. Let d be an arbitrary metric on \mathbf{R}^2 . Suppose that*

$$Z_0 = \{(y_1, y_2, y_3) \in Y^3 : d(y_1, y_2) = d(y_1, y_3)\}$$

is the countable union of sets with lower Minkowski dimension at most α , for $\varepsilon \geq 0$. Then there exists a set $X \subset Y$ with dimension at least $(3 - \alpha)/2$ so that no triple of points $(x_1, x_2, x_3) \in X^3$ form the vertices of an isocles triangle.

Proof. Without loss of generality, by translation and rescaling, we may assume $\pi(Y)$ contains $[0, 1)$. Form the set

$$Z = \pi(Z_0) = \{(\pi(y_1), \pi(y_2), \pi(y_3)) : y \in Z_0\}$$

Then Z is the projection of an α dimensional set, and therefore has dimension at most α . Applying Theorem 18 with $d = 1$ and $n = 3$, we construct a set $X_0 \subset [0, 1)$ with Hausdorff dimension at least $(3 - \alpha)/2$ such that for any distinct $x_1, x_2, x_3 \in X_0$, $(x_1, x_2, x_3) \notin Z$. Thus if we form a set X by picking, from each $x \in X_0$, a single element of $\pi^{-1}(x)$, then X avoids isocles triangles, and has Hausdorff dimension at least as large as X_0 . \square

To see that Theorem 3 indeed generalizes the result of Fraser and the second author, observe that if d is the Euclidean metric, then for every pair of points $x, y \in \mathbf{R}^2$, the set

$$\{z \in \mathbf{R}^2 : d(x, z) = d(y, z)\}$$

is the perpendicular bisector B_{xy} of x and y . If γ is a compact portion of a smooth curve with non-vanishing curvature, then the number of points in $\gamma \cap B_{xy}$ is bounded independantly of x and y . Thus the set Z_0 in the statement of Theorem 3 has Minkowski dimension at most 2, and we can find a set with Hausdorff dimension $(3 - 2)/2 = 1/2$ on the curve avoiding isocles triangles.

Results about slice of measures, such as those detailed in Chapter 6 of [12], show that for any one dimensional set Y , for almost every line L , $L \cap Y$ consists of a finite collection of points. This suggests that if Y is any set with fractional dimension one, then Z_0 has dimension at most 2. This implies that we can find a subset of Y with dimension $1/2$ avoiding curves. We are unsure if this is true for every set with dimension one, but we provide two examples suggesting this is true for a generic set. The first result shows that for any $\varepsilon > 0$, there is an infinite family of Cantor-type sets with dimension $1 + \varepsilon$ such that Z_0 has dimension at most $2 + \varepsilon$. The second shows that for any rectifiable curve, Z_0 has dimension at most 2. Thus Theorem 3 can be applied in settings where Y is incredibly ‘rough’, i.e. totally disconnected.

To study the first example, we consider a probabilistic model for a Cantor-type set which almost surely has the required properties. This model is obtained

by considering a nested decreasing family of discretized random sets $\{C_k\}$, with each C_k a union of sidelength $1/2^k$ squares. First, we fix $p \in [0, 1]$. Then we set $C_0 = [0, 1]^2$. To construct C_{k+1} , we split each sidelength $1/2^k$ cube in C_k into four sidelength $1/2^{k+1}$ squares, and keep each square in C_{k+1} with probability p . To study this model, we employ some results about tail bounds and asymptotics for branching processes.

Lemma 21. *If $p > 1/4$, then with non-zero probability, $\dim_{\mathbf{M}}(C) = 2 - \log_2(1/p)$.*

Proof. Let $p > 1/4$. For each k , let Z_k denote the number of sidelength $1/2^k$ cubes in $[0, 1]^2$. Then the sequence $\{Z_k\}$ is a branching process, where each cube can produce between zero and four subcubes, with each of these four cubes kept with probability p . Thus $\mathbf{E}[Z_{k+1}|Z_k] = (4p)Z_k$. Since $4p > 1$, A simple calculation, summarized in Theorem 8.1 of [7], shows that the process $W_k = Z_k/(4p)^k$ is an L^2 bounded martingale, and so there exists a random variable W such that $W_k \rightarrow W$ almost surely, and $\mathbf{E}(W|W_k) = W_k$ for all k . Whenever W is non-zero,

$$\dim_{\mathbf{M}}(C) = \lim_{k \rightarrow \infty} \frac{\log Z_k}{k \log 2} = \lim_{k \rightarrow \infty} \frac{\log(W_k/W) + \log(W(4p)^k)}{k \log 2} = 2 - \log_2(1/p).$$

Since $\mathbf{E}(W) = \mathbf{E}(\mathbf{E}(W|W_0)) = \mathbf{E}(W_0) = 1$, W is non-zero with positive probability. \square

Similar asymptotics for branching processes show that the set Z_0 associated with C as in Theorem 3 almost surely has Minkowski dimension $3 - \alpha$. We do this first by proving a supplementary result.

Lemma 22. *Let $\{Z_k\}$ be a supercritical branching process with extinction probability q . If we set $\tilde{M} = q + M$, then there exists small positive constants λ and ε , depending only on the offspring law of $\{Z_k\}$, such that $\mathbf{P}(Z_k \geq k\tilde{M}^k) \lesssim \exp(-\lambda k^{1+\varepsilon})$.*

Proof. Let q_0, q_1, \dots, q_M denote the offspring law for the branching process, so $q = q_0$. Then there exists a grid of i.i.d discrete random variables X_{ij} with $\mathbf{P}(X_{ij} = k) = q_k$ such that

$$Z_{k+1} = \sum_{j=1}^{Z_k} X_{ij}.$$

Now consider the branching process $\{\tilde{Z}_k\}$ defined by setting $\tilde{Z}_0 = 1$, and

$$\tilde{Z}_{k+1} = \sum_{j=1}^{\tilde{Z}_k} \max(X_{ij}, 1).$$

We find $Z_k \leq \tilde{Z}_k$, and if $\tilde{M} = q + M$, then

$$\mathbf{E}(\tilde{Z}_{k+1}|\tilde{Z}_k) = \left(q + \sum_{k=1}^N k q_k \right) \tilde{Z}_k = (q + M) \tilde{Z}_k = \tilde{M} \tilde{Z}_k.$$

Most importantly for our purposes, $\{\tilde{Z}_k\}$ has zero chance of extinction. Theorem 5 of [1] implies that for such a supercritical branching process, there exists $\lambda > 0$ depending only on the offspring distribution of \tilde{Z} , such that if $\tilde{W}_k = \tilde{Z}_k / \tilde{M}^k$, and $\tilde{W} = \lim \tilde{W}_k$, then

$$\mathbf{P}(|\tilde{W} - \tilde{W}_k| \geq t) \lesssim \exp(-\lambda t^{2/3} \tilde{M}^{k/3})$$

In particular, this means

$$\begin{aligned} \mathbf{P}(Z_k \geq (1 + \tilde{W}) \tilde{M}^k) &\leq \mathbf{P}(\tilde{Z}_k \geq (1 + \tilde{W}) \tilde{M}^k) \\ &\leq \mathbf{P}(|\tilde{W} \tilde{M}^k - \tilde{Z}_k| \geq \tilde{M}^k) \\ &= \mathbf{P}(|\tilde{W} - \tilde{W}_k| \geq 1) \lesssim \exp(-\lambda \tilde{M}^{k/3}), \end{aligned}$$

Theorem 2 of [3] implies that as $t \rightarrow \infty$, there are constants C and $\varepsilon > 0$ depending only on the offspring distribution of \tilde{Z} such that

$$-\log \mathbf{P}(\tilde{W} \geq t) \geq (C + o(1)) t^{1+\varepsilon}$$

In particular, there exists a small constant λ such that

$$\mathbf{P}(1 + \tilde{W} \geq k) \leq \exp(-\lambda k^{1+\varepsilon})$$

Applying a union bound gives

$$\begin{aligned} \mathbf{P}(Z_k \geq k \tilde{M}^k) &\leq \mathbf{P}(Z_k \geq (1 + \tilde{W}) \tilde{M}^k) + \mathbf{P}(1 + \tilde{W} \geq k) \\ &\lesssim \exp(-\lambda \tilde{M}^{k/3}) + \exp(-\lambda k^{1+\varepsilon}) \lesssim \exp(-\lambda k^{1+\varepsilon}). \quad \square \end{aligned}$$

For a set E , let $E_\delta = \{x : d(x, E) \leq \delta\}$ denote the δ thickened version of E .

Lemma 23. *There exists a constant B such that for any k , we can find lines $L_{k,1}, \dots, L_{k,M}$ with $M \leq B \cdot 8^{kN}$ such that for any line L , there exists i such that $[0, 1]^2 \cap L_{1/2^{kN+1}} \subset (L_{k,i})_{2^{-kN}}$.*

Proof. For each $(x, \theta) \in [0, 1]^2 \times [0, 1]$, let

$$L^{\theta, x} = \{x + t e^{2\pi i \theta} : t \in \mathbf{R}\}$$

Note that

$$L^{\theta,x} \cap [0,1]^2 \subset \{x + te^{2\pi i\theta} : t \in [-2,2]\}$$

Any line intersecting $[0,1]^2$ is equal to $L^{\theta,x}$ for some θ and some x . For any k , we consider the set of $8^5 \cdot 8^{kN}$ points $E = (\mathbf{Z}/2^{kN+5})^3 \cap [0,1]^3$. For any (x_1, θ_1) and (x_2, θ_2) , and $t \in [-2,2]$, we calculate that

$$|(x_1 + te^{2\pi i\theta_1}) - (x_2 + te^{2\pi i\theta_2})| \leq |x_1 - x_2| + 2\pi t|\theta_1 - \theta_2| \leq |x_1 - x_2| + 4\pi|\theta_1 - \theta_2|$$

Thus $L^{\theta_1,x_1} \cap [0,1]^2$ is contained in the $|x_1 - x_2| + 4\pi|\theta_1 - \theta_2|$ thickening of L^{θ_2,x_2} . For any line $L^{\theta,x}$, there exists $(\theta_0, x_0) \in E$ such that $|\theta - \theta_0| \leq 1/2^{kN+2}$ and $|x - x_0| \leq \sqrt{2}/2^{kN+5}$, and so $L^{\theta,x} \cap [0,1]^2$ is contained in the $\sqrt{2}/2^{kN+5} + 4\pi/2^{kN+5} \leq 14/2^{kN+5} \leq 1/2^{kN+1}$, and thus $L^{\theta,x}_{1/2^{kN+1}} \cap [0,1]^2 \subset L_{1/2^{kN}}$. Thus we may set $B = 8^5$, completing the proof. \square

Lemma 24. *Fix N . If $p > 1/2$, then almost surely, there exists a value k_0 such that if $k \geq k_0$, and L is any line, then*

$$\#\mathcal{B}_{2^{-kN}}^2(L_{2^{-kN}}) \leq k \cdot 10^k (2p)^{kN}$$

Proof. Fix N , and for any index k let $\delta_k = 1/2^{Nk}$. Given a line L , let I be a side-length δ_k square intersecting the δ_k thickened line L_{δ_k} . Then L passes from one edge of I to another edge, and so if we split I into 4^N sidelength δ_{k+1} squares, then $L_{\delta_{k+1}}$ intersects at most $5 \cdot 2^N$ of these boxes. Conditioned on I being contained in C_{kN} , each of these subboxes occurs in $C_{(k+1)N}$ with probability p^N . We let $Z_k(L)$ denote the number of sidelength $1/2^{Nk}$ boxes in C_{Nk} intersecting L_{δ_k} . We now find $\tilde{Z}_k(L) \geq Z_k(L)$ by adding *exactly* $5 \cdot 2^N$ potential subboxes of each box at each subsequent stage, then $\tilde{Z}_k(L)$ is a branching process, whose offspring distribution is independent of L . Since each subbox is added with probability p^N , the extinction probability of \tilde{Z}_k is $(1 - p^N)^{5 \cdot 2^N}$. Also, $\mathbf{E}(\tilde{Z}_{k+1}(L) | \tilde{Z}_k(L)) = 5 \cdot (2p)^N \tilde{Z}_k$. Setting $M = 5(2p)^N$ and $q = (1 - p^N)^{5 \cdot 2^N}$, Lemma 22 shows that there exists positive constants λ and ε , independent of L , such that if $\tilde{M} = M + q$,

$$\mathbf{P}(\tilde{Z}_k(L) \geq k\tilde{M}^k) \lesssim \exp(-\lambda k^{1+\varepsilon}).$$

Elementary bounds on the logarithm show that

$$5 \cdot 2^N \log(1 - p^N) \leq \frac{-5(2p)^N}{2 - p^N} \leq -5(2p)^N$$

so if $p > 1/2$,

$$q = (1 - p^N)^{5 \cdot 2^N} \leq e^{-5(2p)^N} \leq 1 \leq 5(2p)^N$$

and this implies $\tilde{M} \leq 10(2p)^N$, so we have shown

$$\mathbf{P}(\tilde{Z}_k(L) \geq k \cdot 10^k (2p)^{kN}) \lesssim \exp(-\lambda k^{1+\varepsilon}).$$

This gives an upper bound on $\tilde{Z}_k(L)$ up to a superexponentially decaying term in k .

For each k , Lemma 23 shows we can find lines $L_{k,1}, \dots, L_{k,M}$, with $M \leq B \cdot 8^{kN}$ such that for any line L , there exists i such that $[0, 1]^2 \cap L_{\delta_k/2} \subset (L_{k,i})_{\delta_k}$. Applying a union bound, we find that

$$\mathbf{P}(\text{there is } i \text{ such that } Z_k(L_{k,i}) \geq k \cdot 10^k (2p)^{kN}) \lesssim B \cdot 8^{kN} \exp(-ck^\lambda)$$

But since $\lambda > 1$, we therefore find

$$\sum_{k=1}^{\infty} \mathbf{P}(\text{there is } i \text{ such that } Z_k(L_{k,i}) \geq k \cdot 10^k (2p)^{kN}) < \infty$$

Applying the Borel-Cantelli lemma, we conclude that almost surely, it is eventually true for sufficiently large k that $Z_k(L_{k,i}) \leq k \cdot 10^k (2p)^{kN}$ for all i . In particular, the number of cubes intersecting $L_{\delta_k/2}$ for any line L is upper bounded by $k \cdot 10^k (2p)^{kN}$. \square

Lemma 25. *There exists a constant A such that if $I, J \in \mathcal{B}_{1/2^{kN}}^2[0, 1]^2$, with $d(I, J) \geq A/2^{kN}$, then*

$$\bigcup \{L_{xy} : x \in I, y \in J\} \subset L_{1/2^{kN+1}}$$

where we recall that L_{xy} is the bisector of the points x and y .

Proof. Denote the bottom left vertices of the boxes I and J by $(N_I, M_I) \cdot 1/2^{kN}$ and $(N_J, M_J) \cdot 1/2^{kN}$, with $0 \leq N_I, M_I, N_J, M_J \leq 2^{kN}$. Let $\Delta_N = |N_I - N_J|$ and $\Delta_M = |M_I - M_J|$. We may assume for simplicity that $N_I \leq N_J$ and $M_I \leq M_J$. Since $d(I, J) \geq A/2^{kN}$,

$$(\Delta_N^2 + \Delta_M^2)^{1/2} \geq A$$

If we let θ_1 and θ_2 denote the minimal and maximal angle between the lines

connecting pairs of points in I and J and the horizontal line, then

$$\begin{aligned}\sin(\theta_2 - \theta_1) &= \sin(\theta_2)\cos(\theta_1) - \sin(\theta_1)\cos(\theta_2) \\ &= \frac{(\Delta_M + 1)(\Delta_N + 1)}{\Delta_N^2 + \Delta_M^2} - \frac{\Delta_N\Delta_M}{\Delta_N^2 + \Delta_M^2} \\ &\leq \frac{\Delta_N + \Delta_M + 1}{\Delta_N^2 + \Delta_M^2} \leq 1/A\end{aligned}$$

Thus $\theta_2 - \theta_1 \leq 2/A$. Thus all the bisectors are contained in a

$$\sqrt{2}/2^{kN} + 4\pi(2/A) \leq 1/2^{kN}$$

□

Theorem 21. *If $p > 1/2$, then almost surely, the set Z_0 associated with C has lower Minkowski dimension at most $5 - 3\log_2(1/p)$.*

Proof. Almost surely, Lemma 21 shows that for sufficiently large k we can cover C_{kN} by $O(2^{\alpha kN})$ boxes $I_1, \dots, I_M \in \mathcal{B}_{2^{-kN}}^2$, with $\alpha = 2 - \log_2(1/p)$. Lemma 24 shows that if k is sufficiently large, then the $1/2^{kN+1}$ thickened line L intersects at most $k \cdot 10^k (2p)^{kN}$ squares in $\mathcal{B}_{2^{-kN}}^2$. There exists a constant A such that for any i and j , if $d(I_i, I_j) > A2^{-kN}$, then there exists a line L_{ij} such that as x ranges over all points in I_i , and y over all points in I_j ,

$$\bigcup_{x,y} B_{xy} \cap [0, 1]^2 \subset (L_{ij})_{\delta_k/2}$$

This means that $Z_0 \cap (I_i \times I_j \times [0, 1]^2) \subset I_i \times I_j \times (L_{ij})_{\delta_k/2}$, and the last paragraph implies that $(L_{ij})_{\delta_k/2}$ is covered by $k \cdot 10^k (2p)^{kN}$ cubes in $\mathcal{B}_{l_k}^2[0, 1]^2$. On the other hand, if $d(I_i, I_j) < A\delta_k$, we can apply the obvious bound that $Z_0 \cap (I_i \times I_j \times [0, 1]^2) \subset I_i \times I_j \times C_{kN}$, and C_{kN} is coverable by $O(2^{\alpha kN})$ boxes. Thus we conclude that almost

surely, for sufficiently large k ,

$$\begin{aligned}
\# \mathcal{B}_{\delta_k}^6(Z_0) &= \sum_{i,j} \mathcal{B}_{\delta_k}^6(Z_0 \cap (I_i \times I_j \times [0, 1]^2)) \\
&\leq \sum_i \left(\sum_{d(I_i, I_j) \leq A\delta_k} \# \mathcal{B}_{\delta_k}^6(Z_0 \cap (I_i \times I_j \times (L_{ij})_{\delta_k/2})) \right) \\
&\quad + \left(\sum_{d(I_i, I_j) > A\delta_k} \# \mathcal{B}_{\delta_k}^6(Z_0 \cap (I_i \times I_j \times C_{kN})) \right) \\
&\leq \sum_i \left(\sum_{d(I_i, I_j) \leq A\delta_k} \# \mathcal{B}_{\delta_k}^2((L_{ij})_{\delta_k/2}) \right) + \left(\sum_{d(I_i, I_j) > A\delta_k} \# \mathcal{B}_{\delta_k}^2(C_{kN}) \right) \\
&\lesssim \sum_i 2^{\alpha kN} \cdot k 10^k (2p)^{kN} + 2^{\alpha kN} \lesssim 2^{2\alpha kN} k 10^k (2p)^{kN}
\end{aligned}$$

Thus we conclude that almost surely,

$$\begin{aligned}
\dim_{\mathbf{M}}(Z_0) &= \lim_{k \rightarrow \infty} \frac{\log(\# \mathcal{B}_{\delta_k}^6(Z_0))}{\log(1/\delta_k)} \\
&\leq \lim_{k \rightarrow \infty} \frac{\log(2^{2\alpha kN} k 10^k (2p)^{kN}) + O(1)}{\log(2^{kN})} \\
&= \lim_{k \rightarrow \infty} \frac{2\alpha kN \log(2) + k \log 10 + kN \log(2p)}{kN \log(2)} \\
&= 2\alpha + 3/N + 1 - \log_2(1/p) \\
&= 5 - 3 \log_2(1/p) + 3/N
\end{aligned}$$

Taking $N \rightarrow \infty$, we obtain the required result. \square

TODO: does a projection of C have non-empty interior almost surely?

Corollary. *If $p = 1/2^{1-\varepsilon}$, then conditioned on C having Minkowski dimension $1 + \varepsilon$, we can find a subset X of C avoiding isocoles triangles with $\dim_{\mathbf{H}}(X) \geq 1/2 - (3/2)\varepsilon$.*

4.6 Relation to Literature, and Future Work

Our result is part of a growing body of work finding general methods to find sets avoiding patterns. The main focus of this section is comparing our method to the

two other major results in the literature. [6] constructs sets with dimension $k/(n-1)$ avoiding the zero sets of rank k C^1 functions. In [11], sets of dimension d/l are constructed avoiding a degree l algebraic hypersurface specified by a polynomial with rational coefficients.

We can view our result as a robust version of Pramanik and Fraser’s result. Indeed, if we try and avoid the zero set of a C^1 rank k function, then we are really avoiding a dimension $dn-k$ dimensional manifold. Our method gives a dimension

$$\frac{dn - (dn - k)}{n - 1} = \frac{k}{n - 1}$$

set, which is exactly the result obtained in [6].

That our result generalizes [6] should be expected because the technical skeleton of our construction is heavily modeled after their construction technique. Their result also reduces the problem to a discrete avoidance problem. But they *deterministically* select a particular side length S cube in every side length R cube. For arbitrary Z , this selection procedure can easily be exploited for a particularly nasty Z , so their method must rely on smoothness in order to ensure some cubes are selected at each stage. Our discrete avoidance technique was motivated by other combinatorial optimization problems, where adding a random quantity prevents inefficient selections from being made in expectation. This allows us to rely purely on combinatorial calculations, rather than employing smoothness, and greatly increases the applicability of the sets Z we can apply our method to. Furthermore, it shows that the underlying problem is robust to changes in dimension; slightly ‘thickening’ Z only slightly perturbs the dimension of X .

One useful technique in [6], and its predecessor [9], is the use of a Cantor set construction ‘with memory’; a queue in the construction algorithm for their sets allows storage of particular discrete versions of the problem to be stored, and then retrieved at a much later stage of the construction process. This enables them to ‘separate’ variables in the discrete version of the problem, i.e. instead of forming a single set F from a set E , they form n sets F_1, \dots, F_n from disjoint sets E_1, \dots, E_n . The fact that our result is more general, yet does not rely on this technique is an interesting anomaly. An obvious advantage is that the description of the technique is much more simple. But an additional advantage is that we can attack ‘one scale’ of the problem at a time, rather than having to rely on stored memory from a vast number of steps before the current one. We believe that we can exploit the single scale approach to the problem to generalize our theorem to a much wider family of ‘dimension α ’ sets Z , which we plan to discuss in a later paper.

As a generalization of the result in [6], our result has the same issues when compared to the result of [11]. When the parameter n is large, the dimension of our result suffers greatly, as with the n fold sum application in the last section. Furthermore, our result can't even beat trivial results if Z is almost full dimensional, as the next example shows.

Example. Consider an α dimensional set of angles Y , and try and find $X \subset \mathbf{R}^2$ such that the angle formed from any collection of three points in X avoids Y . If we form the set

$$Z = \left\{ (x, y, z) : \text{There is } \theta \in Y \text{ such that } \frac{(x-y) \cdot (x-z)}{|x-y||x-z|} = \cos \theta \right\}$$

Then we can find X avoiding Z . But one calculates that Z has dimension $3d + \alpha - 1$, which means X has dimension $(1 - \alpha)/2$. Provided the set of angles does not contain π , the trivial example of a straight line beats our result.

Nonetheless, we still believe our method is a useful inspiration for new techniques in the ‘high dimensional’ setting. Most prior literature studies sets Z only implicitly as zero sets of some regular function f . The features of the function f imply geometric features of Z , which are exploited by these results. But some geometric features are not obvious from the functional perspective; in particular, the fractional dimension of the zero set of f is not an obvious property to study. We believe obtaining methods by looking at the explicit geometric structure of Z should lead to new techniques in the field, and we already have several ideas in mind when Z has geometric structure in addition to a dimension bound, which we plan to publish in a later paper.

We can compare our randomized selection technique to a discrete phenomenon that has been recently noticed, for instance in [2]. There, certain combinatorial problems can be rephrased as abstract problems on hypergraphs, and one can then generalize the solutions of these problems using some strategy to improve the result where the hypergraph is sparse. Our result is a continuous analogue of this phenomenon, where sparsity is represented by the dimension of the set Z we are trying to avoid. One can even view Lemma 1 as a solution to a problem about independent sets in hypergraphs. In particular, we can form a hypergraph by taking the cubes $\mathcal{B}_s^d(E)$ as vertices, and adding an edge (I_1, \dots, I_n) between n distinct cubes $I_k \in \mathcal{B}_s^d(E)$ if $I_1 \times \dots \times I_n$ intersects Z_s . An independent set of cubes in this hypergraph corresponds precisely to a set F with F^n disjoint except on a discretization of the diagonal. And so Lemma 1 really just finds a ‘uniformly cho-

sen' independent set in a sparse graph. Thus we really just applied the discrete phenomenon at many scales to obtain a continuous version of the phenomenon.

Chapter 5

Extensions to Low Rank Configurations

5.1 Boosting the Dimension of Pattern Avoiding Sets by Low Rank Coordinate Changes

We now consider finding subsets of $[0, 1]$ avoiding solutions to the equation $y = f(Tx)$, where T is a rank k linear transformation with integer coefficients with respect to standard coordinates, and f is real-valued and Lipschitz continuous. Fix a constant A bounding the operator norm of T , in the sense that $|Tx| \leq A|x|$ for all $x \in \mathbf{R}^n$, and a constant B such that $|f(x+y) - f(x)| \leq B|y|$ for all x and y for which the equation makes sense (if f is C^1 , this is equivalent to a bound $\|\nabla f\|_\infty \leq B$). Consider sets $J_0, J_1, \dots, J_n, \subset [0, 1]$, which are unions of intervals of length $1/M$, with startpoints lying on integer multiples of $1/M$. The next theorem works as a ‘building block lemma’ used in our algorithm for constructing a set avoiding solutions to the equation with Hausdorff dimension k and full Minkowski dimension.

Theorem 22. *For infinitely many integers N , there exists $S_i \subset J_i$ avoiding solutions to $y = f(Tx)$ with $y \in S_0$ and $x_n \in S_n$, such that*

- *For $n \neq 0$, if we decompose each J_i into length $1/N$ consecutive intervals, S_i contains an initial portion $\Omega(1/N^k)$ of each length $1/N$ interval. This part of the decomposition gives the Hausdorff dimension $1/k$ bound for the set we will construct.*

- If we decompose J_i into length $1/N$ intervals, and then subdivide these intervals into length $\Omega(1/N^k)$ intervals, then S_0 contains a subcollection of these $1/N^k$ intervals which contains a total length $\Omega(1/N)$ of a fraction $1 - 1/M$ of the length $1/N$ intervals. This property gives that our resultant set will have full Minkowski dimension.

The implicit constants in these bounds depend only on A , B , n , and k .

Proof. Split each interval of J_a into length $1/N$ intervals, and then set

$$\mathbf{A} = \{x : x_a \text{ is a startpoint of a } 1/N \text{ interval in } J_a\}$$

Since the startpoints of the intervals are integer multiples of $1/N$, $T(\mathbf{A})$ is contained within a rank k sublattice of $(\mathbf{Z}/N)^m$. The operator norm also guarantees $T(\mathbf{A})$ is contained within the ball B_A of radius A in \mathbf{R}^m . Because of the lattice structure of the image, $|x - y| \gtrsim_n 1/N$ for each distinct pair $x, y \in T(\mathbf{A})$. For any R , we can cover $\Sigma \cap B_A$ by $O_{n,k}((A/R)^k)$ balls of radius R . If $R \gtrsim_n 1/N$, then each ball can contain only a single element of $T(\mathbf{A})$, so we conclude that $|T(\mathbf{A})| \lesssim_{n,k} (AN)^k$. If we define the set of ‘bad points’ to be

$$\mathbf{B} = \{y \in [0, 1] : \text{there is } x \in \mathbf{A} \text{ such that } y = f(T(x))\}$$

Then

$$|\mathbf{B}| = |f(T(\mathbf{A}))| \leq |T(\mathbf{A})| = O_{A,n,k}(N^k)$$

For simplicity, we now introduce an integer constant $C_0 = C_0(A, n, k, M)$ such that $|\mathbf{B}| \leq (C_0/M^2)N^k$. We now split each length $1/M$ interval in J_0 into length $1/N$ intervals, and filter out those intervals containing more than C_0N^{k-1} elements of \mathbf{B} . Because of the cardinality bound we have on \mathbf{B} there can be at most N/M^2 such intervals, so we discard at most a fraction $1/M$ of any particular length $1/M$ interval in J_0 . If we now dissect the remaining intervals into $4C_0N^{k-1}$ intervals of length $1/4C_0N^k$, and discard any intervals containing an element of \mathbf{B} , or adjacent to such an interval, then the remaining such intervals I satisfy $d(I, \mathbf{B}) \geq 1/4C_0N^k \gtrsim_{A,n,M,k} (1/N^k)$, and because of our bound on the number of elements of \mathbf{B} in these intervals, there are at least C_0N^{k-1} intervals remaining, with total length exceeding $C_0N^{k-1}/4C_0N^k = \Omega(1/N)$. If f is C^1 with $\|\nabla f\|_\infty \leq B$, or more generally, if f is Lipschitz continuous of magnitude B , then

$$|f(Tx) - f(Tx')| \leq AB|x - x'|$$

and so we may choose $S_i \subset J_i$ by thickening each startpoint $x \in J_i$ to a length $O(1/N^k)$ interval while still avoiding solutions to the equation $y = f(T(x))$. \square

Remark. *If T is a rank k linear transformation with rational coefficients, then there is some number a such that aT has integer coefficients, and then the equation $y = f(Tx)$ is the same as the equation $y = f_0((aT)(x))$, where $f_0(x) = f(x)/a$. Since f_0 is also Lipschitz continuous, we conclude that we still get the dimension $1/k$ bound if T has rational rather than integral coefficients. More generally, this trick shows the result applies unperturbed if all coefficients of T are integer multiples of some fixed real number. More generally, by varying the lengths of our length $1/N$ decomposition by a constant amount, we can further generalize this to the case where each column of T are integers multiples of some fixed real number.*

Remark. *To form \mathbf{A} , we take startpoints lying at equal spaced $1/N$ points. However, by instead taking startpoints at varying points in the length $1/N$ intervals, we might be able to make points cluster more than in the original algorithm. Maybe the probabilistic method would be able to guarantee the existence of a choice of startpoints whose images are tightly clustered together.*

Remark. *Since the condition $y = f(Tx)$ automatically assumes a kind of ‘non-vanishing derivative’ condition on our solutions, we do not need to assume the regularity of f , and so the theorem extends naturally to a more general class of functions than Rob’s result, i.e. the Lipschitz continuous functions.*

Using essentially the same approach as the last argument shows that we can avoid solutions to $y = f(Tx)$, where y and x are now vectors in some \mathbf{R}^m , and T has rank k . If we consider unions of $1/M$ cubes J_0, \dots, J_n . If we fix startpoints of each x_k forming lattice spaced apart by $\Omega(1/N)$, and consider the space \mathbf{A} of products, then there are $O(N^k)$ points in $T(\mathbf{A})$, and so there are $O(N^k)$ elements in \mathbf{B} . We now split each $1/M$ cube in J_0 into length $1/N$ cubes, and discard those cubes which contain more than $O(N^{k-m})$ bad points, then we discard at most $1 - 1/M$ of all such cubes. We can dissect the remaining length $1/N$ cubes into $O(N^{k-m})$ length $\Omega(1/N^{k/m})$ cubes, and as in the previous argument, the cubes not containing elements of \mathbf{B} nor adjacent to an element have total volume $\Omega(1/N^m)$, which we keep. The startpoints in the other intervals T_i may then be thickened to a length $\Omega(1/N^{m/k})$ portion while still avoiding solutions. This gives a set with full Minkowski dimension and Hausdorff dimension m/k avoiding solutions to $y = f(Tx)$. (I don’t yet understand Minkowski dimension enough to understand this, but the techniques of the appendix make proving the Hausdorff dimension $1/k$ bound easy)

5.2 Extension to Well Approximable Numbers

If the coefficients of the linear transformation T in the equation $y = f(Tx)$ are non-rational, then the images of startpoints under the action of T do not form a lattice, and so points may not overlap so easily when avoiding solutions to the equations $y = f(Tx)$. However, if T is ‘very close’ to a family of rational coefficient linear transformations, then we can show the images of the startpoints are ‘very close’ to a lattice, which will still enable us to find points avoiding solutions by replacing the direct combinatorial approach in the argument for integer matrices with a covering argument.

Suppose that T is a real-coefficient linear transformation with the property that for each coefficient x there are infinitely many rational numbers p/q with $|x - p/q| \leq 1/q^\alpha$, for some fixed α . For infinitely many K , we can therefore find a linear transformation S with coefficients in \mathbf{Z}/K with each coefficient of T differing from the corresponding coefficient in S by at most $1/K^\alpha$. Then for each x , we find

$$\|(T - S)(x)\|_\infty \leq (n/K^\alpha)\|x\|_\infty$$

If we now consider T_0, \dots, T_n , splitting T_1, \dots, T_n into length $1/N$ intervals, and considering \mathbf{A} as in the last section, then $S(\mathbf{A})$ lie in a k dimensional sublattice of $(\mathbf{Z}/KN)^m$, hence containing at most $(2A)^k(KN)^k = O_{T,n}((KN)^k)$ points. By our error term calculation of $T - S$, the elements of $T(\mathbf{A})$ are contained in cubes centered at these lattice points with side-lengths $2n/K^\alpha$, or balls centered at these points with radius $n^{3/2}/K^\alpha$. If $\|\nabla f\| \leq B$, then the images of the radius $n^{3/2}/K^\alpha$ balls under the action of f are contained in length $Bn^{3/2}/K^\alpha$ intervals. Thus the total length of the image of all these balls under f is $(Bn^{3/2}/K^\alpha)(2A)^k(KN)^k = (2A)^k Bn^{3/2} K^{k-\alpha} N^k$. If $k < \alpha$, then we can take K arbitrarily large, so that there exists intervals with $\text{dist}(I, \mathbf{B}) = \Omega_{A,k,M}(1/N^k)$. But I believe that, after adding the explicit constants in, we cannot let $k = \alpha$.

Remark. *One problem is that, if T has rank k , we might not be able to choose S to be rank k as well. Is this a problem? If T has full rank, then the set of all such matrices is open so if T and S are close enough, S also has rank k , but this need not be true if T does not have full rank.*

Example. *If T has rank 1, then Dirichlet’s theorem says that every irrational number x can be approximated by infinitely many p/q with $|x - p/q| < 1/q^2$, so every real-valued rank 1 linear transformation can be avoided with a dimension one bound.*

5.3 Equidistribution and Real Valued Matrices

If T is a non-invertible matrix containing irrational coefficients, then the values Tx , for $x \in \mathbf{Z}^n$, do not form a lattice, and therefore we cannot use the direct combinatorial arguments of the past section to obtain the decomposition lemma. However, without loss of generality, we can write $T(x) = S(x_1) + U(x_2)$, where $x = (x_1, x_2)$, $x_1 \in \mathbf{R}^k$, $x_2 \in \mathbf{R}^{n-k}$, and S has full rank k . Then S is an embedding of \mathbf{R}^k into \mathbf{R}^n , so $\Gamma = S((\mathbf{Z}/N)^k)$ forms a lattice with points spaced apart by a distance on the order of $\Omega(1/N)$. Since T has rank k , the image of U is contained within the image of S . We let \mathbf{T} denote the torus obtained by quotienting the k dimensional subplane forming the image of T by Γ . Then the image of S in \mathbf{T} is contained within an α dimensional subtorus of \mathbf{T} . Note that $\alpha = 0$ precisely when T still has rank k over the rational numbers, so that in a suitable basis T is an integer valued matrix. If $\alpha = 1$, then by an appropriate scaling in the values x_2 we can still make the values of S lie at lattice points, which should give a Hausdorff dimension one set. When $\alpha = 2$, we run into problems.

If $\pi : \mathbf{R}^k \rightarrow T$ is the homomorphism obtained by composing the quotient map onto the torus with the linear map T , then $\pi(x_1, 0) = 0$ for all $x_1 \in (\mathbf{Z}/N)^k$. On the other hand, Ratner's theorem implies that for each $x_2 \in \mathbf{R}^{n-k}$, there is some M such that the sequence $\pi(0, nMx_2) = Mn\pi(0, x_2)$ is equidistributed on a subtorus of T . Equidistribution may be useful in extending the rational matrix result to all real matrices with some dimension loss, since a matrix is rational if and only if $nM\pi(0, x_2)$ is equidistributed on a zero dimensional lattice – it may ensure that points are closely clustered to lattice points.

Lets consider the simplest case, where $n = k + 1$, so $x_2 \in \mathbf{R}$. Consider our setup, with intervals J_0, \dots, J_n , and an equation $y = f(Tx)$, where f is Lipschitz with Lipschitz norm bounded by B . Now if $T = S + U$, then $S(\mathbf{Z}^k)$ forms a rank k lattice, and $U(\mathbf{Z})$ equidistributes over an α dimensional subtorus of the torus generated over the lattice. In particular, since the set $\mathbf{Z} \cap NJ_n$ contains $\Omega(N|J_n|) = \Omega(N)$ consecutive points, for any ε and suitably large N , $\mathbf{Z} \cap NJ_n$ contains $\Omega(Nr^\alpha)$ points x such that $S(x)$ is within a distance r from a lattice point, for any r . Dividing by N tells us that we have $O(Nr^\alpha)$ points x in $\mathbf{Z}/N \cap J_n$ such that $U(x)$ is at a distance r/N from a lattice point in $S((\mathbf{Z}/N)^k)$. If $r = 1/N^\beta$, then we have $O(N^{1-\alpha\beta})$ points at a distance $1/N^{\beta+1}$ from a lattice point. There are $O(N^k)$ points in the lattice, and so provided that $k < 1 + \beta$, we can find a large subset avoiding the images of these startpoints, and we should be able to thicken the startpoints to length h $\Omega(1/N^k)$ intervals, hence we should expect the set we construct to have Hausdorff dimension $1/k(k-1)$ if this process is repeated to construct our solution avoiding

set.

5.4 Applications of Low Rank Coordinate Changes

Example. Our initial exploration of low rank coordinate changes was inspired by trying to find solutions to the equation

$$y - x = (u - w)^2$$

Our algorithm gives a Hausdorff dimension $1/2$ set avoiding solutions to this equation. This equals Mathé's result. But this dimension for us now depends on the shifts involved in the equation, not on the exponent, so we can actually avoid solutions to the equation

$$y - x = (u - w)^n$$

for any n , in a set of Hausdorff dimension $1/2$. More generally, if X is a set, then given a smooth function f of n variables, we can find a set X of Hausdorff dimension $1/n$ such that there is no $x \in X$, and $y_1, \dots, y_n \in X - X$ such that $x = f(y_1, \dots, y_n)$. This is better than the $1/2n$ bound that is obtained by Malabika and Fraser's result.

Example. For any fixed m , we can find a set $X \subset \mathbf{R}^n$ of full Hausdorff dimension which contains no solutions to

$$a_1x_1 + \dots + a_nx_n = 0$$

for any rational numbers a_n which are not all zero. Since Malabika/Fraser's technique's solutions are bounded by the number of variables, they cannot let $n \rightarrow \infty$ to obtain a linearly independent set over the rational numbers. But since the Hausdorff dimension of our sets now only depends on the rank of T , rather than the total number of variables in T , we can let $n \rightarrow \infty$ to obtain full sets linearly independent over the rationals. More generally, for any Lipschitz continuous function $f : \mathbf{R} \rightarrow \mathbf{R}$, we can find a full Hausdorff dimensional set such that there are no solutions

$$f(a_1x_1 + \dots + a_nx_n, y)$$

for any n , and for any rational numbers a_n that are not all zero.

Example. The easiest applications of the low rank coordinate change method are probably involving configuration problems involving pairwise distances be-

tween m points in \mathbf{R}^n , where $m \ll n$, since this can best take advantage of our rank condition. Perhaps one way to encompass this is to avoid m vertex polyhedra in n dimensional space, where $m \ll n$. In order to distinguish this problem from something that can be solved from Mathé's approach, we can probably find a high dimensional set avoiding m vertex polyhedra on a parameterized n dimensional manifold, where $m \ll n$. There is a result in projective geometry which says that every projectively invariant property of m points in \mathbf{RP}^d is expressible as a function in the $\binom{m}{d}$ bracket polynomials with respect to these m points. In particular, our result says that we can avoid a countable collection of such invariants in a dimension $1/\binom{m}{d}$ set. This is a better choice of coordinates than Euclidean coordinates if $\binom{m}{d} \leq m$. Update: I don't think this is ever the case.

Remark. Because of how we construct our set X , we can find a dimension $1/k$ set avoiding solutions to $y = f(Tx)$ for all rank k rational matrices T , without losing any Hausdorff dimension. Maybe this will help us avoid solutions to more general problems?

Example. Given a smooth curve Γ in \mathbf{R}^n , can we find a subset E with high Hausdorff dimension avoiding isocles triangles. That is, if the curve is parameterized by $\gamma: [0, 1] \rightarrow \mathbf{R}^n$, can we find $E \subset [0, 1]$ such that for any t_1, t_2, t_3 , $\gamma(t_1)$, $\gamma(t_2)$, and $\gamma(t_3)$ do not form the vertices of an isocles triangle. This is, in a sense, a non-linear generalization of sets avoiding arithmetic progressions, since if Γ is a line, an isocles triangle is given by arithmetic progressions. Assuming our curve is simple, we must avoid zeroes of the function

$$|\gamma(t_1) - \gamma(t_2)|^2 = |\gamma(t_2) - \gamma(t_3)|^2$$

If we take a sufficiently small segment of this curve, and we assume the curve has non-zero curvature on this curve, we can assume that $t_1 < t_2 < t_3$ in our dissection method.

If the coordinates of γ are given by polynomials with maximum degree d , then the equation

$$|\gamma(t_1) - \gamma(t_2)|^2 - |\gamma(t_2) - \gamma(t_3)|^2$$

is a polynomial of degree $2d$, and so Mathé's result gives a set of dimension $1/2d$ avoiding isocles triangles. In the case where Γ is a line, then the function $f(t_1, t_2, t_3) = \gamma(t_1) + \gamma(t_3) - 2\gamma(t_2)$ avoids arithmetic progressions, and Mathé's result gives a dimension one set avoiding such progressions. Rob and Malabika's algorithm easily gives a set with dimension $1/2$ for any curve Γ . Our algorithm doesn't seem to be able to do much better here.

Example. What is the largest dimension of a set in Euclidean space such that for any value λ , there is at most one pair of points x, y in the set such that $|x - y| = \lambda$.

Example. What is the largest dimension of a set which avoids certain angles, i.e. for which a triplet x, y, z avoids certain planar configurations.

Example. A set of points $x_0, \dots, x_d \in \mathbf{R}^d$ lie in a hyperplane if and only if the determinant formed by the vectors $x_n - x_0$, for $n \in \{1, \dots, d\}$, is zero. This is a degree d polynomial, hence Mathé's result gives a dimension one set with no set of $d + 1$ points lying in a hyperplane. On the other hand, a theorem of Mattila shows that every analytic set E with dimension exceeding one contains $d + 1$ points in a hyperplane. Can we generalize this to a more general example avoiding points on a rotational, translation invariant family of manifolds using our results?

Example. Given a set F not containing the origin, what is the largest Hausdorff dimension of a set E such that for any for any distinct rational a_1, \dots, a_N , the sum $a_1 E + \dots + a_N E$ does not contain any elements of F . Thus the vector space over the rationals generated by E does not contain any elements of F . We can also take the non-linear values $f(a_1 E + \dots + a_N E)$ avoiding elements of F . F must have non-empty interior for the problem to be interesting. Then can we find a smooth function f with non-vanishing derivative which vanishes over F , or a family of smooth functions with non-vanishing derivative around F .

5.5 Idea: Generalizing This Problem to low rank smooth functions

Suppose we are able to find dimension $1/k$ sets avoiding configurations $y = g(f(x))$, where f is a smooth function from $\mathbf{R}^n \rightarrow \mathbf{R}^m$ of rank k . Then given any function $g(f(x))$, where f has rank k , if $g(f(x)) = 0$, then the implicit function theorem guarantees that there is a cover U_α and functions $h_\alpha : U^k \rightarrow \mathbf{R}^{n-k}$ such that for each if $g(f(x)) = 0$, for $x \in U_\alpha$, then there is a subset of k indices I such that $x_{I^c} = h_\alpha(x_I)$.

Then given any function $f(x)$ with rank k , we can use the implicit function theorem to find sets U_α , indices n_α , and functions g_α such that if $f(x) = 0$, for

$x \in U_\alpha$, then $x_{n_\alpha} = g_\alpha(x_1, \dots, x_{n_\alpha}, \dots, x_n)$. Thus we need only avoid this type of configuration to avoid configurations of a general low rank function. If we don't believe that we are able to get dimension $1/(k - 1)$ sets for rank k configurations,

then we shouldn't be able to find sets of Hausdorff dimension $1/k$ avoiding configurations of the form $y = f(x)$, where f has rank k .

Remark. *If the functions g_α are only partially defined, this makes the problem easier than if the functions were globally defined, because the constraint condition is now smaller than the original constraint.*

Remark. *This would solve our problem of avoiding $y - x = (u - v)^2$, since if $f(x, y, u, v) = y - x - (u - v)^2$, then*

$$\nabla f$$

5.6 Idea: Algebraic Number Fields

If $\mathbf{Q}(\omega)$ is a quadratic extension of the rational numbers, then the ring of integers in this field form a lattice. Perhaps we can use this to generalize our approach to avoiding configurations $y = f(Tx)$, where all coefficients of the matrix T lie in some common quadratic extension of the rational numbers.

5.7 A Scheme for Avoiding Configurations

Mathé's result can be reconfigured in terms of a building block strategy for implementation in our algorithm.

Theorem 23. *Let f be a polynomial of degree m , and consider unions of length $1/M$ intervals $T_0, \dots, T_d \subset [0, 1]$, with rational start-points. If $\partial_0 f$ is non-vanishing on $T_0 \times \dots \times T_d$, then there exists arbitrarily large integers N and a constant C not depending on N and sets $S_n \subset T_n$ such that*

- $f(x) \neq 0$ for $x \in S_0 \times \dots \times S_d$.
- If T_0, \dots, T_d are split into length $1/N$ intervals, then S_n contains a length C/N^d region of each interval.

Proof. Without loss of generality (by subdividing the initial intervals), let M be the greatest common divisor of all of the startpoints of the intervals in T_n . Divide each interval T_n into length $1/N$ intervals, and let $\mathbf{A} \subset (\mathbf{Z}/N)^d$ be the cartesian product of all startpoints of these length $1/N$ intervals. Since f has degree m ,

$f(\mathbf{A}) \in \mathbf{Z}/N^m$. If $A_0 \leq |\partial_0 f| \leq A_1$ on $T_0 \times \cdots \times T_d$, then for any $a \in \mathbf{A}$, and δ_0 , there exists δ_1 between 0 and δ_0 for which

$$|f(a + \delta_0 e_0) - f(a)| = \delta_0 |(\partial_0 f)(a + \delta_1)|$$

If K is fixed such that $A_1 \leq (K-1)A_0$, so that we can choose

$$\frac{1/K}{A_0 N^m} \leq \delta_0 \leq \frac{(1-1/K)}{A_1 N^m}$$

Then

$$\frac{1/K}{N^m} \leq |f(a + \delta_0 e_0) - f(a)| \leq \frac{1-1/K}{N^m}$$

Thus $d(f(\mathbf{A} + \delta_0 e_0), \mathbf{Z}/N^m) \geq 1/KN^m$. Thus if we thicken the coordinates of $\mathbf{A} + \delta_0$ to intervals of length $O(1/N^m)$, then we obtain sets S_0, \dots, S_n avoiding solutions. \square

TODO: CAN WE USE THE COMBINATORIAL NULLSTELLENSATZ TO COME UP WITH AN ALTERNATE BUILDING BLOCK LEMMA FOR ARBITRARY FIELDS?

5.8 Square Free Sets

We now look at avoiding solutions to the equation $x - y = (u - v)^2$. We consider two sets I and J . Suppose that we can select a subset \mathbf{S} from $\mathbf{Z} \cap N^2 I$ such that if $x, y \in \mathbf{S}$ are distinct, $x - y$ is not a perfect square, and $|\mathbf{S}| \gtrsim |\mathbf{Z} \cap N^2 I|^\alpha$. Then for any distinct $u, v \in \mathbf{Z} \cap N J$, $(u - v)^2 \notin \mathbf{S} - \mathbf{S}$. But this means that if we thicken the points in \mathbf{S}/N^2 to length $O(1/N^2)$ intervals, and the points in $\mathbf{Z}/N \cap J$ into length $O(1/N)$ intervals, then the resultant set will avoid solutions to $x - y = (u - v)^2$. This should give a dimension α set.

Chapter 6

Constructing Squarefree Sets

6.1 Ideas For New Work

A continuous formulation of the squarefree difference problem is not so clear to formulate, because every positive real number has a square root. Instead, we consider a problem which introduces a similar structure to avoid in the continuous domain rather than the discrete. Unfortunately, there is no direct continuous analogy to the squarefree subset problem on the interval $[0, 1]$, because there is no canonical subset of $[0, 1]$ which can be identified as ‘perfect squares’, unlike in \mathbf{Z} . If we only restrict ourselves to perfect squares of a countable set, like perfect squares of rational numbers, a result of Keleti gives us a set of full Hausdorff dimension avoiding this set. Thus, instead, we say a set $X \subset [0, 1]$ is (continuously) **squarefree** if there are no nontrivial solutions to the equation $x - y = (u - v)^2$, in the sense that there are no $x, y, u, v \in X$ satisfying the equation for $x \neq y, u \neq v$. In this section we consider some blue sky ideas that might give us what we need.

How do we adopt Rusza’s power series method to this continuous formulation of the problem? We want to scale up the problem exponentially in a way we can vary to give a better control of the exponentials. Note that for a fixed m , every elements $x \in [0, 1]$ has an essentially unique m -ary expansion

$$x = \sum_{n=1}^{\infty} \frac{x_n}{m^n}$$

and the pullback to the Haar measure on \mathbf{F}_m^{∞} is measure preserving (with respect to the natural Haar measure on \mathbf{F}_m^{∞}), so perhaps there is a way to reformulate the

problem natural as finding nice subsets of \mathbf{F}_m^∞ avoiding squares. In terms of this expansion, the equation $x - y = (u - v)^2$ can be rewritten as

$$\sum_{n=1}^{\infty} \frac{x_n - y_n}{m^n} = \left(\sum_{k=1}^{\infty} \frac{u_k - v_k}{m^k} \right)^2 = \sum_{n=1}^{\infty} \left(\sum_{k=1}^{n-1} (u_k - v_k)(u_{n-k} - v_{n-k}) \right) \frac{1}{m^n}$$

One problem with this expansion is that the sums of the differences of each element do not remain in $\{0, \dots, m-1\}$, so the sum on the right cannot be considered an equivalent formal expansion to the expansion on the left. Perhaps \mathbf{F}_m^∞ might be a simpler domain to explore the properties of squarefree subsets, in relation to Ruzsa's discrete strategy. What if we now consider the problem of finding the largest subset X of \mathbf{F}_m^∞ such that there do not exist $x, y, u, v \in \mathbf{F}_m^\infty$ such that if $x, y, u, v \in X$, $x \neq y$, $u \neq v$, then for any n

$$x_n - y_n \neq \sum_{k=1}^{n-1} (u_k - v_k)(u_{n-k} - v_{n-k})$$

What if we consider the problem modulo m , so that the convolution is considered modulo m , and we want to avoid such differences modulo m . So in particular, we do not find any solutions to the equation

$$\begin{aligned} x_2 - y_2 &= (u_1 - v_1)^2 \\ x_3 - y_3 &= 2(u_1 - v_1)(u_2 - v_2) \\ x_4 - y_4 &= (u_1 - v_1)(u_3 - v_3) + (u_2 - v_2)^2 \\ &\vdots \end{aligned}$$

which are considered modulo m . The topology of the p -adic numbers induces a power series relationship which 'goes up' and might be useful to our analysis, if the measure theory of the p -adic numbers agrees with the measure theory of normal numbers in some way, or as an alternate domain to analyze the squarefree problem as with \mathbf{F}_m^∞ .

The problem with the squarefree subset problem is that we are trying to optimize over two quantities. We want to choose a set X such that the number of distinct differences $x - y$ as small as possible, while keeping the set as large as possible. This double optimization is distinctly different from the problem of finding squarefree difference subsets of the integers. Perhaps a more natural analogy is to fix a set V , and to find the largest subset X of $[0, 1]$ such that $x - y = (u - v)^2$, where

$x \neq y \in X$, and $u \neq v \in V$. Then we are just avoiding subsets of $[0, 1]$ which avoid a particular set of differences, and I imagine this subset has a large theory. But now we can solve the general subset problem by finding large subsets X such that $(X - X)^2 \subset V$ and X containing no differences in V . Does Ruzsa's method utilize the fact that the problem is a single optimization? Can we adapt Ruzsa's method work to give better results about finding subsets X of the integers such that $X - X$ is disjoint from $(X - X)^2$?

6.2 Squarefree Sets Using Modulus Techniques

We now try to adapt Ruzsa's idea of applying congruences modulo m to avoid squarefree differences on the integers to finding high dimensional subsets of $[0, 1]$ which satisfy a continuous analogy of the integer constraint. One problem with the squarefree problem is that solutions are non-scalable, in the sense that if $X \subset [N]$ is squarefree, αX may not be squarefree. This makes sense, since avoiding solutions to $\alpha(x - y) = \alpha^2(u - v)^2$ is clearly not equivalent to the equation $x - y = (u - v)^2$. As an example, $X = \{0, 1/2\}$ is squarefree, but $2X = \{0, 1\}$ isn't. On the other hand, if X avoids squarefree differences modulo N , it is scalable by a number congruent to 1 modulo N . More generally, if α is a rational number of the form p/q , then αX will avoid nontrivial solutions to $q(x - y) = p(u - v)^2$, and if p and q are both congruent to 1 modulo N , then X is squarefree, so modulo arithmetic enables us to scale down. Since the set of rational numbers with numerator and denominator congruent to 1 is dense in \mathbf{R} , *essentially* all scales of X are continuously squarefree. Since X is discrete, it has Hausdorff dimension zero, but we can 'fatten' the scales of X to obtain a high dimension continuously squarefree set. To initially simplify the situation, we now choose to avoid nontrivial solutions to $y - x = (z - x)^2$, removing a single degree of freedom from the domain of the equation.

So we now fix a subset X of $\{0, \dots, m - 1\}$ avoiding squares modulo m . We now ask how large can we make ε such that nontrivial solutions to $x - y = (x - z)^2$ in the set

$$E = \bigcup_{x \in X} [\alpha x, \alpha x + \varepsilon)$$

occur in a common interval, if α is just short of $1/m^n$. This will allow us to recursively place a scaled, 'fattened' version of X in every interval, and then consider a limiting process to obtain a high dimensional continuously squarefree set. If we have a nontrivial solution triple, we can write it as $\alpha x + \delta_1, \alpha y + \delta_2$, and $\alpha z + \delta_3$,

with $\delta_1, \delta_2, \delta_3 < \varepsilon$. Expanding the solution leads to

$$\alpha(x-y) + (\delta_1 - \delta_2) = \alpha^2(x-z)^2 + 2\alpha(x-z)(\delta_1 - \delta_3) + (\delta_1 - \delta_3)^2$$

If x, y , and z are all distinct, then, as we have discussed, we cannot have $\alpha(x-y) = \alpha^2(x-z)^2$. if α is chosen close enough to $1/m^n$, then we obtain an approximate inequality

$$|\alpha(x-y) - \alpha^2(x-z)^2| \geq \alpha^2$$

(we require α to be close enough to $1/n$ for some n to guarantee this). Thus we can guarantee at least two of x, y , and z are equal to one another if

$$|2\alpha(x-z)(\delta_1 - \delta_3) + (\delta_1 - \delta_3)^2 - (\delta_1 - \delta_2)| < \frac{1}{m^{2n}}$$

We calculate that

$$2\alpha(x-z)(\delta_1 - \delta_3) + (\delta_1 - \delta_3)^2 - (\delta_1 - \delta_2) < 2\alpha(m-1)\varepsilon + \varepsilon^2 + \varepsilon$$

$$(\delta_1 - \delta_2) - 2\alpha(x-z)(\delta_1 - \delta_3) - (\delta_1 - \delta_3)^2 \leq \varepsilon + 2\alpha(m-1)\varepsilon$$

So it suffices to choose ε such that

$$\varepsilon^2 + [2\alpha(m-1) + 1]\varepsilon \leq \alpha^2$$

This is equivalent to picking

$$\varepsilon \leq \sqrt{\left(\frac{2\alpha(m-1) + 1}{2}\right)^2 + \alpha^2} - \frac{2\alpha(m-1) + 1}{2} \approx \frac{\alpha^2}{2\alpha(m-1) + 1}$$

We split the remaining discussion of the bound we must place on ε into the three cases where two of x, y , and z are equal, but one is distinct, to determine how small ε must be to prevent this from happening. Now

- If $y = z$, but x is distinct, then because we know $\alpha(x-y) = \alpha^2(x-y)^2$ has no solution in X , we obtain that (provided α is close enough to $1/m^n$),

$$|\alpha(x-y) - \alpha^2(x-y)^2| \geq \alpha^2$$

and the same inequality that worked for the case where the three equations are distinct now applies for this case.

- If $x = y$, but z is distinct, we are left with the equation

$$\delta_1 - \delta_2 = \alpha^2(x - z)^2 + 2\alpha(x - z)(\delta_1 - \delta_3) + (\delta_1 - \delta_3)^2$$

Now $\alpha^2(x - z)^2 \geq \alpha^2$, and

$$\delta_1 - \delta_2 - 2\alpha(x - z)(\delta_1 - \delta_3) - (\delta_1 - \delta_3)^2 < \varepsilon + 2\alpha(m - 1)\varepsilon$$

so we need the additional constraint $\varepsilon + 2\alpha(m - 1)\varepsilon \leq \alpha^2$, which is equivalent to saying

$$\varepsilon \leq \frac{\alpha^2}{1 + 2\alpha(m - 1)}$$

- If $x = z$, but y is distinct, we are left with the equation

$$\alpha(x - y) + (\delta_1 - \delta_2) = (\delta_1 - \delta_3)^2$$

Now $|\alpha(x - y)| \geq \alpha$, and

$$(\delta_1 - \delta_3)^2 - (\delta_1 - \delta_2) < \varepsilon^2 + \varepsilon$$

$$(\delta_1 - \delta_2) - (\delta_1 - \delta_3)^2 < \varepsilon$$

so to avoid this case, we need $\varepsilon^2 + \varepsilon \leq \alpha$, or

$$\varepsilon \leq \frac{\sqrt{1 + 4\alpha} - 1}{2} \approx \alpha$$

Provided ε is chosen as above, all solutions in E must occur in a common interval. Thus, if we now replace the intervals with a recursive fattened scaling of X , all solutions must occur in smaller and smaller intervals. If we choose the size of these scalings to go to zero, these solutions are required to lie in a common interval of length zero, and thus the three values must be equal to one another. Rigorously, we set $\varepsilon \approx 1/m^2$, and $\alpha \approx 1/m$, we can define a recursive construction by setting

$$E_1 = \bigcup_{x \in X} [\alpha x, \alpha x + \varepsilon_1)$$

and if we then set X_n to be the set of startpoints of the intervals in E_n , then

$$E_{n+1} = \bigcup_{x \in X_n} (x + \alpha^2 E_n)$$

Then $\cap E_n$ is a continuously squarefree subset. But what is its dimension?

6.3 Idea; Delaying Swaps

By delaying the removing in the pattern removal queue, we may assume in our dissection methods that we are working with sets with certain properties, i.e. we can swap an interval with a dimension one set avoiding translates.

6.4 Squarefree Subsets Using Interval Dissection Methods

The main idea of Keleti's proof was that, for a function f , given a method that takes a sequence of disjoint unions of sets J_1, \dots, J_N , each a union of almost disjoint closed intervals of the same length, and gives large subsets $J'_n \subset J_n$, each a union of almost disjoint intervals of a much smaller length, such that $f(x_1, \dots, x_n) \neq 0$ for $x_n \in J'_n$. Then one can find high dimensional subsets K of the real line such that $f(x_1, \dots, x_n) \neq 0$ for a sequence of distinct $x_1, \dots, x_n \in K$. The larger the subsets J'_n are compared to J_n , the higher the Hausdorff dimension of K . We now try and apply this method to construct large subsets avoiding solutions to the equation $f(x, y, z) = (x - y) - (x - z)^2$. In this case, since solutions to the equation above satisfy $y = x - (x - z)^2$, given J_1, J_2, J_3 , finding J'_1, J'_2, J'_3 as in the method above is the same as choosing J'_1 and J'_3 such that the image of $J'_1 \times J'_3$ under the map $g(x, z) = x - (x - z)^2$ is small in J_2 . We begin by discretizing the problem, splitting J_1 and J_3 into unions of smaller intervals, and then choosing large subsets of these intervals, and finding large intervals of J_2 avoiding the images of the startpoints to these intervals.

So suppose that J_1, J_2 , and J_3 are unions of intervals of length $1/M$, for which we may find subsets $A, B \subset [M]$ of the integers such that

$$J_1 = \bigcup_{a \in A} \left[\frac{a}{M}, \frac{a+1}{M} \right] \quad J_3 = \bigcup_{b \in B} \left[\frac{b}{M}, \frac{b+1}{M} \right]$$

If we split J_1 and J_3 into intervals of length $1/NM$, for some $N \gg M$ to be specified later (though we will assume it is a perfect square), then

$$J_1 = \bigcup_{\substack{a \in A \\ 0 \leq k < N}} \left[\frac{Na+k}{NM}, \frac{Na+k}{NM} + \frac{1}{NM} \right] \quad J_3 = \bigcup_{\substack{b \in B \\ 0 \leq l < N}} \left[\frac{Nb+l}{NM}, \frac{Nb+l}{NM} + \frac{1}{NM} \right]$$

We now calculate g over the startpoints of these intervals, writing

$$\begin{aligned} g\left(\frac{Na+k}{NM}, \frac{Nb+l}{NM}\right) &= \frac{Na+k}{NM} - \left(\frac{N(a-b) + (k-l)}{NM}\right)^2 \\ &= \frac{a}{M} - \frac{(a-b)^2}{M^2} + \frac{k}{NM} - \frac{2(a-b)(k-l)}{NM^2} + \frac{(k-l)^2}{(NM)^2} \end{aligned}$$

which splits the terms into their various scales. If we write $m = k - l$, then m can range on the integers in $(-N, N)$, and so, ignoring the first scale of the equation, we are motivated to consider the distribution of the set of points of the form

$$\frac{k}{NM} - \frac{2(a-b)m}{NM^2} + \frac{m^2}{(NM)^2}$$

where k is an integer in $[0, N)$, and m an integer in $(-N, N)$. To do this, fix $\varepsilon > 0$. Suppose that we find some value $\alpha \in [0, 1]$ such that S intersects

$$\left[\alpha, \alpha + \frac{1}{N^{1+\varepsilon}}\right]$$

Then there is k and m such that

$$0 \leq \frac{kNM - 2N(a-b)m + m^2}{(NM)^2} - \alpha \leq \frac{1}{N^{1+\varepsilon}}$$

Write $m = q\sqrt{N} + r$ (remember that we chose N so it's square root is an integer), with $0 \leq r < \sqrt{N}$. Then $m^2 = qN + 2qr\sqrt{N} + r^2$, and if $2qr = Q\sqrt{N} + R$, where $0 \leq R < \sqrt{N}$, then we find

$$-\frac{R}{M^2N^{3/2}} - \frac{r^2}{(NM)^2} \leq \frac{kM - 2(a-b)m + q + Q}{NM^2} - \alpha \leq \frac{1}{N^{1+\varepsilon}} - \frac{R}{M^2N^{3/2}} - \frac{r^2}{(NM)^2}$$

Thus

$$d(\alpha, \mathbf{Z}/NM^2) \leq \max\left(\frac{1}{N^{1+\varepsilon}} - \frac{R}{\sqrt{N}} - \frac{r^2}{N}, \frac{R}{M^2N^{3/2}} + \frac{r^2}{(NM)^2}\right)$$

If we now restrict our attention to the set S consisting of the expressions we are studying where $R \leq (\delta_0/2)\sqrt{N}$, $r \leq \sqrt{\delta_0 N/2}$, then if the interval corresponding to

α intersects S , then

$$d(\alpha, \mathbf{Z}/NM^2) \leq \max\left(\frac{1}{N^{1+\varepsilon}}, \frac{\delta_0}{NM^2}\right)$$

If $N^\varepsilon \geq M^2/\delta_0$, then we can force $d(\alpha, \mathbf{Z}/NM^2) \leq \delta_0/NM^2$ for all α intersecting S . Thus, if we split J_2 into intervals starting at points of the form

$$\frac{k+1/2}{NM^2}$$

each of length $1/N^{1+\varepsilon}$, then provided $\delta_0 < 1/2$, we conclude that these intervals do not contain any points in S , since

$$d\left(\frac{k+1/2}{NM^2}, \mathbf{Z}/NM^2\right) = \frac{1}{2NM^2} > \frac{\delta_0}{NM^2}$$

So we're well on our way to using Pramanik and Fraser's recursive result, since this argument shows that, provided points in J_1 and J_3 are chosen carefully, we can keep $O_M(1/N^{1+\varepsilon})$ of each interval in J_2 , which should lead to a dimension bound arbitrarily close to one.

6.5 Finding Many Startpoints of Small Modulus

To ensure a high dimension corresponding to the recursive construction, it now suffices to show J_1 and J_3 contain many startpoints corresponding to points in S , so that the refinements can be chosen to obtain $O_M(1/N)$ of each of the original intervals. Define T to be the set of all integers $m \in (-N, N)$ with $m = q\sqrt{N} + r$ and $r \leq \sqrt{\delta_0 N/2}$ and $2qr = Q\sqrt{N} + R$ with $R \leq (\delta_0/2)\sqrt{N}$. Because of the uniqueness of the division decomposition, we find T is in one to one correspondence with the set T' of all pairs of integers (q, r) , with $q \in (-\sqrt{N}, \sqrt{N})$ and $r \in [0, \sqrt{N})$, with $r \leq \sqrt{\delta_0 N/2}$, $2qr = Q\sqrt{N} + R$, and $R \leq (\delta_0/2)\sqrt{N}$. Thus we require some more refined techniques to better upper bound the size of this set.

Let's simplify notation, generalizing the situation. Given a fixed ε , We want to find a large number of integers $n \in (-N, N)$ with a decomposition $n = qr$, where $r \leq \varepsilon\sqrt{N}$, and $q \leq \sqrt{N}$. The following result reduces our problem to understanding the distribution of the smooth integers.

Lemma 26. *Fix constants A, B , and let $n \leq AN$ be an integer. If all prime factors of n are $\leq BN^{1-\delta}$, then n can be decomposed as qr with $r \leq \varepsilon\sqrt{N}$ and $q \leq \sqrt{N}$.*

Proof. Order the prime factors of n in increasing order as $p_1 \leq p_2 \leq \dots \leq p_K$. Let $r = p_1 \dots p_m$ denote the largest product of the first prime factors such that $r \leq \varepsilon \sqrt{N}$. If $r = n$, we can set $q = 1$, and we're finished. Otherwise, we know $rp_{m+1} > \varepsilon \sqrt{N}$, hence

$$r > \frac{\varepsilon \sqrt{N}}{p_{m+1}} \geq \frac{\varepsilon \sqrt{N}}{BN^{1-\delta}} = \frac{\varepsilon}{B} N^{\delta-1/2}$$

And if we set $q = n/r$, the inequality above implies

$$q < \frac{nB}{\varepsilon} N^{1/2-\delta} \leq \frac{AB}{\varepsilon} N^{3/2-\delta}$$

But now we run into a problem, because the only way we can set $q < \sqrt{N}$ while keeping A , B , and ε fixed constants is to set $\delta = 1$, and $AB/\varepsilon \leq 1$. \square

Remark. *Should we expect this method to work? Unless there's a particular reason why values of (q, r) should accumulate near $Q = 0$, we should expect to lose all but $N^{-1/2}$ of the N values we started with, so how can we expect to get $\Omega(N)$ values in our analysis. On the other hand, if a number n is suitably smooth, in a linear amount of cases we should be able to divide up primes into two numbers q and r such that r is small and q fits into a suitable value of Q , so maybe this method will still work.*

Regardless of whether the lemma above actually holds through, we describe an asymptotic formula for perfect numbers which might come in handy. If $\Psi(N, M)$ denotes the number of integers $n \leq N$ with no prime factor exceeding M , then Karl Dickman showed

$$\Psi(N, N^{1/u}) = N\rho(u) + O\left(\frac{uN}{\log N}\right)$$

This is essentially linear for a fixed u , which could show the set of (q, r) is $\Omega_\varepsilon(N)$, which is what we want. Additional information can be obtained from Hildebrand and Tenenbaum's survey paper "Integers Without Large Prime Factors".

6.6 A Better Approach

Remember that we can write a general value in our set as

$$x = \frac{-2(a-b)m}{NM^2} + \frac{q}{NM^2} + \frac{Q}{NM^2} + \frac{R}{N^{3/2}M^2} + \frac{r^2}{N^2M^2}$$

with the hope of guaranteeing the existence of many points, rather than forcing R to be small, we now force R to be close to some scaled value of \sqrt{N} ,

$$|R - n\varepsilon\sqrt{N}| = \delta\sqrt{N} \leq \varepsilon\sqrt{N}$$

Then

$$x = \frac{-2(a-b)m + q + Q + n\varepsilon + \delta}{NM^2} + \frac{r^2}{N^2M^2}$$

So

$$d(x, \mathbf{Z}/NM^2 + n\varepsilon/NM^2) \leq \frac{\varepsilon}{NM^2} + \frac{1}{4NM^2} = \frac{\varepsilon + 1/4}{NM^2}$$

By the pidgeonhole principle, since $R < \sqrt{N}$, there are $1/\varepsilon$ choices for n , whereas there are

The choice has the benefit of automatically possessing a lot of points by the pidgeonhole principle,

Chapter 7

Conclusions

Bibliography

- [1] K. Athreya. Large deviation rates for branching processes—i. single type case. *Ann. Appl. Probab.*, 4:779–790, 1994. → page 59
- [2] J. Balogh, R. Morris, and W. Samotij. Independent sets in hypergraphs. *J. Amer. Math. Soc.*, 28:669–709, 2015. → page 65
- [3] J. Biggins and N. Bingham. Large deviations in the supercritical branching process. *Adv. in Appl. Probab.*, 25:757–772. → page 59
- [4] K. Dickman. On the frequency of numbers containing prime factors of a certain relative magnitude.
- [5] K. Falconer. *Fractal Geometry: Mathematical Foundations and Applications*. Wiley, 2003. → page 53
- [6] R. Fraser and M. Pramanik. Large sets avoiding patterns. *Anal. PDE*, 11: 1083–1111, 2018. → pages 43, 44, 45, 53, 55, 56, 64, 65
- [7] T. Harris. *The Theory of Branching Processes*. Springer-Verlag, 1963. → page 58
- [8] N. H. Katz and T. Tao. Some connections between Falconer’s distance set conjecture, and sets of Furstenburg type. *New York J. Math.*, 7:149–187, 2001. → page 46
- [9] T. Keleti. A 1-dimensional subset of the reals that intersects each of its translates in at most a single point. *Real Anal. Exchange*, 24:843–845, 1999. → pages 45, 64
- [10] P. Maga. Full dimensional sets without given patterns. *Real Anal. Exchange*, 36:79–90, 2010. → pages 2, 43

- [11] A. Máthé. Sets of large dimension not containing polynomial configurations. *Adv. Math.*, 316:691–709, 2017. → pages 43, 44, 64, 65
- [12] P. Mattila. *Fourier analysis and Hausdorff dimension*. Cambridge University Press, 2015. → pages 1, 2, 43, 57
- [13] I. Z. Ruzsa. Difference sets without squares.
- [14] B. Sudakov, E. Szemerédi, and V. Vu. On a question of erdős and moser.
- [15] T. Tao. 245 C, Notes 5: Hausdorff dimension, 2009. URL <https://terrytao.wordpress.com/2009/05/19/245c-notes-5-hausdorff-dimension-optional/>.