

Cartesian Products Avoiding Patterns

by

Jacob Denson

A THESIS SUBMITTED IN PARTIAL FULFILLMENT
OF THE REQUIREMENTS FOR THE DEGREE OF

Master of Science

in

THE FACULTY OF SCIENCE

(Mathematics)

The University of British Columbia

(Vancouver)

April 2019

© Jacob Denson, 2019

The following individuals certify that they have read, and recommend to the Faculty of Graduate and Postdoctoral Studies for acceptance, the thesis entitled:

Cartesian Products Avoiding Patterns

submitted by **Jacob Denson** in partial fulfillment of the requirements for the degree of **Master of Science** in **Mathematics**.

Examining Committee:

Malabika Pramanik, Mathematics

Supervisor

Joshua Zahl, Mathematics

Supervisor

Abstract

The pattern avoidance problem seeks to construct a set $X \subset \mathbf{R}^d$ with large fractal dimension that avoids a prescribed pattern, such as three term arithmetic progressions, or more general patterns such as avoiding points $x_1, \dots, x_n \in \mathbf{R}^d$ such that $f(x_1, \dots, x_n) = 0$ for a given function f . Previous work on the subject has considered patterns described by polynomials, or functions f satisfying certain regularity conditions. We provide an exposition of some results in this setting, as well as considering new strategies to avoid ‘rough patterns’. There are several problems that fit into the framework of rough pattern avoidance. As a first application, if $Y \subset [0, 1]$ is a set with Minkowski dimension s , we construct a set $X \subset [0, 1]$ with Hausdorff dimension $1 - s$ so that $X + X$ is disjoint from Y . As a second application, given a set Y with dimension close to one, we can construct a set $X \subset Y$ of dimension $1/2$ that avoids isosceles triangles.

Lay Summary

Geometers are often interested in constructing shapes satisfying certain properties. For instance, given three points, can one find a circle connecting them? Most questions of this type involving shapes like circles and polygons have been answered. But many open questions remain about more modern families of shapes. Here, we focus on *fractals*, a class of shapes whose most well known representatives include the Koch snowflake and the Sierpinski triangle. Fractals often occur in applications such as small scale physics and computer graphics.

This thesis focuses on constructing *large* fractals which avoid the existence of certain configurations. For example, can one construct a large fractal so that one cannot form an equilateral triangle from three points contained on the fractal? We begin with an exposition providing examples of previous results achieved in the literature, and then provide new construction techniques utilizing randomness.

Preface

This thesis gives an exposition by the author, of the pattern avoidance problem and the geometric measure theory required to understand the pattern avoidance problem in the non-discrete setting. In Chapter 4, the author presents details of joint work with his supervisors Dr. Joshua Zahl and Dr. Malabika Pramanik. The results of that section have been submitted for publication in the Springer Series *Harmonic Analysis and Applications*.

Contents

| | |
|--|------------|
| Abstract | iii |
| Lay Summary | v |
| Preface | vii |
| Acknowledgments | xi |
| 1 Introduction | 1 |
| 2 Background | 3 |
| 2.1 Configuration Avoidance | 3 |
| 2.2 Minkowski Dimension | 6 |
| 2.3 Hausdorff Dimension | 7 |
| 2.4 Dyadic Scales | 10 |
| 2.5 Frostman Measures | 12 |
| 2.6 Dyadic Fractional Dimension | 17 |
| 2.7 Beyond Hyperdyadics | 23 |
| 3 Related Work | 27 |
| 3.1 Keleti: A Translate Avoiding Set | 27 |
| 3.2 Fraser/Pramanik: Smooth Configurations | 30 |
| 3.3 Mathé: Polynomial Configurations | 35 |
| 4 Avoiding Rough Sets | 39 |
| 4.1 Avoidance at Discrete Scales | 40 |
| 4.2 Fractal Discretization | 44 |

| | | |
|----------|--|-----------|
| 5 | Applications | 47 |
| 5.1 | Sum-sets avoiding specified sets | 47 |
| 5.2 | Subsets of Lipschitz curves avoiding isosceles triangles | 48 |
| 6 | Future Work | 53 |
| 6.1 | Low Rank Avoidance | 53 |
| 6.2 | Fourier Dimension | 56 |
| | Bibliography | 65 |

Acknowledgments

I am indebted to my advisors, Dr. Malabika Pramanik and Dr. Josh Zahl, for their key insights in the past two years of research. Their advice will help me conduct research for years to come. Without their tough scrutiny of my writing style over the past year, this thesis would be exponentially less legible.

I would also like to give credit to Dr. Zachary Friggstad. Our long discussions have changed the way I think about mathematics, and I look forward to many more. Your influence is felt throughout the research in this thesis.

Thanks to my family, for their support throughout my education. I would especially like to thank my grandfather, Ted McClung. Without your encouragement in my early years of undergraduate education, it is unlikely I would have found my passion for higher mathematics.

Last, but certainly not least, I'd like to thank the UBC mathematics department, and greater student community, for keeping me grounded during many stressful moments over the past two years.

Chapter 1

Introduction

In this thesis, we study a simple family of questions:

How large can Euclidean sets be not containing geometric patterns?.

For instance, what is the maximal size of a set $X \subset \mathbf{R}^d$ that contains no three collinear points? Or what is the maximal size of a set not containing the vertices of an isosceles triangle? Aside from pure geometric interest, these problems provide useful settings to test methods of ergodic theory, additive combinatorics and harmonic analysis.

Sets which avoid the patterns we consider are highly irregular, in many ways behaving like a fractal set. Our understanding of their structure requires techniques from geometric measure theory. In particular, we use various *fractal dimensions* to measure the size of a pattern avoiding set.

At present, many fundamental questions about geometric structure and its relation to fractal dimension remain unsolved. One might expect sets with sufficiently large fractal dimension must contain a given pattern. For example, Theorem 6.8 of [8] shows that any set $X \subset \mathbf{R}^d$ with Hausdorff dimension exceeding one must contain three collinear points. On the other hand, Theorem 2.3 of [5] constructs a set $X \subset \mathbf{R}^2$ with full Hausdorff dimension such that no four points in X form the vertices of a parallelogram. Thus the conjecture that sufficiently large sets must contain a given pattern depends on the particular patterns involved. For most geometric configurations, it remains unknown at what threshold patterns are guaranteed, or whether such a threshold exists at all.

Our goal in this thesis is to derive new methods for constructing sets avoiding patterns. In particular, we expand on a number of general *pattern dissection methods* which have proven useful in the area, originally developed by Keleti but also

studied notably by Fraser, Mathé, and Pramanik. In Chapter 3, we give an exposition of some of their methods, after we establish some background in Chapter 2. Our main contribution is to avoid new classes of patterns using random dissection methods. This enables us to expand the utility of pattern dissection methods from regular families of patterns to a family of *fractal avoidance problems*, that previous methods were completely unavailable to address. Such problems include finding large sets X such that the angles formed by any three distinct points of X avoid a specified set Y of angles, where Y has a fixed fractal dimension. This method is described in Chapter 4. Applications are then described in Chapter 5. Chapter 6 describes various improvements to our results which we hope to develop in the future.

Chapter 2

Background

In this chapter we discuss the required background to understand the techniques of the pattern avoidance problem. The majority of the background in geometric measure theory can be found in other resources, e.g. in [1], [7], or [10], though not geared in particular towards the pattern avoidance problem.

2.1 Configuration Avoidance

Our main focus in this thesis is the *pattern avoidance problem*. In this section, we formalize the notion of a pattern, and what it means to avoid it. Given a set \mathbf{A} , we let

$$\mathcal{C}^n(\mathbf{A}) = \{(a_1, \dots, a_n) \in \mathbf{A}^n : a_i \neq a_j \text{ if } i \neq j\}.$$

and

$$\mathcal{C}(\mathbf{A}) = \bigcup_{n=1}^{\infty} \mathcal{C}^n(\mathbf{A}).$$

We call $\mathcal{C}(\mathbf{A})$ the *configuration space* of \mathbf{A} .

Example (Non-Colinearity). We say a set $X \subset \mathbf{R}^d$ *avoids colinear points* if no three points $x_1, x_2, x_3 \in X$ lie on a common line in \mathbf{R}^d . Define

$$\mathcal{C} = \left\{ (x, x + av, x + 2av) \in \mathcal{C}^3(\mathbf{R}^d) : a \in \mathbf{R} - \{0\}, v \in \mathbf{R}^d - \{0\} \right\}.$$

Then X avoids colinear points if and only if for any $x_1, x_2, x_3 \in X$, $(x_1, x_2, x_3) \notin \mathcal{C}$.

Example (Isosceles Triangle Configuration). We say a set $X \subset \mathbf{R}^2$ *avoids isosceles triangles* if no three points $x_1, x_2, x_3 \in X$ form the vertices of a non-degenerate

isosceles triangle. Define

$$\mathcal{C} = \{(x_1, x_2, x_3) \in \mathcal{C}^3(\mathbf{R}^2) : |x_1 - x_2| = |x_1 - x_3|\}.$$

A set X avoids isosceles triangles if and only for any $x_1, x_2, x_3 \in X$, $(x_1, x_2, x_3) \notin \mathcal{C}$.

Example (Linear Independence Configuration). Let V be a vector space over a field K . We set

$$\mathcal{C} = \bigcup_{n=1}^{\infty} \left\{ (x_1, \dots, x_n) \in \mathcal{C}^n(V) : \begin{array}{l} \text{there is } a_1, \dots, a_n \in K \text{ such that} \\ a_1 x_1 + \dots + a_n x_n = 0 \end{array} \right\}.$$

A set $X \subset V$ is linearly independent in V if, for any n , and any set of n points $x_1, \dots, x_n \in X$, $(x_1, \dots, x_n) \notin \mathcal{C}$. In this thesis, we will be interested in the case where $K = \mathbf{Q}$, and $V = \mathbf{R}$.

Example (Sum Set Configuration). Let G be an abelian group, and fix $Y \subset G$. Set

$$\mathcal{C}^1 = \{g \in \mathcal{C}^1(G) : g + g \in Y\} \quad \text{and} \quad \mathcal{C}^2 = \{(g_1, g_2) \in \mathcal{C}^2(G) : g_1 + g_2 \in Y\}.$$

Define $\mathcal{C} = \mathcal{C}^1 \cup \mathcal{C}^2$. Then $(X + X) \cap Y = \emptyset$ if and only if for any $x_1, x_2 \in X$, $(x_1, x_2) \notin \mathcal{C}^2$, and for any $x \in X$, $x \notin \mathcal{C}^1$.

All the configurations we discuss in this thesis can be specified in terms of subsets of $\mathcal{C}(\mathbf{A})$. Thus we formally define a *configuration* on \mathbf{A} to be a subset of $\mathcal{C}(\mathbf{A})$, and an *n point configuration* if it is a subset of $\mathcal{C}^n(\mathbf{A})$. For a fixed configuration \mathcal{C} on \mathbf{A} , we say a set $X \subset \mathbf{A}$ *avoids* \mathcal{C} if $\mathcal{C}(X)$ is disjoint from \mathcal{C} . The *pattern avoidance problem* asks to find sets X of maximal size avoiding a fixed configuration \mathcal{C} . Often, the configuration \mathcal{C} describes algebraic or geometric structure, and the pattern avoidance problem asks to find the maximal size of a set before it is guaranteed to have such structure.

Depending on the structure of the ambient space \mathbf{A} and the configuration \mathcal{C} , there are various ways of measuring the size of sets $X \subset \mathbf{A}$ for the purpose of the pattern avoidance problem. If \mathbf{A} is finite, for instance, a natural choice is the cardinality of X . But our goal is to study pattern avoidance where $\mathbf{A} = \mathbf{R}^d$. In certain cases, one can use the Lebesgue measure to determine the size of a pattern avoiding set. But this really only works for ‘discrete’ configurations on \mathbf{R}^d , as the next theorem shows, under the often true assumption that \mathcal{C} is *translation invariant*, i.e. that if $(a_1, \dots, a_n) \in \mathcal{C}$ and $b \in \mathbf{R}^d$, $(a_1 + b, \dots, a_n + b) \in \mathcal{C}$.

Theorem 1. *Let \mathcal{C} be a n -point configuration on \mathbf{R}^d . Suppose*

(A) \mathcal{C} is translation invariant.

(B) For any $\varepsilon > 0$, there is $(a_1, \dots, a_n) \in \mathcal{C}$ with $\text{diam}\{a_1, \dots, a_n\} \leq \varepsilon$.

Then no set with positive Lebesgue measure avoids \mathcal{C} .

Proof. Let $X \subset \mathbf{R}^d$ have positive Lebesgue measure. The Lebesgue density theorem shows that there exists a point $x \in X$ such that

$$\lim_{l(Q) \rightarrow 0} \frac{|X \cap Q|}{|Q|} = 1, \quad (2.1)$$

where Q ranges over all axis-oriented cubes in \mathbf{R}^d with $x \in Q$, and $l(Q)$ denotes the sidelength of Q . Fix $\varepsilon > 0$, to be specified later, and choose r small enough that $|X \cap Q| \geq (1 - \varepsilon)|Q|$ for any cube Q with $x \in Q$ and $l(Q) \leq r$. Now let Q_0 denote a cube centered at x with $l(Q_0) \leq r$. Applying Property (B), we find $C = (a_1, \dots, a_n) \in \mathcal{C}$ such that

$$\text{diam}\{a_1, \dots, a_n\} \leq l(Q_0)/2. \quad (2.2)$$

For each $p \in Q_0$, let $C(p) = (a_1(p), \dots, a_n(p))$, where $a_i(p) = p + (a_i - a_1)$. A union bound shows

$$|\{p \in Q_0 : C(p) \notin \mathcal{C}(X)\}| \leq \sum_{i=1}^n |\{p \in Q_0 : a_i(p) \notin X\}|. \quad (2.3)$$

We have $a_i(p) \notin X$ precisely when $p + (a_i - a_1) \notin X$, so

$$|\{p \in Q_0 : a_i(p) \notin X\}| = |(Q_0 + (a_i - a_1)) \cap X^c|. \quad (2.4)$$

Note $Q_0 + (a_i - a_1)$ is a cube with the same sidelength as Q_0 . Equation (2.2) implies $|a_i - a_1| \leq l(Q_0)/2$, so $x \in Q_0 + (a_i - a_1)$. Thus (2.1) shows

$$|Q_0 + (a_i - a_1) \cap X^c| \leq \varepsilon |Q_0|. \quad (2.5)$$

Combining (2.3), (2.4), and (2.5), we find

$$|\{p \in Q_0 : C(p) \notin \mathcal{C}(X)\}| \leq \varepsilon n |Q_0|.$$

Provided $\varepsilon n < 1$, this means there is $p \in Q_0$ with $C(p) \in \mathcal{C}(X)$. Property (A) implies $C(p) \in \mathcal{C}$, so X does not avoid \mathcal{C} . \square

Since no set of positive Lebesgue measure can avoid non-discrete, translation invariant configurations, we cannot use the Lebesgue measure to quantify the size of pattern avoiding sets in this setting. Geometric measure theory provides us with a quantity which can distinguish between the size of sets of measure zero. This is the *fractal dimension* of a set.

There are many variants of fractal dimension. Here we choose to use the Minkowski dimension, and the Hausdorff dimension. They assign the same dimension to any smooth manifold, but can differ for rougher sets. One major difference is that Minkowski dimension measures relative density at a single scale, whereas Hausdorff dimension measures relative density at countably many scales.

2.2 Minkowski Dimension

Given $l > 0$, and a bounded set $E \subset \mathbf{R}^d$, we let $N(l, E)$ denote the *covering number* of E , i.e. the minimum number of sidelength l cubes required to cover E . We define the *lower* and *upper* Minkowski dimension as

$$\dim_{\underline{\mathbf{M}}}(E) = \liminf_{l \rightarrow 0} \left[\frac{\log(N(l, E))}{\log(1/l)} \right] \quad \text{and} \quad \dim_{\overline{\mathbf{M}}}(E) = \limsup_{l \rightarrow 0} \left[\frac{\log(N(l, E))}{\log(1/l)} \right].$$

If $\dim_{\underline{\mathbf{M}}}(E) = \dim_{\overline{\mathbf{M}}}(E)$, then we refer to this common quantity as the *Minkowski dimension* of E , denoted $\dim_{\mathbf{M}}(E)$. Thus $\dim_{\underline{\mathbf{M}}}(E) < s$ if there exists a sequence of lengths $\{l_k\}$ converging to zero such that for each k , E is covered by fewer than $(1/l_k)^s$ sidelength l_k cubes, and $\dim_{\overline{\mathbf{M}}}(E) < s$ if E is covered by fewer than $(1/l)^s$ sidelength l cubes for *any* suitably small l .

Remark. Any cube with sidelength r is covered in $O_d(1)$ balls of radius r . Conversely, any ball of radius r is covered by $O_d(1)$ cubes of sidelength r . Thus for any $r > 0$, if we temporarily define $N_B(r, E)$ to be the optimal number of radius r balls it takes to cover E , then $N(r, E) \sim_d N_B(r, E)$. As $r \rightarrow 0$, this means

$$\frac{\log(N(r, E))}{\log(1/r)} = \frac{\log(N_B(r, E))}{\log(1/r)} + o(1).$$

In particular, $\dim_{\underline{\mathbf{M}}}(E) < s$ if and only if there exists a sequence of lengths $\{r_k\}$ such that E is covered by $(1/r_k)^s$ radius r_k balls, and $\dim_{\overline{\mathbf{M}}}(E) < s$ if and only if E is covered by $(1/r)^s$ radius r balls, for any suitably small $r > 0$.

It is often easy to upper bound the Minkowski dimension of a set, simply by

providing a cover of the set and counting the number of cubes that cover it. Let us now consider an example. We say a set $S \subset \mathbf{R}^d$ is an s dimensional *Lipschitz manifold* if there exists a family of bounded, open subsets $\{U_\alpha\}$ of \mathbf{R}^s , together with a family of bi-Lipschitz maps $\{f_\alpha : U_\alpha \rightarrow S\}$, such that the sets $\{f_\alpha(U_\alpha)\}$ form a relatively open cover of S . Every C^1 manifold in \mathbf{R}^d is a Lipschitz manifold. This example proves useful in Chapter 4.

Theorem 2. *Let $S \subset \mathbf{R}^d$ be a Lipschitz manifold of dimension s . Then for any compact set $K \subset S$, $\dim_{\overline{\mathbf{M}}}(K) \leq s$.*

Proof. Since K is compact, we can find finitely many bi-Lipschitz maps f_1, \dots, f_N such that the family $\{f_i(U_i) : 1 \leq i \leq N\}$ covers K . Then there is C such that for each i , if $x, y \in U_i$,

$$|f_i(x) - f_i(y)| \leq C \cdot |x - y|.$$

Since each U_i is bounded, for any $r > 0$, we can find a family of balls $B_{i,1}, \dots, B_{i,M_i}$ of radius (r/C) covering U_i , such that the centers $x_{i,1}, \dots, x_{i,M_i}$ of the balls also lie in U_i , and $M_i \lesssim (C/r)^s$. But then the balls of radius r with centers lying in

$$\{f_i(x_{i,j}) : 1 \leq i \leq N, 1 \leq j \leq M_i\}$$

cover K , and there are $\sum_{i=1}^N M_i \lesssim (N/C^s)r^{-s}$ such balls. Since C and N are independent of r , and $r > 0$ was arbitrary, this shows $\dim_{\overline{\mathbf{M}}}(K) \leq s$. \square

2.3 Hausdorff Dimension

For $E \subset \mathbf{R}^d$ and $\delta > 0$, we define the *Hausdorff content*

$$H_\delta^s(E) = \inf \left\{ \sum_{k=1}^{\infty} l(Q_k)^s : E \subset \bigcup_{k=1}^{\infty} Q_k, l(Q_k) \leq \delta \right\}.$$

The s -dimensional *Hausdorff measure* of E is

$$H^s(E) = \lim_{\delta \rightarrow 0} H_\delta^s(E) = \sup \{H_\delta^s(E) : \delta > 0\}.$$

It is easy to see H^s is an exterior measure on \mathbf{R}^d , and $H^s(E \cup F) = H^s(E) + H^s(F)$ if the Hausdorff distance $d(E, F)$ between E and F is positive. So H^s is actually a metric exterior measure, and the Caratheodory extension theorem shows all Borel

sets are measurable with respect to H^s . Sometimes, it is convenient to use the exterior measure

$$H_\infty^s(E) = \inf \left\{ \sum_{k=1}^{\infty} l(Q_k)^s : E \subset \bigcup_{k=1}^{\infty} Q_k \right\}.$$

The majority of Borel sets which occur in practice fail to be measurable with respect to H_∞^s , but the exterior measure H_∞^s has the useful property that $H_\infty^s(E) = 0$ if and only if $H^s(E) = 0$.

Lemma 3. *Consider $t < s$, and $E \subset \mathbf{R}^d$.*

(i) *If $H^t(E) < \infty$, then $H^s(E) = 0$.*

(ii) *If $H^s(E) \neq 0$, then $H^t(E) = \infty$.*

Proof. Suppose that $H^t(E) = A < \infty$. Then for any $\delta > 0$, there is a cover of E by a collection of intervals $\{Q_k\}$, such that $l(Q_k) \leq \delta$ for each k , and

$$\sum_{k=1}^{\infty} l(Q_k)^t \leq A < \infty.$$

But then

$$H_\delta^s(E) \leq \sum_{k=1}^{\infty} l(Q_k)^s \leq \sum_{k=1}^{\infty} l(Q_k)^{s-t} l(Q_k)^t \leq \delta^{s-t} A.$$

As $\delta \rightarrow 0$, we conclude $H^s(E) = 0$, proving (i). And (ii) is just the contrapositive of (i), and therefore immediately follows. \square

Corollary 4. *If $s > d$, $H^s = 0$.*

Proof. The measure H^d is just the Lebesgue measure on \mathbf{R}^d , so

$$H^d[-N, N]^d = (2N)^d.$$

If $s > d$, Lemma 3 shows $H^s[-N, N]^d = 0$. By countable additivity, taking $N \rightarrow \infty$ shows $H^s(\mathbf{R}^d) = 0$. Since H^s is a positive measure, $H^s(E) = 0$ for all E . \square

Given any Borel set E , Corollary 4, combined with Lemma 3, implies there is a unique value $s_0 \in [0, d]$ such that $H^s(E) = 0$ for $s > s_0$, and $H^s(E) = \infty$ for $0 \leq s < s_0$. We refer to s_0 as the *Hausdorff dimension* of E , denoted $\dim_{\mathbf{H}}(E)$.

Theorem 5. For any bounded set E , $\dim_{\mathbf{H}}(E) \leq \dim_{\mathbf{M}}(E) \leq \dim_{\overline{\mathbf{M}}}(E)$.

Proof. Given $l > 0$, we have a simple bound $H_l^s(E) \leq N(l, E) \cdot l^s$. If $\dim_{\mathbf{M}}(E) < s$, then there exists a sequence $\{l_k\}$ with $l_k \rightarrow 0$, and $N(l_k, E) \leq (1/l_k)^s$. We conclude that

$$H^s(E) = \lim_{k \rightarrow \infty} H_{l_k}^s(E) \leq \lim_{k \rightarrow \infty} N(l_k, E) \cdot l_k^s \leq 1,$$

Thus Lemma 3 implies $\dim_{\mathbf{H}}(E) \leq s$. Taking infima over all $s > \dim_{\mathbf{M}}(E)$ shows $\dim_{\mathbf{H}}(E) \leq \dim_{\mathbf{M}}(E)$. \square

Remark. If $\dim_{\mathbf{H}}(E) < d$, then $|E| = H^d(E) = 0$. Thus any set with fractal dimension less than d (either Hausdorff or Minkowski) must have measure zero. This means we can use the dimension as a way of distinguishing between sets of measure zero, which is precisely what we need to study the configuration avoidance problem for non-discrete configurations.

The fact that Hausdorff dimension is defined over multiple scales simultaneously makes it more stable under analytical operations. In particular, for any family of at most countably many sets $\{E_k\}$,

$$\dim_{\mathbf{H}} \left\{ \bigcup E_k \right\} = \sup \{ \dim_{\mathbf{H}}(E_k) \}.$$

This need not be true for the Minkowski dimension; a single point has Minkowski dimension zero, but $\mathbf{Q} \cap [0, 1]$, which is a countable union of points, has Minkowski dimension one. An easy way to make Minkowski dimension countably stable is to define the *modified Minkowski dimensions*

$$\dim_{\mathbf{MM}}(E) = \inf \left\{ s : E \subset \bigcup_{i=1}^{\infty} E_i, \dim_{\mathbf{M}}(E_i) \leq s \text{ for each } i \right\}$$

and

$$\dim_{\overline{\mathbf{MM}}}(E) = \inf \left\{ s : E \subset \bigcup_{i=1}^{\infty} E_i, \dim_{\overline{\mathbf{M}}}(E_i) \leq s \text{ for each } i \right\}.$$

This notion of dimension, in a disguised form, appears in Chapter 4.

2.4 Dyadic Scales

It is now useful to introduce the dyadic notation we utilize throughout this thesis. At the cost of losing access to the continuous structure of \mathbf{R}^d , applying dyadic techniques often allows us to elegantly discretize problems in Euclidean space.

Fix an integer N . The classic family of dyadic cubes with *branching factor* N is given by setting, for each integer $k \geq 0$,

$$\mathcal{D}_k^d = \left\{ \prod_{i=1}^d \left[\frac{n_i}{N^k}, \frac{n_i+1}{N^k} \right] : n \in \mathbf{Z}^d \right\},$$

and then setting $\mathcal{D}^d = \bigcup_{k \geq 0} \mathcal{D}_k^d$. Elements of \mathcal{D}^d are known as *dyadic cubes*, and elements of \mathcal{D}_k^d are known as *dyadic cubes of generation k* . The most important properties of the dyadic cubes is that for each k , \mathcal{D}_k^d is a cover of \mathbf{R}^d by cubes of sidelength $1/N^k$, and for any two cubes $Q_1, Q_2 \in \mathcal{D}^d$, either their interiors are disjoint, or one cube is nested in the other.

- For each cube $Q_1 \in \mathcal{D}_{k+1}^d$, there is a unique cube $Q_2 \in \mathcal{D}_k^d$ such that $Q_1 \subset Q_2$. We refer to Q_2 as the *parent* of Q_1 , and Q_1 as a *child* of Q_2 . Each cube in \mathcal{D}^d has exactly N^d children. For $Q \in \mathcal{D}_{k+1}^d$, we let $Q^* \in \mathcal{D}_k^d$ denote its parent.
- We say a set $E \subset \mathbf{R}^d$ is \mathcal{D}_k *discretized* if it is a union of cubes in \mathcal{D}_k^d . If E is \mathcal{D}_k discretized, we define

$$\mathcal{D}_k(E) = \{Q \in \mathcal{D}_k^d : Q \subset E\}.$$

Then $E = \bigcup \mathcal{D}_k(E)$.

- Given $k \geq 0$, and $E \subset \mathbf{R}^d$, we let

$$E(1/N^k) = \bigcup \{Q \in \mathcal{D}_k^d : Q \cap E \neq \emptyset\}.$$

Then $E(1/N^k)$ is the smallest \mathcal{D}_k discretized set containing E in its interior. Our choice of notation invites thinking of $E(1/N^k)$ as a discretized version of the classic $1/N^k$ thickening

$$\{x \in \mathbf{R}^d : d(x, E) < 1/N^k\}.$$

We have no need for the standard notion of thickening in this thesis, so there is no notational conflict.

Since any cube is covered by at most $O_d(1)$ cubes in \mathcal{D} of comparable sidelength, from the perspective of geometric measure theory, working with dyadic cubes is normally equivalent to working with the class of all cubes.

Our main purpose with working with dyadic cubes is to construct *fractal-type sets*. By this, we mean defining sets X as the intersection of a nested family of sets $\{X_k\}$, where each X_k is \mathcal{D}_k discretized, and each successive set X_{k+1} is obtained from X_k by application of a simple, recursive procedure. Such a construction satisfies the following three properties of Falconer's definition of a fractal, as detailed in the introduction to [1]:

- (i) X has detail on arbitrarily small scales.
- (ii) X is too irregular to be described in traditional geometric language.
- (v) X is defined recursively.

This justifies the term 'fractal' when used to referring about these sets.

Example. Let us construct the middle thirds Cantor set C as a fractal-type set. We form C from the family of dyadic cubes with branching factor $N = 3$. We initially set $C_0 = [0, 1]$. Then, given the \mathcal{D}_k discretized set C_k , we consider each $I \in \mathcal{D}_k^1(C_k)$, and let $\mathcal{D}_{k+1}^1(I) = \{I_1, I_2, I_3\}$, where I_1, I_2, I_3 are given in increasing order with respect to their appearance in I . We set

$$C_{k+1} = \bigcup \{I_1 \cup I_3 : I \in \mathcal{D}_k^1(C_k)\}.$$

Then $C = \bigcap_{k \geq 0} C_k$ is the Cantor set.

Unfortunately, a *constant* branching factor is not sufficient to describe the types of recursive procedures we discuss in this thesis. Thus, we introduce a more general family of cubes. Instead of a single branching factor N , we fix a sequence of positive integers $\{N_k : k \geq 1\}$, with $N_k \geq 2$ for all k , which gives the branching factor at each stage of the class of cubes we define.

- For each $k \geq 0$, we define

$$\mathcal{Q}_k^d = \left\{ \prod_{i=1}^d \left[\frac{m_i}{N_1 \dots N_k}, \frac{m_i + 1}{N_1 \dots N_k} \right] : m \in \mathbf{Z}^d \right\}.$$

These are the *dyadic cubes of generation k* . We let $\mathcal{Q}^d = \bigcup_{k \geq 0} \mathcal{Q}_k^d$. Note that any two cubes \mathcal{Q}^d are either nested within one another, or their interiors are disjoint.

- We set $l_k = (N_1 \dots N_k)^{-1}$. Then l_k is the sidelength of the cubes in \mathcal{Q}_k^d .
- Given $Q \in \mathcal{Q}_{k+1}^d$, we let $Q^* \in \mathcal{Q}_k^d$ denote the *parent cube* of Q , i.e. the unique dyadic cube of generation k such that $Q \subset Q^*$.
- We say a set $E \subset \mathbf{R}^d$ is \mathcal{Q}_k *discretized* if it is a union of cubes in \mathcal{Q}_k^d . In this case, we let

$$\mathcal{Q}_k(E) = \{Q \in \mathcal{Q}_k^d : Q \subset E\}$$

denote the family of cubes whose union is E .

- For $E \subset \mathbf{R}^d$ and $k \geq 0$, we let $E(l_k) = \{Q \in \mathcal{Q}_k^d : Q \cap E = \emptyset\}$.

Sometimes, our recursive constructions need a family of ‘intermediary’ cubes that lie between the scales \mathcal{Q}_k^d and \mathcal{Q}_{k+1}^d . In this case, we consider a supplementary sequence $\{M_k : k \geq 1\}$ with $M_k \mid N_k$ for each k .

- For $k \geq 1$, we define

$$\mathcal{R}_k^d = \left\{ \prod_{i=1}^d \left[\frac{m_i}{N_1 \dots N_{k-1} M_k}, \frac{m_i + 1}{N_1 \dots N_{k-1} M_k} \right] : m \in \mathbf{Z}^d \right\}.$$

- We set $r_k = (N_1 \dots N_{k-1} M_k)^{-1}$. Then r_k is the sidelength of a cube in \mathcal{R}_k^d .
- The notions of being \mathcal{R}_k^d discretized, the collection of cubes $\mathcal{R}_k^d(E)$, and the sets $E(r_k)$, are defined as should be expected.

The cubes in \mathcal{R}_k^d are coarser than those in \mathcal{Q}_k^d , but finer than those in \mathcal{Q}_{k-1}^d .

Remark. We note that there is some notational conflict between the cubes \mathcal{D}^d and the cubes \mathcal{Q}^d , but since we never use both families simultaneously in a single argument, it should be clear which notation we are using.

2.5 Frostman Measures

It is often easy to upper bound Hausdorff dimension, but non-trivial to *lower bound* the Hausdorff dimension of a given set. A key technique to finding a lower bound is *Frostman’s lemma*, which says that a set has large Hausdorff dimension if and only if it supports a Borel measure obeying a decay law on small sets. We say a finite Borel measure μ is a *Frostman measure* of dimension s if it is

non-zero, compactly supported, and there exists $C > 0$ such that for any cube Q , $\mu(Q) \leq C \cdot l(Q)^s$. The proof of Frostman's lemma will utilize a technique often useful, known as the *mass distribution principle*. To prove the mass distribution principle, we apply weak convergence.

Lemma 6. *Suppose $\{\mu_i\}$ is a Cauchy sequence of non-negative Borel measures on \mathbf{R}^d , in the sense that for any $f \in C_c(\mathbf{R}^d)$, the sequence*

$$\left\{ \int f d\mu_i \right\}$$

is Cauchy. Then there is a regular Borel measure μ such that $\mu_i \rightarrow \mu$ weakly.

Proof. Fix a compact set K . then we can find a function $\varphi \in C_c(\mathbf{R}^d)$ such that $\mathbf{1}_K \leq \varphi$. This means that

$$\mu_i(K) = \int \mathbf{1}_K d\mu_i \leq \int \varphi d\mu_i.$$

Since $\{\int \varphi d\mu_i\}$ is Cauchy, it is uniformly bounded in i . In particular, $\mu_i(K)$ is uniformly bounded in i . Applying the Banach Alaoglu theorem, we conclude that the collection of measures $\{\mu_i|_K\}$ is contained in a compact subset of the space of non-negative Borel measures with respect to the weak $*$ topology. But every compact subset of a locally convex space is complete, and so we can therefore find a finite Borel measure μ^K supported on K such that $\mu_i|_K \rightarrow \mu^K$ weakly.

If $f \in C_c(\mathbf{R}^d)$ is supported on $K_1 \cap K_2$, for two compact sets K_1 and K_2 , then

$$\int f d\mu^{K_1} = \lim_{i \rightarrow \infty} \int_{K_1} f d\mu_i = \lim_{i \rightarrow \infty} \int_{K_2} f d\mu_i = \int f d\mu^{K_2}.$$

Thus we can define a measure μ such that for each $f \in C_c(\mathbf{R}^d)$, if f is supported on a compact set K , then

$$\int f d\mu = \int f d\mu^K.$$

For any $f \in C_c(\mathbf{R}^d)$, f is supported on some compact set K , and then

$$\int f d\mu = \int f d\mu^K = \lim_{i \rightarrow \infty} \int_K f d\mu_i = \lim_{i \rightarrow \infty} \int f d\mu_i.$$

Since f was arbitrary, $\mu_i \rightarrow \mu$ weakly. □

Theorem 7 (Mass Distribution Principle). *Let $w : \mathcal{Q}^d \rightarrow [0, \infty)$ be a function such that for any $Q_0 \in \mathcal{Q}^d$,*

$$\sum_{Q^*=Q_0} w(Q) = w(Q_0). \quad (2.6)$$

Then there exists a regular Borel measure μ supported on

$$\bigcap_{k=1}^{\infty} \left[\bigcup \{Q \in \mathcal{Q}_k^d : w(Q) > 0\} \right],$$

such that for each $Q \in \mathcal{Q}^d$,

$$\mu(Q^\circ) \leq w(Q) \leq \mu(Q), \quad (2.7)$$

and for any set E and $k \geq 0$, if $\mathcal{E}_k = \mathcal{Q}_k(E(I_k))$, then

$$\mu(E) \leq \sum_{Q \in \mathcal{E}_k} w(Q). \quad (2.8)$$

Proof. For each i , let μ_i be a regular Borel measure such that for each $j \leq i$, and $Q \in \mathcal{Q}_j^d$, $\mu_i(Q) = \mu_i(Q^\circ) = w(Q)$. One such choice is given, for each $\varphi \in C_c(\mathbf{R}^d)$, by the equation

$$\int \varphi d\mu_i = \sum_{Q \in \mathcal{Q}_i^d} \frac{w(Q)}{|Q|} \int_Q \varphi dx.$$

We claim that $\{\mu_i\}$ is a Cauchy sequence. Fix $\varphi \in C_c(\mathbf{R}^d)$, and choose some N such that φ is supported on $[-N, N]^d$. Set $Q' = [-N, N]^d$, and for each i , set $Q'_i = Q_i(Q')$. Since φ is compactly supported, φ is *uniformly continuous*, so for each $\varepsilon > 0$, if i is suitably large, there is a sequence of values $\{a_Q : Q \in \mathcal{Q}_i^d(Q')\}$ such that if $x \in Q$, $|\varphi(x) - a_Q| \leq \varepsilon$. But this means

$$\sum_{Q \in \mathcal{Q}'_i} (a_Q - \varepsilon) \mathbf{I}_Q \leq \varphi \leq \sum_{Q \in \mathcal{Q}'_i} (a_Q + \varepsilon) \mathbf{I}_Q.$$

Thus for any $j \geq i$,

$$\int \varphi d\mu_j \leq \sum_{Q \in \mathcal{Q}'_i} (a_Q + \varepsilon) \mu_j(Q) = \sum_{Q \in \mathcal{Q}'_i} (a_Q + \varepsilon) w(Q)$$

and

$$\int \phi d\mu_j \geq \sum_{Q \in \mathcal{Q}'_i} (a_Q - \varepsilon) \mu_j(Q) = \sum_{Q \in \mathcal{Q}'_i} (a_Q - \varepsilon) w(Q).$$

In particular, if $j, j' \geq i$,

$$\left| \int \phi d\mu_j - \int \phi d\mu_{j'} \right| \leq 2\varepsilon \sum_{Q \in \mathcal{Q}'_i} w(Q).$$

Repeated applications of (2.6) show that

$$\sum_{Q \in \mathcal{Q}'_i} w(Q) = \sum_{Q \in \mathcal{Q}'_0} w(Q),$$

which is therefore bounded independently of i . Since ε and ϕ were arbitrary, this shows $\{\mu_i\}$ is Cauchy.

Applying Lemma 6, we find a regular Borel measure μ such that $\mu_i \rightarrow \mu$ weakly. For any $Q \in \mathcal{Q}^d$, since Q is a closed set,

$$\mu(Q) \geq \limsup_{i \rightarrow \infty} \mu_i(Q) = w(Q),$$

and since Q° is an open set,

$$\mu(Q^\circ) \leq \liminf_{i \rightarrow \infty} \mu_i(Q^\circ) = w(Q).$$

This establishes (2.7). Conversely, if E is arbitrary, then $E \subset E(l_i)^\circ$ for any i , so if $\mathcal{E}_i = \mathcal{Q}_i(E(l_i))$, then

$$\mu(E) \leq \liminf_{i \rightarrow \infty} \mu_i(E(l_i)^\circ) \leq \liminf_{i \rightarrow \infty} \mu_i(E(l_i)) = \sum_{Q \in \mathcal{E}_i} w(Q).$$

This establishes (2.8). □

Remark. The reason why we cannot necessarily find a function μ which *precisely* extends w is that in the weak limit, mass from cubes can ‘leak’ into the mass of adjacent cubes. This is why we must use the weaker ‘extension bounds’ (2.7) and (2.8). This doesn’t cause us to ‘gain’ or ‘lose’ any mass in the weak limit, as (2.7) shows, so the result is still a ‘mass distribution’ result. It just means that the mass specified by w may be shared by adjacent cubes in the weak limit. If, for each

$$Q' \in \mathcal{Q}^d,$$

$$\lim_{k \rightarrow \infty} \sum_{Q \in \mathcal{Q}'_k} w(Q) = 0,$$

where $\mathcal{Q}'_k = \mathcal{Q}_k(Q'(l_k)) - \mathcal{Q}_k(Q')$, then (2.7) and (2.8) together imply we actually have $\mu(Q) = w(Q)$ for all $Q \in \mathcal{Q}$. Thus μ is a measure extending w . One such condition that guarantees this is that $w(Q) \lesssim l(Q)^s$ for some $s > d - 1$. Another is that for any $k \geq 0$, and distinct $Q_0, Q_1 \in \mathcal{Q}_k^d$ with $Q_0 \cap Q_1 \neq \emptyset$, either $w(Q_0) = 0$ or $w(Q_1) = 0$.

Theorem 8 (Frostman's Lemma). *If E is a Borel set, $H^s(E) > 0$ if and only if there exists an s dimensional Frostman measure supported on E .*

Proof. Suppose that μ is s dimensional and supported on E . If $H^s(F) = 0$, then for each $\varepsilon > 0$ there is a sequence of cubes $\{Q_k\}$ whose union covers F , with $\sum_{k=1}^{\infty} l(Q_k)^s \leq \varepsilon$. But then

$$\mu(F) \leq \sum_{k=1}^{\infty} \mu(Q_k) \lesssim \sum_{k=1}^{\infty} l(Q_k)^s \leq \varepsilon.$$

Taking $\varepsilon \rightarrow 0$, we conclude $\mu(F) = 0$. Thus μ is absolutely continuous with respect to H^s . Since $\mu(E) > 0$, this means that $H^s(E) > 0$.

To prove the converse, we will suppose for simplicity that E is compact. We work dyadically with the classical family of dyadic cubes \mathcal{D} , with branching factor $N = 2$. By translating, we may assume that $H^s(E \cap [0, 1]^d) > 0$, and so without loss of generality we may assume $E \subset [0, 1]^d$. For each $Q \in \mathcal{D}^d$, define $w^+(Q) = H_{\infty}^s(E \cap Q)$. Then

$$w^+(Q) \leq l(Q)^s, \tag{2.9}$$

and w^+ is subadditive. We now recursively define a function w such that for any $Q \in \mathcal{Q}^d$, (2.6) and

$$w(Q) \leq w^+(Q), \tag{2.10}$$

are satisfied at each stage of the definition of w . We initially define w by setting $w([0, 1]^d) = w^+([0, 1]^d)$. Given $Q \in \mathcal{D}_k^d$, we enumerate its children as $Q_1, \dots, Q_M \in \mathcal{D}_{k+1}^d$. We then consider any values $A_1, \dots, A_M \geq 0$ such that

$$A_1 + \dots + A_M = w(Q), \tag{2.11}$$

and for each k ,

$$A_k \leq w(Q_k). \tag{2.12}$$

This is feasible to do because $w^+(Q_1) + \dots + w^+(Q_M) \geq w^+(Q)$, and (2.10) holds for the previous definition of w . We then define $w(Q_k) = A_k$ for each k . Equation (2.11) implies (2.6) holds for this new set of definitions, and (2.12) implies (2.10) holds. Thus w is a well defined function. Furthermore, (2.11) implies (2.6) of Lemma 7, and so the mass distribution principle gives the existence of a measure μ supported on E , satisfying (2.8) and (2.7). In particular, (2.8) implies μ is non-zero. For each $Q' \in \mathcal{D}_k^d$, $\#[\mathcal{D}_k^d(Q(l_k))] = 3^d = O_d(1)$, so if we define $Q' = \mathcal{D}_k^d(Q(l_k))$, then (2.7), (2.9), and (2.10) imply

$$\mu(Q') \lesssim_d \max_{Q \in Q'} w^+(Q) \leq 1/2^{ks} = l(Q)^s.$$

where Q' ranges over the cubes in $\mathcal{D}_k^d(Q(l_k))$. Given any cube Q , we find k with $1/2^{k-1} \leq l(Q) \leq 1/2^k$. Then Q is covered by $O_d(1)$ dyadic cubes in \mathcal{D}_k^d , and so $\mu(Q) \lesssim l(Q)^s$. Thus μ is a Frostman measure of dimension s . \square

Given any finite Borel measure μ , we let

$$\dim_{\mathbf{H}}(\mu) = \{s : \mu \text{ is a Frostman measure of dimension } s\}.$$

Frostman's lemma says that for any Borel set E , $\dim_{\mathbf{H}}(E)$ is the supremum of $\dim_{\mathbf{H}}(\mu)$, over all measures μ supported on a closed subset of E . We refer to $\dim_{\mathbf{H}}(\mu)$ as the *Frostman dimension* of the measure μ .

2.6 Dyadic Fractional Dimension

It is often natural for us to establish results about fractal dimension ‘dyadically’, working with the family of cubes \mathcal{Q}^d and branching factors $\{N_k : k \geq 1\}$. We begin with Minkowski dimension. For each m , let $N_{\mathcal{Q}}(m, E)$ denote the minimal number of cubes in \mathcal{Q}_m^d required to cover E . This is often easy to calculate, up to a multiplicative constant, by greedily selecting cubes which intersect E .

Lemma 9. *For any set E ,*

$$N_{\mathcal{Q}}(m, E) \sim_d \#\{Q \in \mathcal{Q}_m^d : Q \cap E \neq \emptyset\} \sim_d N(l_m, E).$$

Proof. Let $\mathcal{E} = \{Q \in \mathcal{Q}_m^d : Q \cap E \neq \emptyset\}$. Then $N(l_m, E) \leq N_{\mathcal{Q}}(m, E) \leq \#(\mathcal{E})$. Conversely, let $\{Q_k\}$ be a minimal cover of E by cubes. Then each cube Q_k intersects at most 3^d cubes in \mathcal{Q}_m^d , so $\#(\mathcal{E}) \leq 3^d \cdot N(l_m, E) \leq 3^d \cdot N_{\mathcal{Q}}(m, E)$. \square

Thus it is natural to ask whether it is true that for any set E ,

$$\begin{aligned} \dim_{\underline{\mathbf{M}}}(E) &= \liminf_{k \rightarrow \infty} \frac{\log[N_{\mathcal{Q}}(k, E)]}{\log[1/l_k]} \\ \text{and} \\ \dim_{\overline{\mathbf{M}}}(E) &= \limsup_{k \rightarrow \infty} \frac{\log[N_{\mathcal{Q}}(k, E)]}{\log[1/l_k]}. \end{aligned} \tag{2.13}$$

The answer depends on the choice of $\{N_k\}$. In particular, a sufficient condition (and as we see later, essentially necessary) is that

$$N_{k+1} \lesssim_{\varepsilon} (N_1 \dots N_k)^{\varepsilon} \quad \text{for any } \varepsilon > 0. \tag{2.14}$$

We will see that this condition allows us to work dyadically in many scenarios when it comes to fractal dimension.

Theorem 10. *If (2.14) holds, then (2.13) holds.*

Proof. Fix a length l , and find k with $l_{k+1} \leq l \leq l_k$. Applying Lemma 9 shows

$$N(l, E) \leq N(l_{k+1}, E) \lesssim_d N_{\mathcal{Q}}(k+1, E)$$

and

$$N(l, E) \geq N(l_k, E) \gtrsim_d N_{\mathcal{Q}}(k, E).$$

Thus

$$\frac{\log[N(l, E)]}{\log[1/l]} \leq \left[\frac{\log(1/l_{k+1})}{\log(1/l_k)} \right] \frac{\log[N_{\mathcal{Q}}(k+1, E)]}{\log[1/l_{k+1}]} + O_d(1/k)$$

and

$$\frac{\log[N(l, E)]}{\log[1/l]} \geq \left[\frac{\log(1/l_k)}{\log(1/l_{k+1})} \right] \frac{\log[N_{\mathcal{Q}}(k, E)]}{\log[1/l_k]} + O_d(1/k).$$

Provided that

$$\frac{\log(1/l_{k+1})}{\log(1/l_k)} \rightarrow 1, \tag{2.15}$$

the conclusion of the theorem is true. But (2.15) is equivalent to the condition that

$$\frac{\log(N_{k+1})}{\log(N_1) + \dots + \log(N_k)} \rightarrow 0,$$

and this is equivalent to (2.14). □

Any constant branching factor satisfies (2.14) for the Minkowski dimension. In particular, we can work fairly freely with the classical dyadic cubes without any problems occurring. But more importantly for our work, we can let the sequence $\{N_k\}$ increase rapidly.

Theorem 11. *If $N_k = 2^{\lfloor 2^{k\psi(k)} \rfloor}$, where $\psi(k)$ is any decreasing sequence of positive numbers tending to zero, such that*

$$\psi(k) \geq \log_2(k)/k, \quad (2.16)$$

then (2.14) holds.

Proof. We note that $\log(N_k) = 2^{k\psi(k)} + O(1)$, and that (2.16) implies that

$$2^{\psi(1)} + \dots + 2^{k\psi(k)} \geq k.$$

Putting these two facts together, we conclude that

$$\begin{aligned} \frac{\log(N_{k+1})}{\log(N_1) + \dots + \log(N_k)} &= \frac{2^{(k+1)\psi(k+1)} + O(1)}{2^{\psi(1)} + 2^{2\psi(2)} + \dots + 2^{k\psi(k)} + O(k)} \\ &\lesssim \frac{2^{(k+1)\psi(k+1)}}{2^{\psi(k)} + 2^{2\psi(k)} + \dots + 2^{k\psi(k)}} \\ &\lesssim \frac{2^{(k+1)\psi(k+1)}}{2^{(k+1)\psi(k)}} (2^{\psi(k)} - 1) \\ &\leq (2^{\psi(k)} - 1) \rightarrow 0. \end{aligned}$$

This is equivalent to (2.14). □

We refer to any sequence $\{l_k\}$ constructed by $\{N_k\}$ satisfying (2.16) for some function ψ as a *subhyperdyadic* sequence. If a sequence $\{l_k\}$ is generated by a sequence $\{N_k\}$ such that for some fixed $c > 0$,

$$N_k = 2^{\lfloor 2^{ck} \rfloor},$$

then the values $\{l_k\}$ are referred to as *hyperdyadic*. The next (counter) example shows that hyperdyadic sequences are essentially the ‘boundary’ for sequences that can be used to measure the Minkowski dimension.

Example. We consider a multi-scale dyadic construction, utilizing the two families \mathcal{Q}^d and \mathcal{R}^d . Fix $0 \leq c < 1$, and define $N_k = 2^{\lfloor 2^{ck} \rfloor}$, and $M_k = 2^{\lfloor c2^{ck} \rfloor}$. Then

$M_k | N_k$ for each k . We recursively define a nested family of sets $\{E_k\}$, with each E_k a \mathcal{Q}_k^d discretized set, and set $E = \bigcap E_k$. We define $E_0 = [0, 1]$. Then, given E_k , for each $Q \in \mathcal{Q}_k(E_k)$, we select a *single* cube $R_Q \in \mathcal{R}_{k+1}(E_k)$, and define $E_{k+1} = \bigcup R_Q$. Then $\#(\mathcal{Q}_0(E_0)) = 1$, and

$$\#(\mathcal{Q}_{k+1}(E_{k+1})) = (N_{k+1}/M_{k+1})\#(\mathcal{Q}_k(E_k)),$$

which we can simplify to read

$$\#(\mathcal{Q}_k(E_k)) = \frac{N_1 \dots N_k}{M_1 \dots M_k}. \quad (2.17)$$

Noting that $\log(N_i) = 2^{ci} + O(1)$, and $\log(M_i) = c2^{ci} + O(1)$, we conclude that

$$\frac{\log \#(\mathcal{Q}_k(E_k))}{\log(1/l_k)} = \frac{(1-c)(2^c + \dots + 2^{ck}) + O(k)}{(2^c + \dots + 2^{ck}) + O(k)} \rightarrow 1 - c.$$

On the other hand, for each k ,

$$\#(\mathcal{R}_{k+1}^d(E_k)) = \#(\mathcal{Q}_k(E_k)) = \frac{N_1 \dots N_k}{M_1 \dots M_k},$$

and so

$$\begin{aligned} \frac{\log \#(\mathcal{R}_{k+1}^d(E_k))}{\log(1/r_{k+1})} &= \frac{(1-c)(2^c + \dots + 2^{ck}) + O(k)}{(2^c + \dots + 2^{ck}) + c2^{c(k+1)} + O(k)} \\ &= \frac{(1-c) \cdot 2^{c(k+1)} + O(k)}{(1-c + c2^c) \cdot 2^{c(k+1)} + O(k)} \\ &\rightarrow \frac{1-c}{1-c + c2^c} < 1 - c. \end{aligned}$$

In particular,

$$\dim_{\mathbf{M}}(E) \neq \liminf_{k \rightarrow \infty} \frac{\log [N(l_k, E)]}{\log(1/l_k)},$$

so measurements at hyperdyadic scales fail to establish general results about the Minkowski dimension.

We now move on to calculating Hausdorff dimension dyadically. The natural

quantity to consider is the measure defined for any set E as

$$H_Q^s(E) = \lim_{m \rightarrow \infty} H_{Q,m}^s(E),$$

where

$$H_{Q,m}^s(E) = \inf \left\{ \sum_k l(Q_k)^s : E \subset \bigcup_k Q_k, Q_k \in \bigcup_{i \geq m} \mathcal{Q}_i^d \text{ for each } k \right\}.$$

A similar argument to the standard Hausdorff measures shows there is a unique s_0 such that $H_Q^s(E) = \infty$ for $s < s_0$, and $H_Q^s(E) = 0$ for $s > s_0$. It is obvious that $H_Q^s(E) \geq H^s(E)$ for any set E , so we certainly have $s_0 \geq \dim_{\mathbf{H}}(E)$. The next lemma guarantees that $s_0 = \dim_{\mathbf{H}}(E)$, under the same conditions on the sequence $\{N_k\}$ as found in Theorem 10.

Lemma 12. *If (2.14) holds, then for any $\varepsilon > 0$, $H_Q^s(E) \lesssim_{s,\varepsilon} H^{s-\varepsilon}(E)$.*

Proof. Fix $\varepsilon > 0$ and m . Let $E \subset \bigcup Q_k$, where $l(Q_k) \leq l_m$ for each k . Then for each k , we can find i_k such that $l_{i_k+1} \leq l(Q_k) \leq l_{i_k}$. Then Q_k is covered by $O_d(1)$ elements of $\mathcal{Q}_{i_k}^d$, and

$$H_{Q,m}^s(E) \lesssim_d \sum l_{i_k}^s \leq \sum (l_{i_k}/l_{i_k+1})^s l(Q_k)^s \leq \sum (l_{i_k}/l_{i_k+1})^s l_{i_k}^\varepsilon l(Q_k)^{s-\varepsilon}. \quad (2.18)$$

By assumption,

$$l_{i_k+1} = \frac{1}{N_1 \dots N_{i_k} N_{i_k+1}} \gtrsim_{s,\varepsilon} (N_1 \dots N_{i_k})^{1+\varepsilon/s} = l_{i_k}^{1+\varepsilon/s}. \quad (2.19)$$

Putting (2.18) and (2.19) together, we conclude that $H_{Q,m}^s(E) \lesssim_{d,s,\varepsilon} \sum l(Q_k)^{s-\varepsilon}$. Since $\{Q_k\}$ was an arbitrary cover of E , we conclude $H_{Q,m}^s(E) \lesssim H^{s-\varepsilon}(E)$, and since m was arbitrary, that $H_Q^s(E) \lesssim H^{s-\varepsilon}(E)$. \square

Finally, we consider computing whether we can establish that a measure is a Frostman measure dyadically.

Theorem 13. *If (2.14) holds, and if μ is a Borel measure such that $\mu(Q) \lesssim l(Q)^s$ for each $Q \in \mathcal{Q}_k^d$, then μ is a Frostman measure of dimension $s - \varepsilon$ for each $\varepsilon > 0$.*

Proof. Given a cube Q , find k such that $l_{k+1} \leq l(Q) \leq l_k$. Then Q is covered by

$O_d(1)$ cubes in \mathcal{Q}_k^d , which shows

$$\mu(Q) \lesssim_d l_k^s = [(l_k/l)^s l^\varepsilon] l^{s-\varepsilon} \leq [l_k^{s+\varepsilon}/l_{k+1}^s] l^{s-\varepsilon} = \left[\frac{N_{k+1}^s}{(N_1 \dots N_k)^\varepsilon} \right] l^{s-\varepsilon} \lesssim_\varepsilon l^{s-\varepsilon}. \quad \square$$

Let's recognize the utility of this approach from the perspective of a dyadic construction. Suppose we have a sequence of nested sets $\{E_k\}$, where E_k is a \mathcal{Q}_k discretized subset of $[0, 1]^d$, and for each $Q_0 \in \mathcal{Q}_k(E_k)$, there is at least one cube $Q \in \mathcal{Q}_{k+1}(E_{k+1})$ with $Q^* = Q_0$. Then we can set $E = \bigcap E_k$ as a 'limit' of the discretizations E_k . We can associate with this construction a finite measure μ supported on E . It is defined by the mass distribution principle with respect to a function $w : \mathcal{Q}^d \rightarrow [0, \infty)$. We set $w([0, 1]^d) = 1$, and for each $Q \in \mathcal{Q}_{k+1}(E_{k+1})$, set

$$w(Q) = \frac{w(Q^*)}{\#\{Q' \in \mathcal{Q}_{k+1}(E_{k+1}) : (Q')^* = Q^*\}}.$$

The mass distribution principle then gives a Borel measure μ , which we refer to as the *canonical measure* associated with this construction. For this measure, it is often easy to show from a combinatorial argument that $w(Q) \lesssim l(Q)^s$ if $Q \in \mathcal{Q}^d$, which together with (2.7) also implies $\mu(Q) \lesssim l(Q)^s$ for $Q \in \mathcal{Q}^d$. This makes Theorem 13 useful.

Remark. The dyadic construction showing that Minkowski dimension cannot be measured only at hyperdyadic scales also shows that a bound on a measure μ on hyperdyadic cubes does not imply the correct bound at all scales. It is easy to show from (2.17) that the canonical measure μ for this example satisfies, for each k , each $Q \in \mathcal{Q}_k(E_k)$, and each $\varepsilon > 0$,

$$\mu(Q) = (\#\mathcal{Q}_k(E_k))^{-1} \lesssim_\varepsilon l(Q)^{1-c-\varepsilon}$$

for all $Q \in \mathcal{Q}_k(E_k)$, yet we know that for the set constructed in that example,

$$\dim_{\mathbf{H}}(E) \leq \dim_{\mathbf{M}}(E) < 1 - c.$$

Frostman's lemma implies we cannot possibly have $\mu(Q) \lesssim_\varepsilon l(Q)^{1-c-\varepsilon}$ for all $\varepsilon > 0$ and *all* cubes Q .

2.7 Beyond Hyperdyadics

If we use a faster increasing sequence of branching factors than that satisfying (2.14), we must exploit some extra property of our construction, which is not always present in general sets. Here, we rely on a *uniform mass distribution* between scales. Given the uniformity assumption, the lengths can decrease as fast as desired. We utilize the multi-scale set of dyadic cubes \mathcal{Q}^d and \mathcal{R}^d introduced in Section 2.4.

Lemma 14. *Let μ be a measure supported on a set E . Suppose that*

- (A) *For any $Q \in \mathcal{Q}_k^d$, $\mu(Q) \lesssim l_k^s$.*
- (B) *For each $R \in \mathcal{R}_{k+1}^d$, $\#[\mathcal{Q}_{k+1}^d(E(l_k) \cap R)] \lesssim 1$.*
- (C) *For any $R \in \mathcal{R}_{k+1}^d$ with parent cube $Q \in \mathcal{Q}_k^d$, $\mu(R) \lesssim (1/M_{k+1})^d \cdot \mu(Q)$.*

Then μ is a Frostman measure of dimension s .

Proof. We establish the general bound $\mu(Q) \lesssim l(Q)^s$ for all cubes Q by separating our analysis into two different cases:

- Suppose there is k with $r_{k+1} \leq l(Q) \leq l_k$. Then $\#(\mathcal{R}_{k+1}^d(Q(r_{k+1}))) \lesssim (l/r_{k+1})^d$. Properties (A) and (C) imply each of these cubes has measure at most $O((r_{k+1}/l_k)^d l_k^s)$, so we obtain that

$$\mu(Q) \lesssim (l/r_{k+1})^d (r_{k+1}/l_k)^d l_k^s = l^d / l_k^{d-s} \lesssim l^s.$$

- Suppose there exists k with $l_k \leq l \leq r_k$. Then $\#[\mathcal{R}_k^d(Q(r_k))] \lesssim 1$. Combining this with Property (C) gives $\#[\mathcal{Q}_k^d(Q(r_k) \cap E(l_k))] \lesssim 1$. This and Property (A) then shows

$$\mu(Q) \lesssim l_k^s \leq l^s.$$

We have addressed all cases, so μ is a Frostman measure of dimension s . □

This theorem is commonly used here when dealing with multi-scale dyadic fractal constructions, whose character is summarized in the following Theorem.

Theorem 15. *let $E = \bigcap E_k$, where $\{E_k\}$ is a nested family of subsets of \mathbf{R}^d , such that E_k is \mathcal{Q}_k discretized for each k . Suppose that*

(A) For each $Q \in \mathcal{Q}_k(E_k)$, there exists a set $\mathcal{R}_Q \subset \mathcal{R}_{k+1}(Q)$ such that $\#(\mathcal{R}_Q) \geq (1/2) \cdot \#(\mathcal{R}_{k+1}(Q))$, and

$$\#(\mathcal{Q}_{k+1}(R \cap E_{k+1})) = \begin{cases} 1 & : R \in \mathcal{R}_Q, \\ 0 & : R \notin \mathcal{R}_Q. \end{cases}$$

(B) For any $\varepsilon > 0$, $N_1 \dots N_k \lesssim_\varepsilon N_{k+1}^\varepsilon$.

(C) There is $0 < s \leq 1$ such that for any $\varepsilon > 0$, $N_{k+1} \lesssim_\varepsilon M_{k+1}^{(1+\varepsilon)/s}$.

Then E has Hausdorff dimension sd .

Remark. Note that Property (B) is essentially the opposite of (2.14).

Proof. Consider the function $w : \mathcal{Q}^d \rightarrow [0, \infty)$ defined with respect to the sequence $\{E_k\}$, which generates the canonical measure μ supported on E using the mass distribution principle. For each $R \in \mathcal{R}_{k+1}(E_k)$ with parent cube $Q \in \mathcal{Q}_k(E_k)$, Property (A) shows

$$w(R) \leq (2/M_{k+1}^d) f(Q). \quad (2.20)$$

Properties (B) and (C) together imply that for each $\varepsilon > 0$,

$$1/M_{k+1} \lesssim_\varepsilon r_{k+1}^{1-\varepsilon} \lesssim_\varepsilon l_{k+1}^{s(1-2\varepsilon)}. \quad (2.21)$$

If $Q \in \mathcal{Q}_{k+1}(E_{k+1})$ has parent cubes $R \in \mathcal{R}_{k+1}(E_k)$ and $Q^* \in \mathcal{Q}_k(E_k)$, then by (2.20), (2.21), and the fact that $w(Q^*) \leq 1$,

$$w(Q) = w(R) \leq (2/M_{k+1}^d) w(Q^*) \leq (2/M_{k+1}^d) \lesssim_\varepsilon l_{k+1}^{ds(1-2\varepsilon)}. \quad (2.22)$$

If Q is a cube with length l_k , then $\#(\mathcal{Q}_k^d(Q(l_k))) = O_d(1)$, which combined with (2.8), shows that

$$\mu(Q) \lesssim_\varepsilon l_{k+1}^{ds(1-2\varepsilon)}. \quad (2.23)$$

Property (B) of Lemma 14 is implied by Property (A). Equation (2.23) is a form of Property (A) in Lemma 14. Together with (2.8), (2.20) implies that the canonical measure μ satisfies Property (C) of Lemma 14. Thus all assumptions of Lemma 14 are satisfied, and so we conclude μ is a Frostman measure of dimension $ds(1 - 2\varepsilon)$ for each $\varepsilon > 0$. Taking $\varepsilon \rightarrow 0$, and applying Frostman's lemma, we conclude that the Hausdorff dimension of the support of μ , which is a subset of E , is greater

than or equal to ds . For each k , and $\varepsilon > 0$,

$$\#(\mathcal{Q}_{k+1}(E_{k+1})) \leq \#(\mathcal{R}_{k+1}(E_k)) \leq (1/M_{k+1})^d \lesssim_{\varepsilon} l_{k+1}^{ds-\varepsilon}.$$

Taking $k \rightarrow \infty$, and then $\varepsilon \rightarrow 0$, we conclude that

$$\dim_{\mathbf{H}}(E) \leq \dim_{\underline{\mathbf{M}}}(E) \leq ds.$$

Since we already know $\dim_{\mathbf{H}}(E) \geq ds$, this completes the proof. \square

Chapter 3

Related Work

Here, we discuss the main papers which influenced our results. In particular, the work of Keleti on translate avoiding sets, Fraser and Pramanik's work on sets avoiding smooth configurations, and Mathé's result on sets avoiding algebraic varieties.

3.1 Keleti: A Translate Avoiding Set

In [3], Keleti constructs a set $X \subset [0, 1]$ with Hausdorff dimension such that for each $t \neq 0$, X intersects $t + X$ in at most one place. The set X is then said to *avoid translates*. This paper contains the core idea behind the *discretization* method adapted in Fraser and Pramanik's paper. We also adapt this technique in our paper, which makes the result of interest.

Lemma 16. *Let X be a set. Then X avoids translates if and only if there do not exist values $x_1 < x_2 \leq x_3 < x_4$ in X with $x_2 - x_1 = x_4 - x_3$. In particular, a set X avoids translates if and only if it avoids the four point configuration*

$$\mathcal{C} = \{(x_1, x_2, x_3, x_4) : x_1 < x_2 \leq x_3 < x_4, x_2 - x_1 = x_4 - x_3\}.$$

Proof. Suppose $(t + X) \cap X$ contains two points $a < b$. Without loss of generality, we may assume that $t > 0$. If $a \leq b - t$, then the equation

$$a - (a - t) = t = b - (b - t)$$

shows that the tuple $(a-t, a, b-t, b)$ lies in \mathcal{C} . We also have

$$(b-t) - (a-t) = b-a,$$

so if $a-t < b-t \leq a < b$, then $(a-t, b-t, a, b) \in \mathcal{C}$. This covers all possible cases. Conversely, if there are $x_1 < x_2 \leq x_3 < x_4$ in X with

$$x_2 - x_1 = t = x_4 - x_3,$$

then $X+t$ contains $x_2 = x_1 + (x_2 - x_1)$ and $x_4 = x_3 + (x_4 - x_3)$. \square

The basic, but fundamental idea of the interval dissection technique is to introduce memory into Cantor set constructions. Keleti constructs a nested family of discrete sets $\{X_k\}$, with X_k a \mathcal{Q}_k discretized set, and with $X = \bigcap X_k$. The sequence $\{N_k\}$ will be specified later, but each N_k will be a multiple of 10. We initialize $X_0 = [0, 1]$. The novel feature of the argument is to incorporate a queuing procedure into the construction, i.e. the algorithm incorporates a list of intervals (known as a queue) that changes over the course of the algorithm, as we remove intervals from the front of the queue, and add intervals to the back of the queue. The queue initially just contains the interval $[0, 1]$, and we let $X_0 = [0, 1]$. To construct the sequence $\{X_k\}$, Keleti iteratively performs the following procedure:

Algorithm 1 Construction of the Sets $\{X_k\}$:

Set $k = 0$.

Repeat

Take off an interval I from the front of the queue.

For all $J \in \mathcal{Q}_k(X_k)$:

Order the intervals in $\mathcal{Q}_{k+1}(J)$ as J_1, \dots, J_N .

If $J \subset I$, add all intervals J_i to X_{k+1} with $i \equiv 0$ modulo 10.

Else add all J_i with $i \equiv 5$ modulo 10.

Add all intervals in \mathcal{Q}_{k+1}^d to the end of the queue.

Increase k by 1.

Each iteration of the algorithm produces a new set X_k , and so leaving the algorithm to repeat infinitely produces a sequence $\{X_k\}$ whose intersection is X .

Lemma 17. *The set X is translate avoiding.*

Proof. If X is not translate avoiding, there is $x_1 < x_2 \leq x_3 < x_4$ with $x_2 - x_1 = x_4 - x_3$. Since $l_k \rightarrow 0$, there is a suitably large integer N such that x_1 is contained in an interval $I \in \mathcal{Q}_N$ not containing x_2, x_3 , or x_4 . At stage N of the algorithm, the interval I is added to the end of the queue, and at a much later stage M , the interval I is retrieved. Find the start points $x_1^\circ, x_2^\circ, x_3^\circ, x_4^\circ \in l_M \mathbf{Z}$ to the intervals in \mathcal{Q}_M containing x_1, x_2, x_3 , and x_4 . Then we can find n and m such that $x_4^\circ - x_3^\circ = (10n)l_M$, and $x_2^\circ - x_1^\circ = (10m + 5)l_M$. In particular, this means that $|(x_4^\circ - x_3^\circ) - (x_2^\circ - x_1^\circ)| \geq 5L_M$. But

$$\begin{aligned} |(x_4^\circ - x_3^\circ) - (x_2^\circ - x_1^\circ)| &= |[(x_4^\circ - x_3^\circ) - (x_2^\circ - x_1^\circ)] - [(x_4 - x_3) - (x_2 - x_1)]| \\ &\leq |x_1^\circ - x_1| + \cdots + |x_4^\circ - x_4| \leq 4L_M \end{aligned}$$

which gives a contradiction. \square

It is easy to see from the algorithm that

$$\#(\mathcal{Q}_k(X_k)) = (N_k/10) \cdot \#(\mathcal{Q}_{k-1}(X_{k-1})).$$

Closing the recursive definition shows

$$\#(\mathcal{Q}_k(X_k)) = \frac{1}{10^k l_k}.$$

In particular, this means $|X_k| = 1/10^k$, so $|X|$ has Lebesgue measure zero regardless of how we choose the parameters $\{N_k\}$.

Theorem 18. *For some sequence $\{N_k\}$, the set X has full Hausdorff dimension.*

Proof. Set $N_k = 10M_k$ for each $k \geq 1$, then one sees that for each $R \in \mathcal{R}_k(X_k)$, $\#(\mathcal{Q}_{k+1}(R \cap X_{k+1})) = 1$. Thus Property (A) of Theorem 15 is satisfied. Furthermore, Property (C) of Theorem 15 is satisfied with $s = 1$. We conclude that if $N_1 \dots N_k \lesssim_\varepsilon N_{k+1}^\varepsilon$, then all assumptions of Theorem 15 is satisfied, and we conclude X is a set with full Hausdorff dimension. This is true, for instance, if we set $N_k = 2^{2^{\lfloor k \log k \rfloor}}$. \square

The most important feature of Keleti's argument is his reduction of a non-discrete configuration avoidance problem to a sequence of discrete avoidance problems on cubes. Let us summarize the result of Keleti's discrete argument in a lemma.

Lemma 19. *Let $T_1, T_2 \subset \mathbf{R}$ be disjoint, \mathcal{Q}_k discretized sets. If $M_{k+1} = N_{k+1}/10$, then we can find $S_1 \subset T_1$ and $S_2 \subset T_2$ such that*

- (i) *For each k , S_k is a \mathcal{Q}_{k+1} discretized subset of T_k .*
- (ii) *If $x_1 \in S_1$ and $x_2, x_3, x_4 \in S_2$, then $x_2 - x_1 \neq x_4 - x_3$.*
- (iii) *For each i , and each cube $R \in \mathcal{R}_{k+1}(S_i)$, $\#(\mathcal{Q}_{k+1}(R \cap S_i)) = 1$.*

Property (i) of Lemma 19 allows Keleti to apply his argument iteratively at each dyadic scale. Property (ii) implies Keleti obtains a configuration avoiding set by iterative the argument infinitely many times. And the reason why Keleti obtains a set with full Hausdorff dimension, relating back to Property (C) in Theorem 15, is because of Property (iii), which allows us to select $N_{k+1} \lesssim M_{k+1}$.

Of course, it is not possible to extend the discrete solution of the configuration argument to general configurations; this part of Keleti's method strongly depends on the arithmetic structure of the configuration. The iterative application of a discrete solution, however, can be applied in generality. Combined with Theorem 15, this technique gives a powerful method to reduce configuration avoidance problems about Hausdorff dimension to discrete avoidance problems on cubes.

3.2 Fraser/Pramanik: Smooth Configurations

Inspired by Keleti's result, in [2], Pramanik and Fraser obtained a generalization of the queue method which allows one to find sets avoiding $n+1$ point configurations given by the zero sets of smooth functions, i.e.

$$\mathcal{C} = \{(x_1, \dots, x_n) \in \mathcal{C}^n[0, 1]^d : f(x_0, \dots, x_n) = 0\},$$

under mild regularity conditions on the function $f : \mathcal{C}^n[0, 1]^d \rightarrow [0, 1]^m$.

Theorem 20 (Pramanik and Fraser). *Fix $m \leq d(n-1)$. Consider a countable family of C^2 functions $\{f_k : [0, 1]^{dn} \rightarrow [0, 1]^m\}$ such that for each k , Df_k has full rank at any $(x_1, \dots, x_n) \in \mathcal{C}^n[0, 1]^d$ where $f_k(x_1, \dots, x_n) = 0$. Then there exists a set $X \subset \mathbf{R}^d$ with Hausdorff dimension $m/(n-1)$ such that X avoids the configuration*

$$\mathcal{C} = \bigcup_k \{(x_1, \dots, x_n) \in \mathcal{C}^n(\mathbf{R}^d) : f_k(x_1, \dots, x_n) = 0\}.$$

Remark. For simplicity, we only prove the result for a single function, rather than a countable family of functions. The only major difference between the two

approaches is the choice of scales we must choose later on in the argument, and a slight modification of the queuing argument.

Just like Keleti, Pramanik and Fraser begin by solving a discrete variant of the configuration problem in Theorem 20, which they can then iteratively apply at each scale. In the discrete setting, rather than making a linear shift in one of the variables, as in Keleti's approach, Pramanik and Fraser must utilize the smoothness properties of the function to find large segments of an interval avoiding. Corollary 23 gives the discrete result that Pramanik and Fraser utilize in a queuing construction to find a set X satisfying the conclusions of Theorem 20. Here, a multi-scale approach proves useful, so we utilize the family of cubes \mathcal{Q} and \mathcal{R} , assuming the existence of two sequences of branching factors $\{N_k\}$ and $\{M_k\}$ which we will specify later on in the argument.

Lemma 21. *Fix $n > 1$. Let $T \subset [0, 1]^d$ and $T' \subset [0, 1]^{(n-1)d}$ be \mathcal{Q}_k discretized sets. Let $B \subset T \times T'$ be \mathcal{Q}_{k+1} discretized. Then there exists a \mathcal{Q}_{k+1} discretized set $S \subset T$, and a \mathcal{Q}_{k+1} discretized set $B' \subset T'$, such that*

$$(A) \quad (S \times T') \cap B \subset S \times B'.$$

$$(B) \quad \text{For every } Q \in \mathcal{Q}_k^d, \text{ there exists } \mathcal{R}(Q) \subset \mathcal{R}_{k+1}^d(Q), \text{ such that}$$

$$\#(\mathcal{R}(Q)) \geq (1/2) \cdot \#(\mathcal{R}_{k+1}^d(Q)),$$

$$\text{and for each } R \in \mathcal{R}_{k+1}^d(Q),$$

$$\#(\mathcal{Q}_{k+1}(R)) = \begin{cases} 1 & : R \in \mathcal{R}(Q), \\ 0 & : R \notin \mathcal{R}(Q). \end{cases}$$

$$(C) \quad \#(\mathcal{Q}_{k+1}(B')) \leq 2(N_1 \dots N_k)^d (M_{k+1}/N_{k+1})^d \cdot \#(\mathcal{Q}_{k+1}(B)).$$

Proof. Fix $Q_0 \in \mathcal{Q}_k(T)$. For each $R \in \mathcal{R}_{k+1}(Q_0)$, define a *slab* $S[R] = R \times T'$, and for each $Q \in \mathcal{Q}_{k+1}(Q_0)$, define a *wafer* $W[Q] = Q \times T'$. We say a wafer $W[Q]$ is *good* if

$$\#(\mathcal{Q}_{k+1}(W[Q] \cap B)) \leq (2/N_{k+1}^d) \cdot \#(\mathcal{Q}_{k+1}(B)). \quad (3.1)$$

Then at most $N_{k+1}^d/2$ wafers are bad. We call a slab *good* if it contains a wafer which is good. Since a slab is the union of $(N_{k+1}/M_{k+1})^d$ wafers, at most $M_{k+1}^d/2 = (1/2) \cdot \#(\mathcal{R}_{k+1}(Q_0))$ slabs are bad. Thus if we set

$$\mathcal{R}(Q_0) = \{R \in \mathcal{R}_{k+1}(Q_0) : S[R] \text{ is good}\},$$

then

$$\#(\mathcal{R}(Q_0)) \geq (1/2) \cdot \#(\mathcal{R}_{k+1}(Q_0)). \quad (3.2)$$

For each $R \in \mathcal{R}(Q_0)$, we pick $Q_R \in \mathcal{Q}_{k+1}(R)$ such that $W[Q_R]$ is good, and define

$$S = \bigcup \{Q_R : R \in \mathcal{R}(Q_0)\}.$$

Equation (3.2) implies S satisfies Property (B).

Let B' be the union of all cubes $Q' \in \mathcal{Q}_{k+1}(T')$ such that there is $Q \in \mathcal{Q}_{k+1}(S)$ with $Q \times Q' \in \mathcal{Q}_{k+1}(B)$. By definition, Property (A) is then satisfied. For each $Q \in \mathcal{Q}_{k+1}(S)$, $W[Q]$ is good, so (3.1) implies

$$\#\{Q' : Q \times Q' \in \mathcal{Q}_{k+1}(B)\} \leq (2/N_{k+1}^d) \cdot \#(\mathcal{Q}_{k+1}(B)).$$

But $\#(\mathcal{Q}_{k+1}(S)) \leq \#(\mathcal{R}_{k+1}(T)) \leq (1/r_{k+1})^d = (N_1 \dots N_k)^d M_{k+1}^d$, so

$$\begin{aligned} \#(\mathcal{Q}_{k+1}(B')) &\leq \#(\mathcal{Q}_{k+1}(S))[(2/N_{k+1}^d) \cdot \#(\mathcal{Q}_{k+1}(B))] \\ &\leq 2(N_1 \dots N_k)^d (M_{k+1}/N_{k+1})^d \#(\mathcal{Q}_{k+1}(B)), \end{aligned}$$

which establishes Property (C). \square

We apply the lemma recursively $n - 1$ times to continually reduce the dimensionality of the avoidance problem we are considering. Eventually, we obtain the case where $n = 0$, and then avoiding the configuration is easy.

Lemma 22. *Fix $n > 1$. Let $T \subset [0, 1]^d$ be \mathcal{Q}_k discretized, and let $B \subset T$ be \mathcal{Q}_{k+1} discretized. Suppose*

$$\# \mathcal{Q}_{k+1}(B) \leq \left[C \cdot 2^{n-1} (N_1 \dots N_k)^{d(n-1)} (M_{k+1}/N_{k+1})^{d(n-1)} \right] (1/l_{k+1})^{dn-m}.$$

and

$$N_{k+1} \geq \left[C \cdot 2^n (N_1 \dots N_k)^{2dn} \right]^{1/m} M_{k+1}^{d(n-1)/m}. \quad (3.3)$$

Then there exists a \mathcal{Q}_{k+1} discretized set $S \subset T$ such that

$$(A) \ S \cap B = \emptyset.$$

$$(B) \ \text{For each } Q_0 \in \mathcal{Q}_k(T), \text{ there is } \mathcal{R}(Q_0) \subset \mathcal{R}_{k+1}(Q_0) \text{ with}$$

$$\#(\mathcal{R}(Q_0)) \geq (1/2) \#(\mathcal{R}_{k+1}(Q_0)),$$

such that for each $R \in \mathcal{R}_{k+1}(Q_0)$,

$$\#(\mathcal{Q}_{k+1}(R \cap S)) = \begin{cases} 1 & : R \in \mathcal{R}(Q_0), \\ 0 & : R \notin \mathcal{R}(Q_0). \end{cases}$$

Proof. For each $Q_0 \in \mathcal{Q}_k(T)$, we set

$$\mathcal{R}(Q_0) = \{R \in \mathcal{R}_{k+1}(Q_0) : \#(\mathcal{Q}_{k+1}(R \cap B)) \leq (2/M_{k+1}^d) \cdot \#(\mathcal{Q}_{k+1}(B))\}.$$

Since $\mathcal{R}_{k+1}(Q_0) = M_{k+1}^d$,

$$\#(\mathcal{R}(Q_0)) \geq \#(\mathcal{R}_{k+1}(Q_0)) - (M_{k+1}^d/2) \geq (1/2) \cdot \#(\mathcal{R}_{k+1}(T)).$$

Now (3.3) implies that for each $R \in \mathcal{R}(Q_0)$,

$$\begin{aligned} \#(\mathcal{Q}_{k+1}(R \cap B)) &\leq (2/M_{k+1}^d) \cdot \#(\mathcal{Q}_{k+1}(B)) \\ &\leq (2/M_{k+1}^d) \left(C \cdot 2^{n-1} (N_1 \dots N_k)^{2dn} (M_{k+1}/N_{k+1})^{d(n-1)} \right) \\ &= \left[2^n C (N_1 \dots N_k)^{2dn} \right] \left(M_{k+1}^{d(n-2)} / N_{k+1}^{m-d} \right) \\ &< (N_{k+1}/M_{k+1})^d \\ &= \mathcal{Q}_{k+1}(R) \end{aligned}$$

Thus for each $R \in \mathcal{R}(Q_0)$, we can find $Q_R \in \mathcal{Q}_{k+1}(R)$ such that $Q_R \cap B = \emptyset$. And so if we set

$$S = \bigcup \{Q_R : R \in \mathcal{R}(Q_0), Q_0 \in \mathcal{Q}_k(T)\},$$

then (A) and (B) are satisfied. \square

Corollary 23. *Let $f : [0, 1]^{dn} \rightarrow [0, 1]^m$ be C^2 , and have full rank at every point $(x_1, \dots, x_n) \in \mathcal{C}^n(\mathbf{R}^d)$ such that $f(x_1, \dots, x_n) = 0$. Then there exists a universal constant C depending only on f such that, if (3.3) is satisfied, then for any disjoint, \mathcal{Q}_k discretized sets $T_1, \dots, T_n \subset [0, 1]^d$, we can find \mathcal{Q}_{k+1} discretized sets $S_1 \subset T_1, \dots, S_n \subset T_n$ such that*

(A) *If $x_1 \in S_1, \dots, x_n \in S_n$, then $f(x_1, \dots, x_n) \neq 0$.*

(B) *For each k , and for each $Q_0 \in \mathcal{Q}_k(T_k)$, there is $\mathcal{R}(Q_0) \subset \mathcal{R}_{k+1}(Q_0)$ with*

$$\#(\mathcal{R}(Q_0)) \geq (1/2) \cdot \#(\mathcal{R}_{k+1}(Q_0)),$$

and for each $R \in \mathcal{R}_{k+1}(Q_0)$,

$$\#(\mathcal{Q}_{k+1}(R \cap S)) = \begin{cases} 1 & : R \in \mathcal{R}(Q_0), \\ 0 & : R \notin \mathcal{R}(Q_0). \end{cases}$$

Proof. Since f is C^2 and has full rank on the set

$$V(f) = \{(x_1, \dots, x_n) \in \mathcal{C}^n(\mathbf{R}^d) : f(x_1, \dots, x_n) = 0\},$$

the implicit function theorem implies $V(f)$ is a smooth manifold of dimension $nd - m$ in \mathbf{R}^{dn} , and the co-area formula implies the existence of a constant C such that for each k ,

$$\#\{Q \in \mathcal{Q}_k^{dn} : Q \cap V(f) \neq \emptyset\} \leq C/l_k^{dn-m}.$$

To apply Lemma 21 and 22, we set

$$B = \#\{Q \in \mathcal{Q}_k^{dn} : Q \cap V(f) \neq \emptyset\}.$$

Applying Lemma 21 iteratively $n - 1$ times, then finishing with an application of Lemma 22 constructs the required sets S_1, \dots, S_n . \square

Just like in Keleti's proof, Pramanik and Fraser's technique applies a discrete result, Corollary 23, iteratively at many scales, with the help of a queuing process, to obtain a high dimensional set avoiding the zeros of a function. We construct a nested family $\{X_k : k \geq 0\}$ of \mathcal{Q}_k discretized sets, converging to a set X , which we will show is translate avoiding. We initialize $X_0 = [0, 1]$. Our queue shall consist of n tuples of disjoint intervals (T_1, \dots, T_n) , all of the same length, which initially consists of all possible tuples of intervals in $\mathcal{Q}_1^d([0, 1]^d)$. To construct the sequence $\{X_k\}$, we perform the following iterative procedure:

Algorithm 2 Construction of the Sets $\{X_k\}$

Set $k = 0$

Repeat

Take off an n tuple (T'_1, \dots, T'_n) from the front of the queue

Set $T_i = T'_i \cap X_k$ for each i

Apply Corollary 23 to the sets T_1, \dots, T_d , obtaining \mathcal{Q}_{k+1} discretized sets S_1, \dots, S_n satisfying Properties (A), (B), and (C) of that Lemma.

Set $X_{k+1} = X_k - \bigcup_{i=1}^n T_i - S_i$.

Add all n tuples of disjoint cubes (T'_1, \dots, T'_n) in $\mathcal{Q}_{k+1}^d(X_{k+1})$ to the back of the queue.

Increase k by 1.

Lemma 24. *The set X constructed by the procedure avoids the configuration*

$$\mathcal{C} = \{(x_1, \dots, x_n) \in \mathcal{C}^n(\mathbf{R}^d) : f(x_1, \dots, x_n) = 0\}.$$

Proof. Suppose $x_1, \dots, x_n \in X$ are distinct. Then at some stage k , x_1, \dots, x_n lie in disjoint cubes $T'_1, \dots, T'_n \in \mathcal{Q}_k^d(X_k)$, for some large k . At this stage, (T'_1, \dots, T'_n) is added to the back of the queue, and therefore, at some much later stage N , the tuple (T'_1, \dots, T'_n) is taken off the front. Sets $S_1 \subset T'_1, \dots, S_n \subset T'_n$ are constructed satisfying Property (A) of Corollary 23. Since $x_1, \dots, x_n \in X$, we must have $x_i \in S_i$ for each i , so $f(x_1, \dots, x_n) \neq 0$. \square

What remains is to show that for some sequence of parameters $\{N_k\}$ and $\{M_k\}$ satisfying (3.3), we can apply Theorem 15. The conclusion of Corollary 23 shows Property (A) of Theorem 15 is always satisfied. If we set

$$N_{k+1} = \left\lceil \left[C \cdot 2^n (N_1 \dots N_k)^{2dn} \right]^{1/m} M_{k+1}^{d(n-1)/m} \right\rceil,$$

then (3.3) holds, so we can apply Lemma 20 at each scale. Properties (B) and (C) of Theorem 15 are satisfied with $s = m/d(n-1)$ if $N_1 \dots N_k \lesssim_\varepsilon M_{k+1}^\varepsilon$ for each $\varepsilon > 0$. This is true, for instance, if we set $M_k = 2^{\lfloor 2^{k \log k} \rfloor}$, and Theorem 15 then shows the resultant set X has Hausdorff dimension $m/(n-1)$.

3.3 Math  : Polynomial Configurations

Math  's result [6] constructs sets avoiding low degree algebraic hypersurfaces.

Theorem 25 (Mathé). *Let $\{f_k : \mathbf{R}^{nd} \rightarrow \mathbf{R}\}$ be a countable family of rational coefficient polynomials with degree at most m . Then there exists a set $X \subset [0, 1]^d$ with Hausdorff dimension d/m which avoids the configuration*

$$\mathcal{C} = \bigcup_k \{(x_1, \dots, x_n) \in \mathcal{C}^n(\mathbf{R}^d) : f_k(x_1, \dots, x_n) = 0\}.$$

Originally, Mathé's result does not explicitly use a discretization method analogous to Keleti and Pramanik and Fraser, but his proof strategy can be reconfigured to work in this setting. For the purpose of brevity, we do not carry out the complete argument, merely giving the discretization method below. By first trying to avoid the zero sets of the partial derivatives of the function f , one can reduce to the case where a partial derivative of f is non-vanishing on the \mathcal{Q}_k discretized sets we start with. The key technique is that f maps discrete lattices of points to a discrete, 'one dimensional lattice' in \mathbf{R}^d , and the degree of the polynomial gives us the difference in lengths between the two lattices. This idea was present in Keleti's work, and one can view Mathé's result as a generalization along these lines. We revisit this idea in Chapter 6. As in Pramanik and Fraser's result, we utilize the multi-scale dyadic notations \mathcal{Q} and \mathcal{R} , with an implicitly chosen sequence $\{N_k\}$ and $\{M_k\}$.

Theorem 26. *Let $f : [0, 1]^{dn} \rightarrow \mathbf{R}$ be a rational coefficient polynomial of degree m , and disjoint, \mathcal{Q}_k discretized sets $T_1, \dots, T_n \subset [0, 1]^d$, such that $\inf |\partial_1 f| \neq 0$ on $T_1 \times \dots \times T_n$. Then there exists a constant C , depending only on f , such that if*

$$N_{k+1} \geq C \cdot (N_1 \dots N_k)^{m-1} \cdot M_{k+1}^m, \quad (3.4)$$

then there exists \mathcal{Q}_{k+1} discretized sets $S_1 \subset T_1, \dots, S_n \subset T_n$ such that

(A) $f(x) \neq 0$ for $x \in S_1 \times \dots \times S_n$.

(B) For each i , and for each $R \in \mathcal{R}_{k+1}^d(T_i)$, $\#(\mathcal{Q}_{k+1}^d(R \cap S_i)) = 1$.

Proof. Without loss of generality, by considering an appropriate integer multiple of f , we may assume f has integer coefficients. Let $\mathbf{A} \subset (r_{k+1} \cdot \mathbf{Z})^d$. Since f has degree m , $f(\mathbf{A}) \subset r_{k+1}^m \cdot \mathbf{Z}$. Suppose that $c_0 \leq |\partial_1 f| \leq C_0$ on $T_1 \times \dots \times T_n$. Then the mean value theorem guarantees that if $\delta \leq r_{k+1}$, then

$$c_0 \delta \leq |f(a + \delta e_1) - f(a)| \leq C_0 \delta.$$

Pick $\varepsilon \in (0, 1)$ small enough that $\varepsilon/c_0 < (1 - \varepsilon)/C_0$. If

$$l_{k+1} \leq [(1 - \varepsilon)/C_0 - \varepsilon/c_0]r_{k+1}^m \quad (3.5)$$

we can find $\delta \in l_{k+1} \cdot \mathbf{Z}$ is such that

$$[\varepsilon/c_0]r_{k+1}^m \leq \delta \leq [(1 - \varepsilon)/C_0]r_{k+1}^m.$$

Equation (3.5) is guaranteed by (3.4) for a sufficiently large constant C . We then find $d(f(\mathbf{A} + \delta e_1), r_{k+1}^m \cdot \mathbf{Z}) \geq \varepsilon r_{k+1}^m$. For $i > 1$, define

$$S_i = \bigcup [a_1, a_1 + l_{k+1}] \times \cdots \times [a_d, a_d + l_{k+1}],$$

where $a = (a_1, \dots, a_d)$ ranges over all start points to intervals

$$[a_1, a_1 + r_{k+1}] \times \cdots \times [a_d, a_d + r_{k+1}] \in \mathcal{R}_{k+1}^d(T_i).$$

Similarly, define

$$S_1 = \bigcup [a_1 + \delta e_1, a_1 + \delta e_1 + s] \times [a_2, a_2 + l_{k+1}] \times \cdots \times [a_d, a_d + l_{k+1}],$$

where a ranges over all start points to intervals

$$[a_1, a_1 + r_{k+1}] \times \cdots \times [a_d, a_d + r_{k+1}] \in \mathcal{R}_{k+1}^d(T_1).$$

Because f is C^1 , we can find C_2 such that if $x, y \in \mathbf{R}^d$ satisfy $|x_i - y_i| \leq t$ for all i , then $|f(x) - f(y)| \leq C_2 t$. But then for a sufficiently large constant C , (3.4) implies that

$$d(f(S_1 \times \cdots \times S_n), r_{k+1}^m \cdot \mathbf{Z}) \geq \varepsilon r_{k+1}^m - C_2 l_{k+1} \geq (\varepsilon/2)r_{k+1}^m.$$

In particular, this implies Property (A). □

Chapter 4

Avoiding Rough Sets

In the previous chapter, we saw that many authors have considered the pattern avoidance problem for configurations \mathcal{C} which take the form of many general classes of smooth shapes; in Mathé's work, \mathcal{C} can take the form of an algebraic variety of low degree, and in Pramanik and Fraser's work, \mathcal{C} can take the form of a smooth manifold. In this chapter, we consider the pattern avoidance problem for an even more general class of 'rough' patterns, that are the countable union of sets with controlled lower Minkowski dimension.

Theorem 27. *Let $s \geq d$, and suppose $\mathcal{C} \subset \mathcal{C}^n(\mathbf{R}^d)$ is the countable union of pre-compact sets, each with lower Minkowski dimension at most s . Then there exists a set $X \subset [0, 1]^d$ with Hausdorff dimension at least $(nd - s)/(n - 1)$ avoiding \mathcal{C} .*

Remarks.

1. When $s < d$, avoiding the configuration \mathcal{C} is trivial. If we define $\pi : \mathcal{C}^n(\mathbf{R}^d) \rightarrow \mathbf{R}^d$ by $\pi(x_1, \dots, x_n) = x_1$, then the set $X = [0, 1]^d - \pi(\mathcal{C})$ is full dimension and avoids \mathcal{C} . Note that obtaining a full dimensional set in the case $s = d$, however, is still interesting.
2. Theorem 27 is trivial when $s = dn$, since we can set $X = \emptyset$. We will therefore assume that $s < dn$ in our proof of the theorem.
3. Let $f : \mathbf{R}^{dn} \rightarrow \mathbf{R}^m$ be a C^1 map such that f has full rank at any point $(x_1, \dots, x_n) \in \mathcal{C}^n(\mathbf{R}^d)$ with $f(x_1, \dots, x_n) = 0$. If we set

$$\mathcal{C} = \{x \in \mathcal{C}^n(\mathbf{R}^d) : f(x) = 0\},$$

Then \mathcal{C} is a C^1 submanifold of $\mathcal{C}^n(\mathbf{R}^d)$ of dimension $nd - m$. A submanifold of Euclidean space is σ compact, so we can write $\mathcal{C} = \bigcup K_i$, where each K_i is a compact set. Applying Theorem 2 shows $\dim_{\mathbf{M}}(K_i) \leq nd - m$ for each i , so we can apply Theorem 27 with $s = nd - m$ to yield a set in \mathbf{R}^d with Hausdorff dimension at least

$$\frac{nd - s}{n - 1} = \frac{m}{n - 1}.$$

This recovers Theorem 20, making Theorem 27 a generalization of Pramanik and Fraser's result.

4. Since Theorem 27 does not require any regularity assumptions on the set \mathcal{C} , it can be applied in contexts that cannot be addressed using previous methods in the literature. Two such applications, new to the best of our knowledge, have been recorded in Chapter 5; see Theorems 31 and 32 there.

Like with the results considered in the last chapter, we construct the set X in Theorem 27 by repeatedly applying a discrete avoidance result at the scales corresponding to cubes \mathcal{Q}^d , constructing a Frostman measure with equal mass at intermediary scales, and applying Lemma 14. However, our method has several innovations that simplify the analysis of the resulting set $X = \bigcap X_k$ than from previous results. In particular, through a probabilistic selection process we are able to use a simplified queuing technique then that used in [3] and [2], that required storage of data from each step of the iterated construction to be retrieved at a much later stage of the construction process.

4.1 Avoidance at Discrete Scales

In this section we describe a method for avoiding a discretized version of \mathcal{C} at a single scale. We apply this technique in Section 4.2 at many scales to construct a set X avoiding \mathcal{C} at all scales. In the discrete setting, \mathcal{C} is replaced by a union of cubes in \mathcal{Q}_{k+1}^{dn} denoted by B . We say a cube $Q = Q_1 \times \cdots \times Q_n \in \mathcal{Q}_{k+1}^{dn}$ is *strongly non-diagonal* if the n cubes Q_1, \dots, Q_n are distinct. Given a \mathcal{Q}_k discretized set $T \subset \mathbf{R}^d$, our goal is to construct a \mathcal{Q}_{k+1} discretized set $S \subset T$, such that $\mathcal{Q}_{k+1}^{dn}(F^n)$ does not contain any strongly non-diagonal cubes of $\mathcal{Q}_{k+1}^{dn}(B)$.

Lemma 28. *Fix $k, s \in [1, dn)$, and $\varepsilon \in [0, (dn - s)/2)$. Let $T \subset \mathbf{R}^d$ be a nonempty,*

\mathcal{Q}_k discretized set, and let $B \subset \mathbf{R}^{dn}$ be a nonempty \mathcal{Q}_{k+1} discretized set such that

$$\#(\mathcal{Q}_{k+1}(B)) \leq N_{k+1}^{s+\varepsilon}.$$

Then there exists a constant $C(s, d, n) > 0$, depending only on s , d , and n , such that, provided

$$N_{k+1} \geq C(s, d, n) \cdot M_{k+1}^{\frac{d(n-1)}{dn-s-\varepsilon}}, \quad (4.1)$$

then there is a \mathcal{Q}_{k+1} discretized set $S \subset T$ satisfying the following three properties:

(A) For any collection of n distinct cubes $Q_1, \dots, Q_n \in \mathcal{Q}_{k+1}(S)$,

$$Q_1 \times \dots \times Q_n \notin \mathcal{Q}_{k+1}(B).$$

(B) For each $Q \in \mathcal{Q}_k(T)$, there exists $\mathcal{R}_Q \subset \mathcal{R}_{k+1}(Q)$ such that

$$\#(\mathcal{R}_Q) \geq \#(\mathcal{R}_{k+1}(Q)),$$

and if $R \in \mathcal{R}_{k+1}(Q)$,

$$\#(\mathcal{Q}_{k+1}(R \cap S)) = \begin{cases} 1 & : R \in \mathcal{R}_Q \\ 0 & : R \notin \mathcal{R}_Q. \end{cases}$$

Proof. For each $R \in \mathcal{R}_{k+1}(T)$, pick Q_R uniformly at random from $\mathcal{Q}_{k+1}(R)$; these choices are independent as R ranges over $\mathcal{R}_{k+1}(T)$. Define

$$A = \bigcup \{Q_R : R \in \mathcal{R}_{k+1}(T)\},$$

and

$$\mathcal{K}(A) = \{K \in \mathcal{Q}_{k+1}(B) \cap \mathcal{Q}_{k+1}(A^n) : K \text{ strongly non-diagonal}\}.$$

The sets A and $\mathcal{K}(A)$ are random, in the sense that they depend on the random variables $\{Q_R\}$. Define

$$S(A) = \bigcup \left[\mathcal{Q}_{k+1}(A) - \{\pi(K) : K \in \mathcal{K}(A)\} \right], \quad (4.2)$$

where $\pi : \mathbf{R}^{dn} \rightarrow \mathbf{R}^d$ is the projection map $(x_1, \dots, x_n) \mapsto x_1$, for $x_i \in \mathbf{R}^d$.

Given any strongly non-diagonal cube $K = K_1 \times \dots \times K_n \in \mathcal{Q}_{k+1}(B)$, either

$K \notin \mathcal{Q}_{k+1}(A^n)$, or $K \in \mathcal{Q}_{k+1}(A^n)$. If the former occurs then $K \notin \mathcal{Q}_{k+1}(S(A))$ since $S(A) \subset A$, so $\mathcal{Q}_{k+1}(S(A)^n) \subset \mathcal{Q}_{k+1}(A^n)$. If the latter occurs then $K \in \mathcal{K}(A)$, and since $\pi(K) = K_1$, $K_1 \notin \mathcal{Q}_{k+1}(S(A))$. In either case, $K \notin \mathcal{Q}_{k+1}(S(A)^n)$, so $S(A)$ satisfies Property (A). By construction, we know that for each $R \in \mathcal{R}_{k+1}(T)$,

$$\#(\mathcal{Q}_{k+1}(S(A) \cap R)) \leq \#(\mathcal{Q}_{k+1}(A \cap R)) = 1. \quad (4.3)$$

To conclude that $S(A)$ satisfies Property (B), it therefore suffices to show that

$$\#(\mathcal{K}(A)) \leq (1/2) \cdot M_{k+1}^d.$$

We now show this is true with non-zero probability.

For each cube $Q \in \mathcal{Q}_{k+1}(T)$, there is a unique ‘parent’ cube $R \in \mathcal{R}_{k+1}(T)$ such that $Q \subset R$. Note that (4.1) implies, if $C(s, d, n) \geq 4d$, that $N_{k+1} \geq 4d \cdot M_{k+1}$. Since Q_R is chosen uniformly from $\mathcal{Q}_{k+1}(R)$, for any $Q \in \mathcal{Q}_{k+1}(R)$,

$$\mathbf{P}(Q \subset A) = \mathbf{P}(Q_R = Q) = (\mathcal{Q}_{k+1}(R))^{-1} = (M_{k+1}/N_{k+1})^d.$$

Suppose $K_1, \dots, K_n \in \mathcal{Q}_{k+1}(T)$ are distinct cubes. If the cubes have distinct parents in \mathcal{R}_{k+1}^d , we can apply the independence of the random cubes $\{Q_R\}$ to conclude that

$$\mathbf{P}(K_1, \dots, K_n \subset A) = (M_{k+1}/N_{k+1})^{dn}.$$

If the cubes K_1, \dots, K_n do not have distinct parents, (4.3) shows

$$\mathbf{P}(K_1, \dots, K_n \subset A) = 0.$$

In either case, we conclude that

$$\mathbf{P}(K_1, \dots, K_n \subset A) \leq (M_{k+1}/N_{k+1})^{dn}. \quad (4.4)$$

Let $K = K_1 \times \dots \times K_n$ be a strongly non-diagonal cube in $\mathcal{Q}_{k+1}(B)$. We deduce from (4.4) that

$$\mathbf{P}(K \subset A^n) = \mathbf{P}(K_1, \dots, K_n \subset A) \leq (M_{k+1}/N_{k+1})^{dn}. \quad (4.5)$$

If

$$C(s, d, n) \geq 4^{\frac{1}{dn-s}} \geq 2^{\frac{1}{dn-s-\varepsilon}},$$

by (4.5), linearity of expectation, and (4.1), we conclude

$$\begin{aligned}
\mathbf{E}(\#(\mathcal{K}(A))) &= \sum_{K \in \mathcal{Q}_{k+1}(B)} \mathbf{P}(K \subset A^n) \\
&\leq \#(\mathcal{Q}_{k+1}(B)) \cdot (M_{k+1}/N_{k+1})^{dn} \\
&\leq M_{k+1}^{dn} / N_{k+1}^{dn-s-\varepsilon} \\
&\leq (1/2) \cdot M_{k+1}^d.
\end{aligned}$$

In particular, there exists at least one (non-random) set A_0 such that

$$\#(\mathcal{K}(A_0)) \leq \mathbf{E}(\#(\mathcal{K}(A))) \leq (1/2) \cdot M_{k+1}^d. \quad (4.6)$$

In other words, $S(A_0) \subset A_0$ is obtained by removing at most $(1/2) \cdot M_{k+1}^d$ cubes in \mathcal{B}_s^d from A_0 . For each $Q \in \mathcal{B}_l^d(T)$, we know that $\# \mathcal{Q}_{k+1}(Q \cap A_0) = M_{k+1}^d$. Combining this with (4.6), we arrive at the estimate

$$\begin{aligned}
\# \mathcal{Q}_{k+1}(Q \cap S(A_0)) &= \mathcal{Q}_{k+1}(Q \cap A_0) - \#\{\pi(K) : K \in \mathcal{K}(A_0), \pi(K) \in A_0\} \\
&\geq \mathcal{Q}_{k+1}(Q \cap A_0) - \#(\mathcal{K}(A_0)) \\
&\geq M_{k+1}^d - (1/2) \cdot M_{k+1}^d \geq (1/2) \cdot M_{k+1}^d
\end{aligned}$$

Thus, $S(A_0)$ satisfies Property (B). Setting $S = S(A_0)$ completes the proof. \square

Remarks.

1. While Lemma 28 uses probabilistic arguments, the proof of the lemma is still constructive. In particular, one can find a suitable S constructively by checking every possible choice of A (there are finitely many) to find one particular choice A_0 which satisfies (4.6), and then defining S by (4.2). Thus the set we obtain in Theorem 27 exists by purely constructive means.

2. As with the proofs in Chapter 3, the fact that we can choose

$$N_{k+1} \sim M_{k+1}^{(1-\varepsilon)/t}$$

where $t = (dn - s)/d(n - 1)$, implies we should be able to iteratively apply Lemma 28 to obtain a set with Hausdorff dimension $(dn - s)/(n - 1)$.

4.2 Fractal Discretization

In this section we construct the set X from Theorem 27 by applying Lemma 28 at many scales. Let us start by fixing a strong cover of \mathcal{C} , as well as the branching factors $\{N_k\}$, that we will work with in the sequel.

Lemma 29. *Let $\mathcal{C} \subset \mathcal{C}^n(\mathbf{R}^d)$ be a countable union of bounded sets with Minkowski dimension at most s , and let $\varepsilon_k \searrow 0$ with $\varepsilon_k < (dn - s)/2$ for all k . Then there exists a choice of branching factors $\{N_k : k \geq 1\}$, and a sequence of sets $\{B_k : k \geq 1\}$, with $B_k \subset \mathbf{R}^{dn}$ for all k , such that*

- (A) Strong Cover: *The interiors $\{B_k^\circ\}$ of the sets $\{B_k\}$ form a strong cover of \mathcal{C} .*
- (B) Discreteness: *For all $k \geq 1$, B_k is a \mathcal{Q}_k discretized subset of \mathbf{R}^{dn} .*
- (C) Sparsity: *For all $k \geq 1$, $\mathcal{Q}_k(B_k) \leq N_k^{s+\varepsilon_k}$.*
- (D) Rapid Decay: *For all $\varepsilon > 0$ and $k \geq 1$, N_k is a power of two such that*

$$N_1 \dots N_{k-1} \lesssim_\varepsilon N_k^\varepsilon,$$

and for all $k \geq 1$,

$$N_k \geq C(s, d, n).$$

Proof. We can write $\mathcal{C} = \bigcup_{i=1}^\infty Y_i$, with $\dim_{\mathbf{M}}(Y_i) \leq s$ for each i . Let $\{i_k\}$ be a sequence of integers that repeats each integer infinitely often. The branching factors $\{N_k\}$ and sets $\{B_k\}$ are defined inductively. Suppose that the lengths N_1, \dots, N_{k-1} have been chosen. Since $\dim_{\mathbf{M}}(Y_{i_k}) < s + \varepsilon_k$, there are arbitrarily small lengths $l \leq l_{k-1}$ such that if $N_k = l_{k-1}/l$,

$$\#(\mathcal{Q}_k(Y_{i_k}(l))) \leq (1/l)^{s+(\varepsilon_k/2)}. \quad (4.7)$$

In particular, we can choose l small enough that $l \leq l_{k-1}^{2s/\varepsilon+2}$, which together with (4.7) implies

$$\#(\mathcal{Q}_k(Y_{i_k}(l))) \leq N_k^{s+\varepsilon_k}. \quad (4.8)$$

We can certainly also choose l small enough that

$$N_k \geq \max(C(s, d, n), (N_1 \dots N_{k-1})^{1/\varepsilon_k}).$$

We know that l_{k-1} is a power of two, so we can ensure N_k is a power of two. We then set $l_k = l$. With this choice, Property (D) is satisfied. And if $B_k =$

$\bigcup \mathcal{Q}_k(Y_{i_k}(l_k))$, then this choice of B_k clearly satisfies Property (B). And Property (C) is precisely Equation (4.8).

It remains to verify that the sets $\{B_k^\circ\}$ strongly cover \mathcal{C} . Fix a point $z \in \mathcal{C}$. Then there exists an index i such that $z \in Y_i$, and there is a subsequence k_1, k_2, \dots such that $i_{k_j} = i$ for each j . But then $z \in Y_i \subset B_{i_{k_j}}^\circ$, so z is contained in each of the sets $B_{i_{k_j}}^\circ$, and thus $z \in \limsup B_i^\circ$. Thus Property (A) is proved. \square

To construct X , we consider a nested, decreasing family of sets $\{X_k\}$, where each X_k is an l_k discretized subset of \mathbf{R}^d . We then set $X = \bigcap X_k$. The goal is to choose X_k such that X_k^n does not contain any *strongly non diagonal* cubes in B_k .

Lemma 30. *For each k , let $B_k \subset \mathbf{R}^{dn}$ be a \mathcal{Q}_k discretized set, such that the interiors $\{B_k^\circ\}$ strongly cover \mathcal{C} . For each index k , let X_k be a \mathcal{Q}_k discretized set such that $\mathcal{Q}_k(X_k^n) \cap \mathcal{Q}_k(B_k)$ contains no strongly non diagonal cubes. If $X = \bigcap X_k$, then X avoids B .*

Proof. Let $x = (x_1, \dots, x_n) \in \mathcal{C}$ be a point such that x_1, \dots, x_n are distinct. Define

$$\Delta = \{(y_1, \dots, y_n) \in \mathbf{R}^{dn} : \text{there exists } i \neq j \text{ such that } y_i = y_j\}.$$

Then $d(\Delta, x) > 0$, where d is the Hausdorff distance between Δ and x . Since $\{B_k\}$ strongly covers \mathcal{C} , there is a subsequence $\{k_m\}$ such that $x \in B_{k_m}^\circ$ for every index m . Since l_k converges to 0 and thus l_{k_m} converges to 0, if m is sufficiently large then $\sqrt{dn} \cdot l_{k_m} < d(\Delta, x)$. Note that $\sqrt{dn} \cdot l_{k_m}$ is the diameter of a cube in \mathcal{Q}_{k_m} . For such a choice of m , any cube $Q \in \mathcal{Q}_{k_m}^d$ which contains x is strongly non-diagonal. Thus $x \in B_{k_m}^\circ$. Since X_{k_m} and B_{k_m} share no cube which contains x , this implies $x \notin X_{k_m}$. In particular, this means $x \notin X^n$. \square

All that remains is to apply the discrete lemma to choose the sequence $\{X_k\}$. Given \mathcal{C} , apply Lemma 29 to choose a sequence $\{N_k\}$ and a sequence $\{B_k\}$. We recursively define the sequence $\{X_k\}$. Set $X_0 = [0, 1]^d$. Then, Property (D) of Lemma 29 implies

$$\#(\mathcal{Q}_{k+1}(B_{k+1})) \leq N_{k+1}^{s+\varepsilon_k}$$

Let M_{k+1} be the largest power of two smaller than

$$\left(\frac{N_{k+1}}{C(s, d, n)} \right)^{\frac{dn-s-\varepsilon_k}{d(n-1)}} \quad (4.9)$$

Then $1 \leq M_{k+1} \leq N_{k+1}$, and (4.1) is satisfied. Set $B = B_{k+1}$, and $T = X_k$. Then we can apply Lemma 28 to find a set S satisfying all Properties of that Lemma, and set $X_{k+1} = S$.

Property (A) of Lemma 28 together with Lemma 30 implies that the set $X = \bigcap X_k$ avoids \mathcal{C} . Property (B) of Lemma 28 shows that the sequence $\{X_k\}$ satisfies Property (A) of Theorem 15. Property (D) of Lemma 29 implies Property (B) of Theorem 15 is satisfied. And by definition, i.e. the choice of $\{M_k\}$ as given by (4.9),

$$N_k \leq M_k^{\frac{d(n-1)}{dn-s-\varepsilon_k}} \lesssim_\varepsilon M_k^{\frac{1+\varepsilon}{t}} \quad (4.10)$$

where $t = (nd - s)/d(n - 1)$. (4.10) is a form of Property (C) of Theorem 15. Thus all assumptions of Theorem 15 are satisfied, and so we find X has Hausdorff dimension $(nd - s)/(n - 1)$, completing the proof of Theorem 27.

Chapter 5

Applications

As discussed in the introduction, Theorem 27 generalizes Theorems 1.1 and 1.2 from [2]. In this chapter, we present two applications of Theorem 27 in settings where previous methods do not yield any results.

5.1 Sum-sets avoiding specified sets

Theorem 31. *Let $Y \subset \mathbf{R}^d$ be a countable union of sets with lower Minkowski dimension at most t . Then there exists a set $X \subset \mathbf{R}^d$ with Hausdorff dimension at least $d - t$ such that $X + X$ is disjoint from Y .*

Proof. Define $\mathcal{C} = \mathcal{C}_1 \cup \mathcal{C}_2$, where

$$\mathcal{C}_1 = \{(x, y) : x + y \in Y\} \quad \text{and} \quad \mathcal{C}_2 = \{(x, y) : y \in Y/2\}.$$

Since Y is a countable union of sets with lower Minkowski dimension at most t , \mathcal{C} is a countable union of sets with lower Minkowski dimension at most $d + t$. Applying Theorem 27 with $n = 2$ and $s = d + t$ produces a set $X \subset \mathbf{R}^d$ with Hausdorff dimension $2d - (d + t) = d - t$ such that $(x, y) \notin \mathcal{C}$ for all $x, y \in X$ with $x \neq y$. We claim that $X + X$ is disjoint from Y . To see this, first suppose $x, y \in X$, $x \neq y$. Since X avoids Z_1 , we conclude that $x + y \notin Y$. Suppose now that $x = y \in X$. Since X avoids Z_2 , we deduce that $X \cap (Y/2) = \emptyset$, and thus for any $x \in X$, $x + x = 2x \notin Y$. This completes the proof. \square

5.2 Subsets of Lipschitz curves avoiding isosceles triangles

In [2], Fraser and the second author prove that there exists a set $S \subset [0, 1]$ with dimension $\log_3 2$ such that for any simple C^2 curve $\gamma: [0, 1] \rightarrow \mathbf{R}^n$ with bounded non-vanishing curvature, $\gamma(S)$ does not contain the vertices of an isosceles triangle. Our method enables us to obtain a result that works for Lipschitz curves with small Lipschitz constants. The dimensional bound that we provide is slightly worse than [2] ($1/2$ instead of $\log_3 2$), and the set we obtain only works for a single Lipschitz curve, not for many curves simultaneously.

Theorem 32. *Let $f: [0, 1] \rightarrow \mathbf{R}^{n-1}$ be Lipschitz with*

$$\|f\|_{Lip} := \sup\{|f(x) - f(y)|/|x - y| : x, y \in [0, 1], x \neq y\} < 1.$$

Then there is a set $X \subset [0, 1]$ of Hausdorff dimension $1/2$ so that the set

$$\{(t, f(t)) : t \in X\}$$

does not contain the vertices of an isosceles triangle.

Corollary 33. *Let $f: [0, 1] \rightarrow \mathbf{R}^{n-1}$ be C^1 . Then there is a set $X \subset [0, 1]$ of Hausdorff dimension $1/2$ so that the set*

$$\{(t, f(t)) : t \in X\}$$

does not contain the vertices of an isosceles triangle.

Proof of Corollary 33. The graph of any C^1 function can be locally expressed, after possibly a translation and rotation, as the graph of a Lipschitz function with small Lipschitz constant. In particular, there exists an interval $I \subset [0, 1]$ of positive length so that the graph of f restricted to I , after being suitably translated and rotated, is the graph of a Lipschitz function $g: [0, 1] \rightarrow \mathbf{R}^{n-1}$ with Lipschitz constant at most $1/2$. Since isosceles triangles remain invariant under these transformations, the corollary is a consequence of Theorem 32. \square

Proof of Theorem 32. Set

$$\mathcal{C} = \left\{ (x_1, x_2, x_3) \in \mathcal{C}^3[0, 1] : \begin{array}{l} \text{The points } p_j = (x_j, f(x_j)) \text{ form} \\ \text{the vertices of an isosceles triangle} \end{array} \right\}. \quad (5.1)$$

In the next lemma, we show \mathcal{C} has lower Minkowski dimension at most two. By Theorem 27, there is a set $X \subset [0, 1]$ of Hausdorff dimension $1/2$ so that X avoids \mathcal{C} . This is precisely the statement that for each $x_1, x_2, x_3 \in X$, the points $(x_1, f(x_1))$, $(x_2, f(x_2))$, and $(x_3, f(x_3))$ do not form the vertices of an isosceles triangle. \square

Lemma 34. *Let $f: [0, 1] \rightarrow \mathbf{R}^{n-1}$ be Lipschitz with $\|f\|_{Lip} < 1$. Then the set \mathcal{C} given by (5.1) satisfies $\dim_{\overline{\mathbf{M}}}(\mathcal{C}) \leq 2$.*

Proof. First, notice that three points $p_1, p_2, p_3 \in \mathbf{R}^n$ form an isosceles triangle, with p_3 as the apex, if and only if $p_3 \in H_{p_1, p_2}$, where

$$H_{p_1, p_2} = \left\{ x \in \mathbf{R}^n : \left(x - \frac{p_1 + p_2}{2} \right) \cdot (p_2 - p_1) = 0 \right\}. \quad (5.2)$$

To prove \mathcal{C} has Minkowski dimension at most two, it suffices to show the set

$$W = \{x \in [0, 1]^3 : p_3 = (x_3, f(x_3)) \in H_{p_1, p_2}\}$$

has upper Minkowski dimension at most 2. This is because \mathcal{C} is covered by three copies of W , obtained by permuting coordinates. We work with the family of dyadic cubes \mathcal{D}^n . To bound the upper Minkowski dimension of W , we prove the estimate

$$\#(\mathcal{D}_k(W(1/2^k))) \leq Ck4^k \quad \text{for all } k \geq 1, \quad (5.3)$$

where C is a constant independent of k . Then for any $\varepsilon > 0$, (5.3) implies that for suitably large k ,

$$\#(\mathcal{D}_k(W(1/2^k))) \leq 2^{(2+\varepsilon)k}.$$

Since ε was arbitrary, this shows $\dim_{\overline{\mathbf{M}}}(W) \leq 2$.

To establish (5.3), we write

$$\#(\mathcal{D}_k(W(1/2^k))) = \sum_{m=0}^{2^k} \sum_{\substack{I_1, I_2 \in \mathcal{D}_k[0,1] \\ d(I_1, I_2) = m/2^k}} \#(\mathcal{D}_k(W(1/2^k) \cap (I_1 \times I_2 \times [0, 1])). \quad (5.4)$$

Our next task is to bound each of the summands in (5.4). Let $I_1, I_2 \in \mathcal{D}_k[0, 1]$, and let $m = 2^k \cdot d(I_1, I_2)$. Let x_1 be the midpoint of I_1 , and x_2 the midpoint of I_2 . Let

$(y_1, y_2, y_3) \in W \cap (I_1 \times I_2 \times [0, 1])$. Then it follows from (5.2) that

$$\left(y_3 - \frac{y_1 + y_2}{2}\right) \cdot (y_2 - y_1) + \left(f(y_3) - \frac{f(y_2) + f(y_1)}{2}\right) \cdot (f(y_2) - f(y_1)) = 0.$$

We know $|x_1 - y_1|, |x_2 - y_2| \leq 1/2^{k+1}$, so

$$\begin{aligned} & \left| \left(y_3 - \frac{y_1 + y_2}{2}\right) (y_2 - y_1) - \left(y_3 - \frac{x_1 + x_2}{2}\right) (x_2 - x_1) \right| \\ & \leq \frac{|y_1 - x_1| + |y_2 - x_2|}{2} |y_2 - y_1| + \left(|y_1 - x_1| + |y_2 - x_2|\right) \left|y_3 - \frac{x_1 + x_2}{2}\right| \quad (5.5) \\ & \leq (1/2^{k+1}) \cdot 1 + (1/2^k) \cdot 1 \leq 3/2^{k+1}. \end{aligned}$$

Conversely, $|f(x_1) - f(y_1)|, |f(x_2) - f(y_2)| \leq 1/2^{k+1}$ because $\|f\|_{\text{Lip}} \leq 1$, and a similar calculation yields

$$\begin{aligned} & \left| \left(f(y_3) - \frac{f(y_1) + f(y_2)}{2}\right) \cdot (f(y_2) - f(y_1)) \right. \\ & \quad \left. - \left(f(y_3) - \frac{f(x_1) + f(x_2)}{2}\right) \cdot (f(x_2) - f(x_1)) \right| \leq 3/2^{k+1}. \end{aligned} \quad (5.6)$$

Putting (5.5) and (5.6) together, we conclude that

$$\begin{aligned} & \left| \left(y_3 - \frac{x_1 + x_2}{2}\right) (x_2 - x_1) \right. \\ & \quad \left. + \left(f(y_3) - \frac{f(x_2) + f(x_1)}{2}\right) \cdot (f(x_2) - f(x_1)) \right| \leq 3/2^k. \end{aligned} \quad (5.7)$$

Since $|(x_2 - x_1, f(x_2) - f(x_1))| \geq |x_2 - x_1| \geq m/2^k$, we can interpret (5.7) as saying the point $(y_3, f(y_3))$ is contained in a $3/k$ thickening of the hyperplane $H_{(x_1, f(x_1)), (x_2, f(x_2))}$. Given another value $y' \in W \cap (I_1 \cap I_2 \cap [0, 1])$, it satisfies a variant of the inequality (5.7), and we can subtract the difference between the two inequalities to conclude

$$\left| (y_3 - y'_3) (x_2 - x_1) + (f(y_3) - f(y'_3)) \cdot (f(x_2) - f(x_1)) \right| \leq 6/2^k. \quad (5.8)$$

The triangle difference inequality applied with (5.8) implies

$$\begin{aligned} (f(y_3) - f(y'_3)) \cdot (f(x_2) - f(x_1)) &\geq |y_3 - y'_3| |x_2 - x_1| - 6/2^k \\ &= \frac{(m+1) \cdot |y_3 - y'_3| - 6}{2^k}. \end{aligned} \quad (5.9)$$

Conversely,

$$\begin{aligned} (f(y_3) - f(y'_3)) \cdot (f(x_2) - f(x_1)) &\leq \|f\|_{\text{Lip}}^2 \cdot |y_3 - y'_3| |x_2 - x_1| \\ &= \|f\|_{\text{Lip}}^2 \cdot (m+1)/2^k \cdot |y_3 - y'_3|. \end{aligned} \quad (5.10)$$

Combining (5.9) and (5.10) and rearranging, we see that

$$|y_3 - y'_3| \leq \frac{6}{(m+1)(1 - \|f\|_{\text{Lip}}^2)} \lesssim \frac{1}{m+1}, \quad (5.11)$$

where the implicit constant depends only on $\|f\|_{\text{Lip}}$. We conclude that

$$\# \mathcal{Q}_k(W(1/2^k) \cap (I_1 \times I_2 \times [0, 1])) \lesssim \frac{2^k}{m+1}, \quad (5.12)$$

which holds uniformly over any value of m .

We are now ready to bound the sum from (5.4). Note that for each value of m , there are at most 2^{k+1} pairs (I_1, I_2) with $d(I_1, I_2) = m/2^k$. Indeed, there are 2^k choices for I_1 and then at most two choices for I_2 . Equation (5.12) shows

$$\begin{aligned} \# \mathcal{Q}_k(W(1/2^k)) &= \sum_{m=0}^{2^k} \sum_{\substack{I_1, I_2 \in \mathcal{Q}_k[0,1] \\ d(I_1, I_2) = m/2^k}} \# \mathcal{Q}_k(W(1/2^k) \cap (I_1 \times I_2 \times [0, 1])) \\ &\lesssim 4^k \sum_{m=0}^{2^k} \frac{1}{m+1} \lesssim k/4^k. \end{aligned}$$

In the above inequalities, the implicit constants depend on $\|f\|_{\text{Lip}}$, but they are independent of k . This establishes (5.3) and completes the proof. \square

Chapter 6

Future Work

To conclude this thesis, we describe a rough sketch of some ideas which we hope will lead to future developments from the ideas of ‘rough sets avoiding patterns’ we detailed in Chapter 4. Section 6.1 attempts to exploit additional geometric information about rough configurations to find sets with large Hausdorff dimension avoiding patterns, and section 6.2 studies the problem of finding a set avoiding a rough configuration which supports a measure with large Fourier decay.

6.1 Low Rank Avoidance

Another way we can attempt to generalize the results of Chapter 4 is by attempting to improve the result if the rough configuration \mathcal{C} is given additional geometric structure. In Chapter 4, we showed a technique to find large sets avoid configurations with low Minkowski dimension. This means precisely that these configurations are efficiently covered by cubes at all scales. Here, we discuss whether one can obtain large sets avoiding configurations efficiently covered by other families of geometric shapes, e.g. by thickened lines, thickened planes, or more generally, by thickened variants of any family of algebraic hypersurfaces.

Theorem 35. *Let $\mathcal{C} \subset \mathcal{C}(\mathbf{R})$ be the countable union of sets $\{C_i\}$ such that*

- *For each i , there exists n_i such that C_i is a pre-compact subset of $\mathcal{C}^{n_i}(\mathbf{R})$.*
- *There exists an integer $k_i > 0$ and $s \in [0, k_i)$, together with a rational-coefficient linear transformation $M : \mathbf{R}^{n_i} \rightarrow \mathbf{R}^{k_i}$ such that $M(C_i)$ has Minkowski dimension s .*

Then there exists a set $X \subset [0, 1]$ avoiding \mathcal{C} with Hausdorff dimension at least

$$\inf_i \left(\frac{k_i - s_i}{k_i} \right).$$

Remark. Rescaling the problem by a constant does not affect any of the quantities in the problem. In particular, in our proof we may assume that the matrix we study is integer valued, which can be obtained by scaling down by a large integer.

Here, we only describe a discrete building block lemma, which can be fleshed out into a full proof by techniques analogous to that given in Chapters 3 and 4. But first, we must ‘change coordinates’ in the problem, so our description of the lemma is simpler. First, since our transformation M has full rank, we may find indices

$$i_1, \dots, i_k \in \{1, \dots, n\}$$

such that the transformation M is invertible when restricted to these indices the span of $\{e_{i_1}, \dots, e_{i_k}\}$. By an affine change of coordinates in the range of M , which preserves the Minkowski dimension of any set, we may assume that $M(e_{i_j}) = e_j$ for each $1 \leq j \leq k$.

Theorem 36. Fix $s \in [0, k)$ and $\varepsilon \in [0, (k-s)/2)$. Let $T_1, \dots, T_n \subset [0, 1]$ be disjoint, \mathcal{Q}_k discretized sets, and let $B \subset \mathbf{R}^k$ be a \mathcal{Q}_{k+1} discretized set such that

$$\#(\mathcal{Q}_{k+1}(B)) \leq N_{k+1}^{s+\varepsilon}.$$

Then there exists a constant $C(s, n, k, M) > 0$, and an integer constant $A(M) > 0$, such that if $A(M) \mid N_{k+1}$, and

$$N_{k+1} \geq C(s, n, k, M) \cdot M_{k+1}^{\frac{k}{s-k}}, \quad (6.1)$$

then there exists sets $S_1 \subset T_1, \dots, S_n \subset T_n$ such that

Proof. For each $i \notin \{i_1, \dots, i_k\}$, there are rational numbers $a_{ij} = p_{ij}/q_{ij} \in \mathbf{Q}$ such that $M(e_i) = \sum a_{ij} e_j$. For each interval $R \in \mathcal{R}_{k+1}(T_i)$, we let $a(R) \in \{0, \dots, N_1 \dots N_k M_{k+1} - 1\}$ be the integer such that

$$R = \left[\frac{a(R)}{N_1 \dots N_k M_{k+1}}, \frac{a(R) + 1}{N_1 \dots N_k M_{k+1}} \right].$$

Let $X \in \{0, \dots, N_{k+1}/M_{k+1} - 1\}^k$. For each $1 \leq j \leq k$, define

$$S_{i_j}(X) = \bigcup_{R \in \mathcal{R}_{k+1}(T_{i_j})} \left[\frac{a(R)}{N_1 \dots N_k M_{k+1}} + \frac{X_j}{N_1 \dots N_{k+1}}, \frac{a(R)}{N_1 \dots N_k M_{k+1}} + \frac{X_j + 1}{N_1 \dots N_{k+1}} \right].$$

For $i \notin \{i_1, \dots, i_k\}$, define

$$S_i(X) = \bigcup_{\substack{R \in \mathcal{R}_{k+1}(T_i) \\ \prod q_{ij} \mid a(R)}} \left[\frac{a(R)}{N_1 \dots N_k M_{k+1}}, \frac{a(R)}{N_1 \dots N_k M_{k+1}} + \frac{1}{N_1 \dots N_{k+1}} \right]$$

For each i , we let $S_i(X)$ denote the set of startpoints to intervals in S_i , and

$$\mathcal{A}(X) = M(S_1(X) \times \dots \times S_n(X)).$$

Then $\mathcal{A}(X) \subset (\mathbf{Z}^k + X)/(N_1 \dots N_{k+1})$. In particular, this implies that for $X \neq X'$, $d(\mathcal{A}(X), \mathcal{A}(X')) \geq 1/N_1 \dots N_{k+1}$. Applying the pigeonhole principle and applying (6.1), this means that there exists a constant C depending only on M and k , and some value of X_0 such that

$$\# \left\{ a \in \mathcal{A}(X_0) : d(a, B) \leq \frac{2}{\sqrt{d}\|M\|} \frac{1}{N_1 \dots N_{k+1}} \right\} \leq C \cdot \frac{N_{k+1}^{s+\varepsilon}}{(N_{k+1}/M_{k+1})^k} = C \cdot \frac{M_{k+1}^k}{N_{k+1}^{k-(s+\varepsilon)}} < 1.$$

In particular, this means that there is some X_0 such that

$$d(a, B) \geq \frac{2}{\sqrt{d}\|M\|} \frac{1}{N_1 \dots N_{k+1}}.$$

for all $a \in \mathcal{A}(X_0)$. But this means that $M(S_1(X_0) \times \dots \times S_n(X_0))$ is disjoint from B . Taking $S_i = S_i(X_0)$ for each i completes the proof. \square

Much remains on research related to this result. For instance, it is unclear whether one can improve the bound in Theorem 36 to

$$\inf \frac{k_i - s}{k_i - 1},$$

which would match the bound in Theorem 27, and how this theorem generalizes to sets avoiding patterns in \mathbf{R}^d , instead of just in \mathbf{R}^1 .

6.2 Fourier Dimension

A popular technique in current research in geometric measure theory is exploiting Fourier analysis to obtain additional structural information about configurations in sets. A key insight to this technique is that the Frostman dimension of any finite Borel measure μ is equal to

$$\sup \left\{ s > 0 : \int \frac{|\widehat{\mu}(\xi)|^2}{|\xi|^{d-s}} d\xi < \infty \right\}.$$

For brevity, we leave the proof to other sources, e.g. [8, Section 3.5]. Note that if

$$\int \frac{|\widehat{\mu}(\xi)|^2}{|\xi|^{d-s}} d\xi < \infty,$$

then there exists a constant C such that for *most* values $\xi \in \mathbf{R}^d$,

$$|\widehat{\mu}(\xi)| \leq C|\xi|^{-s/2}. \quad (6.2)$$

We obtain a strengthening of the Frostman measure condition if we require (6.2) to hold for *all* values ξ . In particular, we say a measure μ has *Fourier dimension* s if (6.2) holds for all $\xi \in \mathbf{R}^d$. If this is true, then for all $t < s$,

$$\int \frac{|\widehat{\mu}(\xi)|^2}{|\xi|^{d-t}} d\xi < \infty$$

so μ has Frostman dimension t for all $t < s$. Thus if we define the *Fourier dimension* of a set E as

$$\dim_{\mathbf{F}}(E) = \sup \left\{ s : \begin{array}{l} \text{there is a Borel probability measure } \mu \\ \text{supported on } E \text{ with Fourier dimension } s \end{array} \right\},$$

then $\dim_{\mathbf{F}}(E) \leq \dim_{\mathbf{H}}(E)$. We view the Fourier dimension as a refinement of the Hausdorff dimension which gives greater structural control on the set in the ‘frequency domain’.

There is much interest in determining whether sets with large Fourier dimension can avoid configurations. Results published recently include [4] and [9]. Most classical examples of fractals, like the middle-thirds Cantor set, have Fourier dimension zero. Nonetheless, it is a general principle that *random* families of sets tend to have Fourier dimension equal to the Hausdorff dimension. Since the main

technique of Theorem 27 involves a random selection strategy, we might hope that a modification of the procedure produces a set with large Fourier dimension.

Theorem 37. *Let $s \geq d$, and suppose \mathcal{C} is the countable union of pre-compact sets, each with lower Minkowski dimension at most s . Then there exists a set $X \subset [0, 1]^d$ with Fourier dimension at least $(nd - s)/n$ avoiding \mathcal{C} .*

Remark. Comparing Theorem 37 with Theorem 27, we see that our result here is slightly suboptimal, giving a dimension $(nd - s)/n$ set rather than the $(nd - s)/(n - 1)$ dimensional set we obtained in the Hausdorff dimension setting. The hope to obtain a more optimal result in the near future.

We begin with a lemma which simplifies the analysis of the Fourier decay of the probability measures we study to the analysis of frequencies in \mathbf{Z}^d .

Lemma 38. *Suppose μ is a compactly supported finite Borel measure, consider $s > 0$, and suppose there exists a constant C such that for each $m \in \mathbf{Z}^d$,*

$$|\widehat{\mu}(m)| \leq C|m|^{-s/2}. \quad (6.3)$$

Then there exists a constant C' such that for each $|\xi| \in \mathbf{R}^d$,

$$|\widehat{\mu}(\xi)| \lesssim C'|\xi|^{-s/2}. \quad (6.4)$$

In particular, μ has Fourier dimension s .

Proof. Without loss of generality, we may assume that μ is supported on a compact subset of $[1/3, 2/3]^d$, since every compactly supported measure is a finite sum of translates of measures of this form. Consider the distribution $\Lambda = \sum_{m \in \mathbf{Z}^d} \delta_m$, where δ_m is the Dirac delta distribution at m . Then the Poisson summation formula says that the Fourier transform of Λ is itself. If $\psi \in C_c(\mathbf{R}^d)$ is a bump function supported on $[0, 1]^d$, with $\psi(x) = 1$ for $x \in [1/3, 2/3]^d$, then $\mu = \psi(\Lambda * \mu)$, so

$$\begin{aligned} |\widehat{\mu}(\xi)| &= |[\widehat{\psi} * (\Lambda \widehat{\mu})](\xi)| \\ &= \left| \sum_{m \in \mathbf{Z}^d} \widehat{\mu}(m) (\widehat{\psi} * \delta_m)(\xi) \right| \\ &= \left| \sum_{m \in \mathbf{Z}^d} \widehat{\mu}(m) \widehat{\psi}(\xi - m) \right|. \end{aligned} \quad (6.5)$$

Since ψ is smooth, we know that for all $\eta \in \mathbf{R}^d$, $|\widehat{\psi}(\eta)| \lesssim 1/|\eta|^{d+1}$. If we perform a dyadic decomposition, we find

$$\begin{aligned}
\sum_{1 \leq |m-\xi| \leq |\xi|/2} |\widehat{\mu}(m)| |\widehat{\psi}(\xi-m)| &\lesssim \sum_{1 \leq |m-\xi| \leq |\xi|/2} |\xi|^{-s/2} |\widehat{\psi}(\xi-m)| \\
&\lesssim \sum_{k=1}^{\log |\xi|} \sum_{\frac{|\xi|}{2^{k+1}} \leq |m-\xi| \leq \frac{|\xi|}{2^k}} |\xi|^{-s/2} \left(2^k/|\xi|\right)^{d+1} \\
&\lesssim \sum_{k=1}^{\log |\xi|} |\xi|^{-s/2} (2^k/|\xi|) \lesssim |\xi|^{-s/2}.
\end{aligned} \tag{6.6}$$

There are $O(1)$ points $m \in \mathbf{Z}^d$ with $|m-\xi| \leq 1$, so if $|\xi| \geq 2$,

$$\sum_{|m-\xi| \leq 1} |\widehat{\mu}(m)| |\widehat{\psi}(m-\xi)| \lesssim |\xi|^{-s/2}. \tag{6.7}$$

We can also perform another dyadic decomposition, using the fact that for all $\eta \in \mathbf{R}^d$, $|\widehat{\psi}(\eta)| \lesssim 1/|\eta|^{2d}$, to find that

$$\begin{aligned}
\sum_{|m-\xi| \geq |\xi|/2} |\widehat{\mu}(m)| |\widehat{\psi}(m-\xi)| &\lesssim \sum_{k=0}^{\infty} \sum_{|\xi|2^{k-1} \leq |m-\xi| \leq |\xi|2^k} \frac{|\widehat{\mu}(m)|}{|\xi|^{2d} 2^{2dk}} \\
&\lesssim \sum_{k=0}^{\infty} |\xi|^{-d} 2^{-dk} \lesssim |\xi|^{-d}.
\end{aligned} \tag{6.8}$$

Combining (6.6), (6.7), and (6.8) with (6.5), we conclude that if $|\xi| \geq 2$,

$$|\widehat{\mu}(\xi)| \lesssim |\xi|^{-s/2}, \tag{6.9}$$

and since $\widehat{\mu}$ is bounded, (6.9) actually holds for all $\xi \in \mathbf{R}^d$. \square

Our goal now is to carefully modify the discrete selection strategy and discretized probability measures we use so that with high probability, we have sharp control over the Fourier transform of these measures. Lemma 38 implies that we only need control over integer-valued frequencies. The discretized measures $\{\nu_k\}$ we select are, for each k , a sum of point mass distributions at the points $(\mathbf{Z}/N_1 \dots N_k)^d$. Therefore, $\widehat{\nu}_k$ will be $N_1 \dots N_k$ periodic, in the sense that for any

$m \in (N_k \mathbf{Z})^d$ and $\xi \in \mathbf{R}^d$, $\widehat{v}_k(\xi + m) = \widehat{v}_k(\xi)$. Since we are only concerned with integer-valued frequencies, it will therefore suffice to obtain control of \widehat{v}_k on frequencies lying in $\{1, \dots, N_1 \dots N_k\}^d$.

In the discrete lemma below, we rely on a variant of the proof strategy of Theorem 2.1 of [9], but modified so that we can allow the branching factors $\{N_k\}$ to increase arbitrarily fast. We still use the families \mathcal{Q}_k^d and \mathcal{R}_k^d , together with parameters $\{N_k\}$ and $\{M_k\}$, in order to perform a multi-scale analysis. For each \mathcal{Q}_k discretized set $E \subset [0, 1]^d$, we define a probability measure

$$v_E = \frac{1}{\#(\mathcal{Q}_k(E))} \sum_{Q \in \mathcal{Q}_k(E)} \delta(a(Q)),$$

where for each $x \in \mathbf{R}^d$, $\delta(x)$ is the Dirac delta measure at x , and for each $Q \in \mathcal{Q}_k^d$, $a(Q)$ is the left-hand corner of the cube Q . Also, for each k , we define a probability measure

$$\eta_k = \frac{1}{N_{k+1}^d} \sum_{i_1, \dots, i_d=0}^{N_{k+1}} \delta\left(\frac{i}{N_1 \dots N_{k+1}}\right).$$

The purpose of introducing η_k is so that, given a measure μ which is a sum of point mass distributions in $(\mathbf{Z}/N_1 \dots N_k)^d$, the probability measure $\mu * \eta_k$ is a sum of point mass distributions in $(\mathbf{Z}/N_1 \dots N_{k+1})^d$, uniformly distributed at the scale $1/N_1 \dots N_{k+1}$.

Lemma 39. *Fix $s \in [1, dn)$, and $\varepsilon \in [0, (dn - s)/4)$. Let $T \subset \mathbf{R}^d$ be a nonempty, \mathcal{Q}_k discretized set, and let $B \subset \mathbf{R}^{dn}$ be a nonempty \mathcal{Q}_{k+1} discretized set such that*

$$\#(\mathcal{Q}_{k+1}(B)) \leq N_{k+1}^{s+\varepsilon}. \quad (6.10)$$

Then there exists a constant $A(d, n, s)$ such that, provided

$$M_{k+1}^{\frac{dn}{dn-s-2\varepsilon}} \leq N_{k+1} \leq 2M_{k+1}^{\frac{dn}{dn-s-2\varepsilon}}, \quad (6.11)$$

there exists a random \mathcal{Q}_k discretized set $S \subset T$, such that with probability

$$1 - \frac{1}{N_{k+1}^\varepsilon} - \left(\left\lceil \frac{A(d, n, s)}{\varepsilon} \right\rceil ! \right)^3 \frac{(N_1 \dots N_k)^d}{N_{k+1}^d},$$

the set S satisfies the following two properties:

(A) For any collection of n distinct cubes $Q_1, \dots, Q_n \in \mathcal{Q}_{k+1}(S)$,

$$Q_1 \times \dots \times Q_n \notin \mathcal{Q}_{k+1}(B).$$

(B) For any $m \in \mathbf{Z}^d$,

$$|\widehat{v}_S(m) - \widehat{\eta}_{k+1}(m)\widehat{v}_T(m)| \leq N_{k+1}^{-(1-\varepsilon)\frac{dn-s-2\varepsilon}{2n}}.$$

Proof. For each $R \in \mathcal{R}_{k+1}(T)$, let Q_R be randomly selected from $\mathcal{Q}_{k+1}(R)$, independently from all other selections $Q_{R'}$. Then, set $S = \bigcup \{Q_R : R \in \mathcal{R}_{k+1}(T)\}$. We then have

$$\#(\mathcal{Q}_{k+1}(S)) = \#(\mathcal{R}_{k+1}(T)) = M_{k+1}^d \mathcal{Q}_k(T).$$

Without loss of generality, removing cubes from B if necessary, we may assume that every cube $Q_1 \times \dots \times Q_n \in \mathcal{Q}_{k+1}(B)$, the values Q_1, \dots, Q_n are distinct. In particular, given any such cube, just as in Lemma 28, we have

$$\mathbf{P}(Q_1 \times \dots \times Q_n \in \mathcal{Q}_{k+1}(S^n)) = (M_{k+1}/N_{k+1})^{dn}.$$

Thus (6.10) and (6.11) imply

$$\mathbf{E}(\#(\mathcal{Q}_{k+1}(B) \cap \mathcal{Q}_{k+1}(S^n))) \leq M_{k+1}^{dn}/N_{k+1}^{dn-(s+\varepsilon)} \leq 1/N_{k+1}^\varepsilon.$$

Markov's inequality implies

$$\begin{aligned} \mathbf{P}(\mathcal{Q}_{k+1}(B) \cap \mathcal{Q}_{k+1}(S^n) \neq \emptyset) &= \mathbf{P}(\#(\mathcal{Q}_{k+1}(B) \cap \mathcal{Q}_{k+1}(S^n)) \geq 1) \\ &\leq 1/N_{k+1}^\varepsilon. \end{aligned} \tag{6.12}$$

Thus $\mathcal{Q}_{k+1}(S^n)$ is disjoint from $\mathcal{Q}_{k+1}(B)$ with high probability.

Now we analyze the Fourier transform of the measure v_S . For each cube $R \in \mathcal{R}_{k+1}(T)$, and for each m , let

$$A_R(m) = e^{\frac{-2\pi i m \cdot a(Q_R)}{N_1 \dots N_{k+1}}} - \frac{1}{N_{k+1}^d} \sum_{k_1, \dots, k_d=0}^N e^{\frac{-2\pi i m \cdot [N_{k+1} a(Q) + k]}{N_1 \dots N_{k+1}}}.$$

Then $\mathbf{E}[A_R(m)] = 0$, $|A_R(m)| \leq 2$ for each m , and

$$\widehat{v}_S(m) - \widehat{\eta}_{k+1}(m)\widehat{v}_T(m) = \frac{1}{\#(\mathcal{R}_{k+1}(T))} \sum_{R \in \mathcal{R}_{k+1}(T)} A_R(m).$$

Now fix a particular value of m . Since the random variables $A_R(m)$ are independent from one another as R ranges over $\mathcal{R}_{k+1}(T)$, we can apply Hoeffding's inequality to conclude that for each $t > 0$,

$$\mathbf{P}(|\widehat{v}_S(m) - \widehat{\eta}_{k+1}(m)\widehat{v}_T(m)| \geq t) \leq e^{-\#(\mathcal{R}_{k+1}(T))t^2/2} = e^{-\#(\mathcal{Q}_k(T))M_{k+1}^d t^2/2}.$$

In particular,

$$\mathbf{P}\left(|\widehat{v}_S(m) - \widehat{\eta}_{k+1}(m)\widehat{v}_T(m)| \geq M_{k+1}^{-d/2-\varepsilon}\right) \leq \exp(-\#(\mathcal{Q}_k(T))M_{k+1}^\varepsilon/2).$$

The function $\widehat{v}_S - \widehat{\eta}_{k+1}$ is $N_1 \dots N_{k+1}$ periodic. Thus, to uniformly bound $\widehat{v}_S(m) - \widehat{\eta}_{k+1}(m)$, we need only bound the function over $(N_1 \dots N_{k+1})^d$ values. Applying a union bound and applying (6.11), we therefore find that

$$\begin{aligned} \mathbf{P}\left(\|\widehat{v}_S - \widehat{\eta}_{k+1}\widehat{v}_T\|_{L^\infty(\mathbf{Z}^d)} \geq M_{k+1}^{-d/2-\varepsilon}\right) &\leq (N_1 \dots N_{k+1})^d \exp(-\#(\mathcal{Q}_k(T))M_{k+1}^\varepsilon/2) \\ &\leq 2^{\lceil \frac{4d^2n}{\varepsilon(dn-s)} \rceil} \left\lceil \frac{4d^2n}{\varepsilon(dn-s)} \right\rceil! \frac{(N_1 \dots N_{k+1})^d}{M_{k+1}^{4d^2n/(dn-s)}} \\ &\leq 2^{\lceil 8d^2n/(dn-s) \rceil} \left\lceil \frac{4d^2n}{\varepsilon(dn-s)} \right\rceil! \frac{(N_1 \dots N_k)^d}{N_{k+1}^d} \\ &\leq \left(\left\lceil \frac{4d^2n}{\varepsilon(dn-s)} \right\rceil! \right)^3 \frac{(N_1 \dots N_k)^d}{N_{k+1}^d}. \end{aligned} \tag{6.13}$$

Note that

$$M_{k+1}^{-d/2-\varepsilon} \lesssim N_{k+1}^{-(1-2\varepsilon/d)\frac{dn-s-2\varepsilon}{2n}} \leq N_{k+1}^{-(1-\varepsilon)\frac{dn-s-2\varepsilon}{2n}}.$$

If we set $A(d, n, s) = (4d^2n/(dn-s))$, then (6.12) and (6.13) allow us to conclude that with probability at least

$$1 - \frac{1}{N_{k+1}^\varepsilon} - \left[\left(\frac{A(d, n, s)}{\varepsilon} \right)! \right]^3 \frac{(N_1 \dots N_k)^d}{N_{k+1}^d},$$

the sets $\mathcal{Q}_{k+1}(S^n)$ and $\mathcal{Q}_{k+1}(B)$ are disjoint, and

$$\|\widehat{v}_S - \widehat{\eta}_{k+1}\widehat{v}_T\|_{L^\infty(\mathbf{Z}^d)} \leq N_{k+1}^{-(1-\varepsilon)\frac{dn-s-2\varepsilon}{2n}}. \quad \square$$

The remainder of the construction follows essentially the construction of the configuration avoiding set in Chapter 4, obtaining a family of discretized sets $\{X_k\}$ with $X = \bigcap X_k$. The only difference is we do not ‘terminate randomness’ at each application of the discrete lemma, instead just choosing intervals randomly at each stage and not worrying whether the sets we choose actually satisfy Properties (A) and (B) at each stage. Nonetheless, we know that these properties are satisfied with large probability, and so provided the parameters $\{N_k\}$ and ε_k satisfy

$$\sum_{k=1}^{\infty} \frac{1}{N_{k+1}^{\varepsilon}} + \left(\left\lceil \frac{A(d, n, s)}{\varepsilon_k} \right\rceil ! \right)^3 \frac{(N_1 \dots N_{k-1})^d}{N_k^d} < \infty,$$

the Borel-Cantelli lemma implies that almost surely, Properties (A) and (B) of Lemma 39 hold at all sufficiently large stages of the construction. This is sufficient to guarantee that the set X avoids the configuration \mathcal{C} . We will show that under the additional constraint that for each $\varepsilon > 0$,

$$(N_1 \dots N_k) \lesssim_{\varepsilon} N_{k+1}^{\varepsilon}, \quad (6.14)$$

the set X has Fourier dimension $(nd - s)/n$.

We must now construct a measure on X with the appropriate Fourier decay. The measures ν_S do not ‘quite’ suffice for this purpose, since their Fourier transforms do not actually have any decay. We will fix this by convolving ν_S with an appropriate bump function. Let ψ be a compactly supported, positive, smooth function supported on $[0, 1]^d$ with

$$\int \psi(x) dx = 1.$$

For each k , let

$$\psi_k(x) = (N_1 \dots N_k)^d \psi(N_1 \dots N_k x).$$

Then ψ_k is supported on $[0, 1/N_1 \dots N_k]^d$, $\int \psi_k(x) dx = 1$, and for any $\xi \in \mathbf{R}^d$,

$$\widehat{\psi_k}(\xi) = \widehat{\psi}(\xi/N_1 \dots N_k).$$

Note that, because ψ is smooth and compactly supported, for any $m > 0$,

$$|\widehat{\psi_k}(\xi)| \lesssim_m (N_1 \dots N_k)^m / |\xi|^m.$$

An important heuristic, a variant of the Heisenberg uncertainty principle, is that

$\widehat{\psi}_k$ is ‘essentially’ supported on the box $[-N_k, N_k]^d$.

Lemma 40. *Suppose that (6.14) holds. For any \mathcal{Q}_k discretized set E , the measure $\mu_E = \nu_E * \psi_k$ is supported on E , and for each $\varepsilon > 0$, there is a universal constant C_ε , independant of k and E , such that if $|\xi| \geq N_k^{1+\varepsilon}$, then*

$$|\widehat{\mu}_E(\xi)| \leq C_\varepsilon |\xi|^{-d/2}.$$

Proof. It is simple to see that μ_E is supported on E , since by general properties of convolution it is supported on the sum of the support of ψ_k , which is $[0, 1/N_k]^d$, and the support of ν_E , which is the union of the left-hand corners of the cubes in $\mathcal{Q}_k(E)$. We also note that

$$\widehat{\mu}_E(\xi) = \widehat{\nu}_E(\xi) \widehat{\psi}(\xi/N_k).$$

Since ν_E is a probability measure, $\|\widehat{\nu}_E\|_{L^\infty(\mathbf{R}^d)} = 1$. Now pick $m \geq d(1 + 1/\varepsilon)$. If $|\xi| \geq N_k^{1+\varepsilon}$, we calculate, using (6.14), to conclude

$$|\widehat{\mu}_E(\xi)| \leq |\widehat{\psi}(\xi/N_k)| \lesssim_m \frac{(N_1 \dots N_k)^m}{|\xi|^m} \lesssim_\varepsilon \frac{N_k^{m(1+\varepsilon/2)}}{|\xi|^{m-d/2}} \frac{1}{|\xi|^{d/2}} \leq \frac{1}{|\xi|^{d/2}}. \quad \square$$

Lemma 41. *Almost surely, for each $\varepsilon > 0$, there exists a constant $C_\varepsilon > 0$ and an integer $K > 0$, such that for any integer $k \geq K$, and any $m \in \mathbf{Z}^d$,*

$$|\widehat{\mu}_{X_k}(m)| \leq C_\varepsilon k \cdot |m|^{-\left(\frac{dn-s}{2n} - \varepsilon\right)}.$$

Proof. By the Borel-Cantelli lemma, almost surely, there exists some K such that Property (A) and (B) of Lemma 39 is satisfied at every stage of the construction past the K th stage. Without loss of generality, by choosing a larger K if necessary, that $\varepsilon_k \leq \varepsilon[2n/(dn - s + 2)]$. Because μ_{X_k} is smooth and compactly supported, there is certainly a constant C_ε such that for any $m \in \mathbf{Z}^d$,

$$|\widehat{\mu}_{X_k}(m)| \leq C_\varepsilon |m|^{-\left(\frac{dn-s}{2n} - \varepsilon\right)} \leq C_\varepsilon K \cdot |m|^{-\left(\frac{dn-s}{2n} - \varepsilon\right)}.$$

This constant should also be chosen larger than the constant in Lemma 40. The remainder of the argument is established by induction. Suppose that for each $m \in \mathbf{Z}^d$,

$$|\widehat{\mu}_{X_l}(m)| \leq C_\varepsilon l \cdot |m|^{-\left(\frac{dn-s}{2n} - \varepsilon\right)}.$$

For $|m| \leq N_{l+1}^{1+\varepsilon}$,

$$\begin{aligned} |\widehat{\mu_{X_{l+1}}}(m)| &\leq N_{l+1}^{-(1-\varepsilon_{l+1})\frac{dn-s-2\varepsilon_{l+1}}{2n}} + C_\varepsilon l \cdot |m|^{-\left(\frac{dn-s}{2n}-\varepsilon\right)} \\ &\leq (l+1)|m|^{-\left(\frac{dn-s}{2n}-\varepsilon\right)}. \end{aligned}$$

If $|m| \geq N_{k+1}^{1+\varepsilon}$, we can apply Lemma 40 to conclude that

$$|\widehat{\mu_{X_{l+1}}}(m)| \leq C_\varepsilon |m|^{-\left(\frac{dn-s}{2n}-\varepsilon\right)} \leq C_\varepsilon (l+1) |m|^{\left(\frac{dn-s}{2n}-\varepsilon\right)}.$$

This completes the proof. \square

The same argument as in Theorem 7 shows that the sequence $\{\mu_{X_k}\}$ is a Cauchy sequence under the weak $*$ topology. If we form the limit $\mu_X = \lim_{k \rightarrow \infty} \mu_{X_k}$, then the Fourier transforms $\widehat{\mu_{X_k}}$ converge pointwise to $\widehat{\mu_X}$. In particular, for each $\varepsilon > 0$ and $n \in \mathbf{Z}^d$, we conclude that

$$|\widehat{\mu_X}(n)| \leq \limsup |\widehat{\mu_{X_k}}(n)| \leq C_\varepsilon |n|^{-\left(\frac{nd-s}{n}-\varepsilon\right)}.$$

Combined with Lemma 38, this implies X almost surely has Fourier dimension $(nd-s)/n$.

Bibliography

- [1] K. Falconer. *Fractal Geometry: Mathematical Foundations and Applications*. Wiley, 2003. → pages 3, 11
- [2] R. Fraser and M. Pramanik. Large sets avoiding patterns. *Anal. PDE*, 11: 1083–1111, 2018. → pages 30, 40, 47, 48
- [3] T. Keleti. A 1-dimensional subset of the reals that intersects each of its translates in at most a single point. *Real Anal. Exchange*, 24:843–845, 1999. → pages 27, 40
- [4] I. Łaba and M. Pramanik. Arithmetic progressions in sets of fractional dimension. *Geom. Funct. Anal.*, 19(2):429–456, Sep 2009. → page 56
- [5] P. Maga. Full dimensional sets without given patterns. *Real Anal. Exchange*, 36:79–90, 2010. → page 1
- [6] A. Máthé. Sets of large dimension not containing polynomial configurations. *Adv. Math.*, 316:691–709, 2017. → page 35
- [7] P. Mattila. *Geometry of Sets & Measures in Euclidean Space*. 1999. → page 3
- [8] P. Mattila. *Fourier analysis and Hausdorff dimension*. Cambridge University Press, 2015. → pages 1, 56
- [9] P. Shmerkin. Salem sets with no arithmetic progressions. *Int. Math. Res. Not. IMRN*, 2017, 10 2015. → pages 56, 59
- [10] T. Tao. 245 C, Notes 5: Hausdorff dimension, 2009. URL <https://terrytao.wordpress.com/2009/05/19/245c-notes-5-hausdorff-dimension-optional/>. → page 3

