

Detangling a Twisted Form in L^4

Jacob Denson and Jake Fiedler

University of Wisconsin Madison

September 24, 2023

- Georgiev, Mukherjee, *Nodal Geometry, Heat Diffusion, and Brownian Motion*, Anal. PDE. **12** (2017), 133-148.
- Steinerberger, *Lower Bounds on Nodal Sets of Eigenfunctions via the Heat Flow*, Comm. Partial Differential Equations. **39** (2014), 2240-2261.
- Øksendal, *Stochastic Differential Equations*, Springer, 2003.
- Chung, *Green, Brown, and Probability and Brownian Motion*, World Scientific Publishing Company, 2002.

Nodal Domains

Goal

Study 'asymptotic geometry' of D_λ as $\lambda \rightarrow \infty$.

Main Result

- **Theorem:** There is $c_M > 0$ such that for any 'good' k -dimensional submanifold Σ of M , then

$$N(\Sigma, c_M/\lambda) = \{x \in M : d(x, \Sigma) < c_M/\lambda\}$$

doesn't contain D_λ .

- Consider the radius $1/\lambda$ tubular neighborhood

$$T_{1/\lambda}\Sigma = \bigcup_{x \in \Sigma} \{v \in (T_x \Sigma)^\perp : |v|_g \leq 1/\lambda\}.$$

The submanifold Σ is 'good' if the geodesic map $T_{1/\lambda}\Sigma \rightarrow N(\Sigma, 1/\lambda)$ is an embedding.

- Local condition: All principal curvatures of Σ are $\lesssim \lambda$.
- But no cheating globally!

Main Result

- **Theorem:** There is $c_M > 0$ such that for any ‘good’ k -dimensional submanifold Σ of M , then

$$N(\Sigma, c_M/\lambda) = \{x \in M : d(x, \Sigma) < c_M/\lambda\}$$

doesn't contain D_λ .

- Can replace Σ with a finite union of $\Omega(1/\lambda)$ separated ‘good’ submanifolds. Or allow finite unions with ‘transverse enough’ intersections.
- There is $C_M > 0$ such that $D_\lambda \subset N(Z_\lambda, C_M/\lambda)$.
- Heuristic: Elliptic methods work for $O(1/\lambda)$ localized results. We study stochastic diffusions, which provide cool tools to analyze eigenfunctions!

Uncertainty Principle on Manifolds?

- What would an analogous result look like on \mathbb{R}^d ?
- **Theorem:** Let D_λ be a nodal domain in \mathbb{R}^d . Then there is $c_d > 0$ such that if Σ is a finite union of $O(1/\lambda)$ -separated k dimensional planes, then D_λ is not contained in $N(\Sigma, c_d/\lambda)$.
- Stronger Result: D_λ contains a ball of radius $O(1/\lambda)$.
- Version on Manifolds: Paper proves for any $\varepsilon > 0$, there is $r_0 > 0$ such that if $x_0 \in D_\lambda$ maximizes $|e_\lambda(x_0)|$ in D_λ , then D_λ contains $1 - \varepsilon_0$ percent of $B(x_0, r_0\lambda^{-1/2})$.

Continuous Stochastic Processes

- Here are three ways to define continuous stochastic processes:
 - As a Borel-measurable function

$$X : \Omega \rightarrow C([0, \infty), M).$$

- As a family of correlated random variables

$$\{X_t : \Omega \rightarrow M : t \in [0, \infty)\}.$$

- As a law predicting future behaviour from present behaviour, i.e. by defining quantities such as

$$\mathbb{E}^x(f(X)) = \mathbb{E}[f(X)|X_0 = x]$$

$$\mathbb{P}^x(P(X)) = \mathbb{P}(P(X)|X_0 = x).$$

Brownian Motion on \mathbb{R}^d

- A stochastic process $\{B_t\}$ such that:
 - For any $I = [t, s]$, given $B_t = x$, the random variable $d_I B = B_s - B_t$ is normally distributed with mean x and variance $s - t$.
 - For any family of disjoint intervals $I_1, \dots, I_N \subset [0, \infty)$, with $I_k = [t_k, s_k]$, the random variables $d_{I_k} B$ are independent from one another.

Itô Diffusions

- Brownian Motion where diffusion is not radially symmetric.
- For each $x \in \mathbb{R}^d$, let $A(x)$ be a $d \times d$ positive semidefinite matrix. Then we have an Itô diffusion $\{X_t\}$ given in law by the 'Stochastic differential equation' $dX = A(X)dB$.
- For practical purposes, we have

$$X_{t+\delta} - X_t \approx A(X_t)[B_{t+\delta} - B_t]$$

where the difference between the LHS and RHS is a random variable with mean $o(\delta)$, and variance $O(\delta)$.

- Diffuses faster in directions where A has large eigenvalues.

Itô Diffusions

- Can define Itô diffusions on compact Riemannian manifolds M given a section $A : M \rightarrow \text{Hom}(TM)$ of positive definite matrices.
- We can define Brownian motion on a Riemannian manifold such that Brownian motion locally diffuses along geodesics at unit speed.

Connection to Elliptic Operators

- For any diffusion X , we can associate a semielliptic operator L , the *generator* of X , such that for $f \in C^\infty(M)$,

$$Lf(x) = \partial_t \{ \mathbb{E}^x[f(X_t)] \} |_{t=0} = \lim_{t \rightarrow 0^+} \frac{\mathbb{E}^x[f(X_t)] - f(x)}{t}.$$

- Second order because paths of X are 'half differentiable'.
- For Brownian motion (on \mathbb{R}^d or a manifold M), $L = \Delta/2$.
- 'Morally' apply the Fundamental Theorem of Calculus to get *Dynkin's Formula*

$$\mathbb{E}^x[f(X_T)] = f(x) + \mathbb{E}^x \left[\int_0^T (Lf)(X_s) ds \right].$$

Application: Escape Times

- In Dynkin's formula, T can be a 'stopping time', i.e. any $[0, \infty)$ valued function of X which doesn't 'predict the future', i.e. if T stops at a time t , it must only stop because of the properties of X on $[0, T]$, and not behaviour on (T, ∞) .
- Given an open, bounded set U , let

$$T_U = \inf\{t : X_t \notin U\}$$

be the *escape time* of U .

- If B is Brownian motion on \mathbb{R}^d , and U is the escape time of a ball of radius $R^{1/2}$ centered at x , $\mathbb{E}^x[T_U] = R/n$.
- If B is Brownian motion on M , escape time will be slower if volume expands (negative curvature) and faster if volume contracts (positive curvature). But irrelevant for the values R we care about.

Feynman Kac Formula

- Reverses Dynkin's Formula: Solves PDEs via Diffusions.
- Physically Intuitive Situations:

- (1) If $\partial_t u = Lu$ on M with $u_0 = f$, then

$$u(x, t) = \mathbb{E}^x[f(X_t)].$$

- (2) $\partial_t u = Lu$ on $D \subset M$ with $u_0 = f$ and $u = 0$ on ∂D ,

$$u(x, t) = \mathbb{E}^x[f(X_t)\chi_t],$$

where $\chi_t = \mathbb{I}(T_D > t)$ kills paths absorbed by ∂D .

- (3) If $Lu = 0$ on $D \subset M$ with $u = \phi$ on ∂D , then

$$u(x) = \mathbb{E}^x[\phi(X_{T_D})].$$

- Can also solve $\partial_t u = Lu$ with $\partial u / \partial \eta = 0$ on ∂D using 'reflection on Brownian motion', but a little more technical with singularities.

The Proof

- **Theorem:** There is $c_M > 0$ such that for any 'good' k -dimensional submanifold Σ of M , then

$$N(\Sigma, c_M/\lambda) = \{x \in M : d(x, \Sigma) < c_M/\lambda\}$$

doesn't contain D_λ .

- Assume $e_\lambda \geq 0$ on D_λ . Let $x^* = \operatorname{argmax}\{e_\lambda(x)\}$.
- Let $p(x, t)$ and $u(x, t)$ solve $\partial_t = \Delta$ with initial / boundary conditions:
 - $p_0 = 0$ and $p = 1$ on ∂D_λ .
 - $u_0 = e_\lambda$, and $u = 0$ on ∂D_λ .

- Given a non-negative matrix A_0 , alternatively apply row and column normalization, obtaining a sequence

$$A_0 \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- Given a non-negative matrix A_0 , alternatively apply row and column normalization, obtaining a sequence

$$A_0 \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- Claim: If $\text{Per}(A_0) > 0$, $d(A_i, \mathbf{S}) \rightarrow 0$, where \mathbf{S} is the family of doubly stochastic matrices.

- Given a non-negative matrix A_0 , alternatively apply row and column normalization, obtaining a sequence

$$A_0 \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- Claim: If $\text{Per}(A_0) > 0$, $d(A_i, \mathbf{S}) \rightarrow 0$, where \mathbf{S} is the family of doubly stochastic matrices.
- For even (odd) i , let γ_{ij} be the j th row (column) sum of A_i , so that $\text{Per}(A_{i+1}) = (\gamma_{i1} \dots \gamma_{in})^{-1} \cdot \text{Per}(A_i)$.

- Given a non-negative matrix A_0 , alternatively apply row and column normalization, obtaining a sequence

$$A_0 \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- Claim: If $\text{Per}(A_0) > 0$, $d(A_i, \mathbf{S}) \rightarrow 0$, where \mathbf{S} is the family of doubly stochastic matrices.
- For even (odd) i , let γ_{ij} be the j th row (column) sum of A_i , so that $\text{Per}(A_{i+1}) = (\gamma_{i1} \dots \gamma_{in})^{-1} \cdot \text{Per}(A_i)$.
- Two Key Facts Ensuring Convergence:**

- Given a non-negative matrix A_0 , alternatively apply row and column normalization, obtaining a sequence

$$A_0 \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- Claim: If $\text{Per}(A_0) > 0$, $d(A_i, \mathbf{S}) \rightarrow 0$, where \mathbf{S} is the family of doubly stochastic matrices.
- For even (odd) i , let γ_{ij} be the j th row (column) sum of A_i , so that $\text{Per}(A_{i+1}) = (\gamma_{i1} \dots \gamma_{in})^{-1} \cdot \text{Per}(A_i)$.
- Two Key Facts Ensuring Convergence:**
 - (1) $\text{Per}(A) \leq 1$ if A is partially normalized.

- Given a non-negative matrix A_0 , alternatively apply row and column normalization, obtaining a sequence

$$A_0 \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- Claim: If $\text{Per}(A_0) > 0$, $d(A_i, \mathbf{S}) \rightarrow 0$, where \mathbf{S} is the family of doubly stochastic matrices.
- For even (odd) i , let γ_{ij} be the j th row (column) sum of A_i , so that $\text{Per}(A_{i+1}) = (\gamma_{i1} \dots \gamma_{in})^{-1} \cdot \text{Per}(A_i)$.
- Two Key Facts Ensuring Convergence:**
 - $\text{Per}(A) \leq 1$ if A is partially normalized.
 - If $\Delta_i = \sum_j (\gamma_{ij} - 1)^2$, $\text{Per}(A_{i+1}) \geq (1 + C\Delta_i) \cdot \text{Per}(A_i)$.

- Given a non-negative matrix A_0 , alternatively apply row and column normalization, obtaining a sequence

$$A_0 \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- Claim: If $\text{Per}(A_0) > 0$, $d(A_i, \mathbf{S}) \rightarrow 0$, where \mathbf{S} is the family of doubly stochastic matrices.
- For even (odd) i , let γ_{ij} be the j th row (column) sum of A_i , so that $\text{Per}(A_{i+1}) = (\gamma_{i1} \dots \gamma_{in})^{-1} \cdot \text{Per}(A_i)$.
- Two Key Facts Ensuring Convergence:**
 - $\text{Per}(A) \leq 1$ if A is partially normalized.
 - If $\Delta_i = \sum_j (\gamma_{ij} - 1)^2$, $\text{Per}(A_{i+1}) \geq (1 + C\Delta_i) \cdot \text{Per}(A_i)$.
- Thus $\text{Per}(A_i)$ is bounded, monotonic, converges to $P \leq 1$.

- Given a non-negative matrix A_0 , alternatively apply row and column normalization, obtaining a sequence

$$A_0 \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$$

- Claim: If $\text{Per}(A_0) > 0$, $d(A_i, \mathbf{S}) \rightarrow 0$, where \mathbf{S} is the family of doubly stochastic matrices.
- For even (odd) i , let γ_{ij} be the j th row (column) sum of A_i , so that $\text{Per}(A_{i+1}) = (\gamma_{i1} \dots \gamma_{in})^{-1} \cdot \text{Per}(A_i)$.
- Two Key Facts Ensuring Convergence:**
 - $\text{Per}(A) \leq 1$ if A is partially normalized.
 - If $\Delta_i = \sum_j (\gamma_{ij} - 1)^2$, $\text{Per}(A_{i+1}) \geq (1 + C\Delta_i) \cdot \text{Per}(A_i)$.
- Thus $\text{Per}(A_i)$ is bounded, monotonic, converges to $P \leq 1$.
- If $\text{Per}(A_i) \geq P - \varepsilon$ for $\varepsilon \ll 1$, then

$$P \geq \text{Per}(A_{i+1}) \geq (1 + C \cdot \Delta_i) \cdot \text{Per}(A_i) \geq (1 + C \cdot \Delta_i)(P - \varepsilon).$$

Thus $\Delta_i \lesssim \varepsilon$. Taking $\varepsilon \rightarrow 0$ shows $\Delta_i \rightarrow 0$.

Sinkhorn Iteration

- **Proof that $\text{Per}(A_i) \leq 1$:**

Sinkhorn Iteration

- **Proof that $\text{Per}(A_i) \leq 1$:**
 - Simple inductive argument: the hypothesis is that if each row of a matrix B all sum up to less than one, then $\text{Per}(B) \leq 1$.

Sinkhorn Iteration

- **Proof that $\text{Per}(A_i) \leq 1$:**
 - Simple inductive argument: the hypothesis is that if each row of a matrix B all sum up to less than one, then $\text{Per}(B) \leq 1$.
- **Proof that $\text{Per}(A_{i+1}) \geq (1 + C\Delta_i) \cdot \text{Per}(A_i)$:**

Sinkhorn Iteration

- **Proof that $\text{Per}(A_i) \leq 1$:**
 - Simple inductive argument: the hypothesis is that if each row of a matrix B all sum up to less than one, then $\text{Per}(B) \leq 1$.
- **Proof that $\text{Per}(A_{i+1}) \geq (1 + C\Delta_i) \cdot \text{Per}(A_i)$:**
 - Really just more robust form of AGM inequality.

Sinkhorn Iteration

- **Proof that $\text{Per}(A_i) \leq 1$:**
 - Simple inductive argument: the hypothesis is that if each row of a matrix B all sum up to less than one, then $\text{Per}(B) \leq 1$.
- **Proof that $\text{Per}(A_{i+1}) \geq (1 + C\Delta_i) \cdot \text{Per}(A_i)$:**
 - Really just more robust form of AGM inequality.
 - If γ_{ij} are the row sums, then because the column sums are all one,

$$\frac{1}{n} \sum_j \gamma_{ij} = 1.$$

Sinkhorn Iteration

- **Proof that $\text{Per}(A_i) \leq 1$:**
 - Simple inductive argument: the hypothesis is that if each row of a matrix B all sum up to less than one, then $\text{Per}(B) \leq 1$.
- **Proof that $\text{Per}(A_{i+1}) \geq (1 + C\Delta_i) \cdot \text{Per}(A_i)$:**
 - Really just more robust form of AGM inequality.
 - If γ_{ij} are the row sums, then because the column sums are all one,

$$\frac{1}{n} \sum_j \gamma_{ij} = 1.$$

- AGM implies $\gamma_{i1} \dots \gamma_{in} \geq 1$, and monotonicity follows from

$$\text{Per}(A_{i+1}) = (\gamma_{i1} \dots \gamma_{in})^{-1} \text{Per}(A_i).$$

And now, back to our regularly scheduled programming

$$\text{BL}(B, p) = \sqrt{\sup_{A_1, \dots, A_m \succ 0} \frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i \cdot B_i^* A_i B_i)}}.$$

And now, back to our regularly scheduled programming

$$\text{BL}(B, p) = \sqrt{\sup_{A_1, \dots, A_m \succ 0} \frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i \cdot B_i^* A_i B_i)}}.$$

- Goal: Rescale our inputs so that

And now, back to our regularly scheduled programming

$$\text{BL}(B, p) = \sqrt{\sup_{A_1, \dots, A_m \succ 0} \frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i \cdot B_i^* A_i B_i)}}.$$

- Goal: Rescale our inputs so that
 - (Isotropy) $\sum p_i B_i^* B_i = I$.

And now, back to our regularly scheduled programming

$$\text{BL}(B, p) = \sqrt{\sup_{A_1, \dots, A_m \succ 0} \frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i \cdot B_i^* A_i B_i)}}.$$

- Goal: Rescale our inputs so that
 - (Isotropy) $\sum p_i B_i^* B_i = I$.
 - (Projection) $B_i B_i^* = I$ for each i .

Iteration of Operator Rescaling

- Sinkhorn: Alternately apply the following two procedures:

Iteration of Operator Rescaling

- Sinkhorn: Alternately apply the following two procedures:
 - (Isotropy Normalization)

Iteration of Operator Rescaling

- Sinkhorn: Alternately apply the following two procedures:
 - (Isotropy Normalization)
 - Let $M = \sum_i p_i B_i^* B_i$.

Iteration of Operator Rescaling

- Sinkhorn: Alternately apply the following two procedures:
 - (Isotropy Normalization)
 - Let $M = \sum_i p_i B_i^* B_i$.
 - Replace B_i with $B'_i = B_i M^{-1/2}$.

Iteration of Operator Rescaling

- Sinkhorn: Alternately apply the following two procedures:
 - (Isotropy Normalization)
 - Let $M = \sum_i p_i B_i^* B_i$.
 - Replace B_i with $B'_i = B_i M^{-1/2}$.
 - Then $\sum p_i (B'_i)^* B'_i = 1$, i.e. isotropy holds.

Iteration of Operator Rescaling

- Sinkhorn: Alternately apply the following two procedures:
 - (Isotropy Normalization)
 - Let $M = \sum_i p_i B_i^* B_i$.
 - Replace B_i with $B'_i = B_i M^{-1/2}$.
 - Then $\sum p_i (B'_i)^* B'_i = 1$, i.e. isotropy holds.
 - (Projection Normalization)

Iteration of Operator Rescaling

- Sinkhorn: Alternately apply the following two procedures:
 - (Isotropy Normalization)
 - Let $M = \sum_i p_i B_i^* B_i$.
 - Replace B_i with $B'_i = B_i M^{-1/2}$.
 - Then $\sum p_i (B'_i)^* B'_i = 1$, i.e. isotropy holds.
 - (Projection Normalization)
 - Let $M_i = B_i B_i^*$.

Iteration of Operator Rescaling

- Sinkhorn: Alternately apply the following two procedures:
 - (Isotropy Normalization)
 - Let $M = \sum_i p_i B_i^* B_i$.
 - Replace B_i with $B'_i = B_i M^{-1/2}$.
 - Then $\sum p_i (B'_i)^* B'_i = 1$, i.e. isotropy holds.
 - (Projection Normalization)
 - Let $M_i = B_i B_i^*$.
 - Replace B_i with $B'_i = M_i^{-1/2} B_i$.

Iteration of Operator Rescaling

- Sinkhorn: Alternately apply the following two procedures:
 - (Isotropy Normalization)
 - Let $M = \sum_i p_i B_i^* B_i$.
 - Replace B_i with $B'_i = B_i M^{-1/2}$.
 - Then $\sum p_i (B'_i)^* B'_i = 1$, i.e. isotropy holds.
 - (Projection Normalization)
 - Let $M_i = B_i B_i^*$.
 - Replace B_i with $B'_i = M_i^{-1/2} B_i$.
 - Then $(B'_i)^* B'_i = I$ for each i .

Iteration of Operator Rescaling

- Sinkhorn: Alternately apply the following two procedures:
 - (Isotropy Normalization)
 - Let $M = \sum_i p_i B_i^* B_i$.
 - Replace B_i with $B'_i = B_i M^{-1/2}$.
 - Then $\sum p_i (B'_i)^* B'_i = 1$, i.e. isotropy holds.
 - (Projection Normalization)
 - Let $M_i = B_i B_i^*$.
 - Replace B_i with $B'_i = M_i^{-1/2} B_i$.
 - Then $(B'_i)^* B'_i = I$ for each i .
- We obtain a sequence $B \rightarrow B_1 \rightarrow B_2 \rightarrow \dots$

Iteration of Operator Rescaling

- **Two Key Facts Ensuring Convergence:**

Iteration of Operator Rescaling

- **Two Key Facts Ensuring Convergence:**

(1) $BL(B, p) \geq 1$ if (B, p) is partially normalized.

Iteration of Operator Rescaling

- **Two Key Facts Ensuring Convergence:**

- (1) $\text{BL}(B, p) \geq 1$ if (B, p) is partially normalized.
- (2) For some $r \leq \text{Poly}(\text{Bits}(B), d, n)$, if

$$\text{BL}(B_i, p) \geq 1 + \varepsilon,$$

then

$$\text{BL}(B_{i+1}, p) \leq (1 - C\varepsilon^r)\text{BL}(B_i, p).$$

Iteration of Operator Rescaling

- **Two Key Facts Ensuring Convergence:**

- (1) $\text{BL}(B, p) \geq 1$ if (B, p) is partially normalized.
- (2) For some $r \leq \text{Poly}(\text{Bits}(B), d, n)$, if

$$\text{BL}(B_i, p) \geq 1 + \varepsilon,$$

then

$$\text{BL}(B_{i+1}, p) \leq (1 - C\varepsilon^r)\text{BL}(B_i, p).$$

- Thus convergence occurs as with Sinkhorn iteration provided that $\text{BL}(B, p) < \infty$.

Iteration of Operator Rescaling

- **Two Key Facts Ensuring Convergence:**

- (1) $\text{BL}(B, p) \geq 1$ if (B, p) is partially normalized.
- (2) For some $r \leq \text{Poly}(\text{Bits}(B), d, n)$, if

$$\text{BL}(B_i, p) \geq 1 + \varepsilon,$$

then

$$\text{BL}(B_{i+1}, p) \leq (1 - C\varepsilon^r)\text{BL}(B_i, p).$$

- Thus convergence occurs as with Sinkhorn iteration provided that $\text{BL}(B, p) < \infty$.
- (1) and (2) follow from techniques in the study of *positive operators*.

Positive Operators

- A linear map $T : M_n \rightarrow M_n$ is *completely positive* if there are $n \times n$ matrices B_1, \dots, B_K and $p_i > 0$ such that

$$T(A) = \sum p_i B_i A B_i^*.$$

Positive Operators

- A linear map $T : M_n \rightarrow M_n$ is *completely positive* if there are $n \times n$ matrices B_1, \dots, B_K and $p_i > 0$ such that

$$T(A) = \sum p_i B_i A B_i^*.$$

- $T : M_n \rightarrow M_n$ is *positive* if $A \succeq 0$ implies $T(A) \succeq 0$.

Positive Operators

- A linear map $T : M_n \rightarrow M_n$ is *completely positive* if there are $n \times n$ matrices B_1, \dots, B_K and $p_i > 0$ such that

$$T(A) = \sum p_i B_i A B_i^*.$$

- $T : M_n \rightarrow M_n$ is *positive* if $A \succeq 0$ implies $T(A) \succeq 0$.
- Can reduce the study of non-negative matrices to positive operators: For a non-negative matrix S , $T(A)$ is the diagonal matrix whose entries are precisely the vector Sa , where a is the vector formed from the diagonal entries of A .

Positive Operators

- A linear map $T : M_n \rightarrow M_n$ is *completely positive* if there are $n \times n$ matrices B_1, \dots, B_K and $p_i > 0$ such that

$$T(A) = \sum p_i B_i A B_i^*.$$

- $T : M_n \rightarrow M_n$ is *positive* if $A \succeq 0$ implies $T(A) \succeq 0$.
- Can reduce the study of non-negative matrices to positive operators: For a non-negative matrix S , $T(A)$ is the diagonal matrix whose entries are precisely the vector Sa , where a is the vector formed from the diagonal entries of A .
- Given T , we have $T^*(A) = \sum p_i B_i^* A B_i$.

Further Connections

- For simplicity, look at Brascamp Lieb where all spaces have the same dimension (all B_i are square matrices).

Further Connections

- For simplicity, look at Brascamp Lieb where all spaces have the same dimension (all B_i are square matrices).
- $\text{BL}(B, p) < \infty$ can only hold if $\sum p_i = 1$.

Further Connections

- For simplicity, look at Brascamp Lieb where all spaces have the same dimension (all B_i are square matrices).
- $\text{BL}(B, p) < \infty$ can only hold if $\sum p_i = 1$.
- Consider optimizing the quantity

$$\inf_{A \succ 0} \frac{\det(\sum p_i B_i^* A B_i)}{\det(A)}$$

analogous to

$$\text{BL}(B, p) = \sup_{A_1, \dots, A_m \succ 0} \sqrt{\frac{\prod_i \det(A_i)^{p_i}}{\det(\sum p_i \cdot B_i^* A_i B_i)}},$$

if all A_i are equal.

Capacity of Operators

$$\inf_{A \succ 0} \frac{\det(\sum p_i B_i^* A B_i)}{\det(A)}$$

Capacity of Operators

$$\inf_{A \succ 0} \frac{\det(\sum p_i B_i^* A B_i)}{\det(A)}$$

- (Gurvits, 2004) The *capacity* of $T : M_n \rightarrow M_n$ is

$$\text{Cap}(T) = \inf_{A \succ 0} \frac{\det(TA)}{\det(A)}.$$

Capacity of Operators

$$\inf_{A \succ 0} \frac{\det(\sum p_i B_i^* A B_i)}{\det(A)}$$

- (Gurvits, 2004) The *capacity* of $T : M_n \rightarrow M_n$ is

$$\text{Cap}(T) = \inf_{A \succ 0} \frac{\det(TA)}{\det(A)}.$$

- For any Brascamp-Lieb data (B, p) , there exists a positive $T : M_n \rightarrow M_n$ and k such that $\text{Cap}(T) = 1/\text{BL}(B, p)^{2k}$.

Capacity of Operators

$$\inf_{A \succ 0} \frac{\det(\sum p_i B_i^* A B_i)}{\det(A)}$$

- (Gurvits, 2004) The *capacity* of $T : M_n \rightarrow M_n$ is

$$\text{Cap}(T) = \inf_{A \succ 0} \frac{\det(TA)}{\det(A)}.$$

- For any Brascamp-Lieb data (B, p) , there exists a positive $T : M_n \rightarrow M_n$ and k such that $\text{Cap}(T) = 1/\text{BL}(B, p)^{2k}$.
- Positive operators are well studied in the quantum information theory literature, so reduction of BL to this theory is useful.

Doubly Stochastic Positive Operators

- (Isotropy) Let $T(A) = \sum p_i B_i^* A B_i$.

Doubly Stochastic Positive Operators

- (Isotropy) Let $T(A) = \sum p_i B_i^* A B_i$.
 - $\sum p_i B_i^* B_i = I$ holds iff $T(I) = I$.

Doubly Stochastic Positive Operators

- (Isotropy) Let $T(A) = \sum p_i B_i^* A B_i$.
 - $\sum p_i B_i^* B_i = I$ holds iff $T(I) = I$.
- (Projection) Let $T(A) = B_i^* A B_i$.

Doubly Stochastic Positive Operators

- (Isotropy) Let $T(A) = \sum p_i B_i^* A B_i$.
 - $\sum p_i B_i^* B_i = I$ holds iff $T(I) = I$.
- (Projection) Let $T(A) = B_i^* A B_i$.
 - $B_i B_i^* = I$ if and only if $T^*(I) = I$.

Doubly Stochastic Positive Operators

- (Isotropy) Let $T(A) = \sum p_i B_i^* A B_i$.
 - $\sum p_i B_i^* B_i = I$ holds iff $T(I) = I$.
- (Projection) Let $T(A) = B_i^* A B_i$.
 - $B_i B_i^* = I$ if and only if $T^*(I) = I$.
- If (B, p) is a Brascamp-Lieb datum with associated operator $T : M_n \rightarrow M_n$, then (B, p) is geometric if and only if T is *doubly stochastic*, i.e. $T(I) = I$ and $T^*(I) = I$.

Operator Rescaling for Positive Operators

- If T is doubly stochastic, $\text{Cap}(T) = 1$.

Operator Rescaling for Positive Operators

- If T is doubly stochastic, $\text{Cap}(T) = 1$.
- We can rescale. If

$$T_{M_1 M_2}(A) = M_2^* T(M_1^* A M_1) M_2,$$

then $\text{Cap}(T_{M_1, M_2}) = \det(M_1)^2 \det(M_2)^2 \cdot \text{Cap}(T)$.

Operator Rescaling for Positive Operators

- If T is doubly stochastic, $\text{Cap}(T) = 1$.
- We can rescale. If

$$T_{M_1 M_2}(A) = M_2^* T(M_1^* A M_1) M_2,$$

then $\text{Cap}(T_{M_1, M_2}) = \det(M_1)^2 \det(M_2)^2 \cdot \text{Cap}(T)$.

- Sinkhorn says to iterate

$$T \mapsto T_{I, T(I)^{-1/2}} \quad \text{and} \quad T \mapsto T_{T^*(I)^{-1/2}, I}.$$

Operator Rescaling for Positive Operators

- If T is doubly stochastic, $\text{Cap}(T) = 1$.
- We can rescale. If

$$T_{M_1 M_2}(A) = M_2^* T(M_1^* A M_1) M_2,$$

then $\text{Cap}(T_{M_1, M_2}) = \det(M_1)^2 \det(M_2)^2 \cdot \text{Cap}(T)$.

- Sinkhorn says to iterate

$$T \mapsto T_{I, T(I)^{-1/2}} \quad \text{and} \quad T \mapsto T_{T^*(I)^{-1/2}, I}.$$

- If $\text{Cap}(T) > 0$, iteration yields a rescaling arbitrarily close to a doubly stochastic operator, in $\text{Poly}(\text{Bits}(B), 1/\varepsilon)$ time.

Upper Bounds For Capacity

Upper Bounds For Capacity

- To guarantee efficiency, we need to show that for partially normalized T , $\text{Cap}(T) \geq 1/e^{\text{Poly}(\text{Bits}(B))}$.

Upper Bounds For Capacity

- To guarantee efficiency, we need to show that for partially normalized T , $\text{Cap}(T) \geq 1/e^{\text{Poly}(\text{Bits}(B))}$.
- (Gurvits, 2004) If $T(A) = \sum B_i A B_i^*$, and $\det(\sum B_i) \neq 0$, then $\text{Cap}(T) \gtrsim (\text{Bits}(B) \cdot n)^{-O(n)}$.

Upper Bounds For Capacity

- To guarantee efficiency, we need to show that for partially normalized T , $\text{Cap}(T) \geq 1/e^{\text{Poly}(\text{Bits}(B))}$.
- (Gurvits, 2004) If $T(A) = \sum B_i A B_i^*$, and $\det(\sum B_i) \neq 0$, then $\text{Cap}(T) \gtrsim (\text{Bits}(B) \cdot n)^{-O(n)}$.
- If $\text{Cap}(T) > 0$, there is $d > 0$ and $d \times d$ matrices C_i s.t.
$$\det(\sum C_i \otimes B_i) \neq 0 \quad \text{and} \quad \text{Bits}(C) \leq \text{Poly}(d, \text{Bits}(B)).$$

Upper Bounds For Capacity

- To guarantee efficiency, we need to show that for partially normalized T , $\text{Cap}(T) \geq 1/e^{\text{Poly}(\text{Bits}(B))}$.
- (Gurvits, 2004) If $T(A) = \sum B_i A B_i^*$, and $\det(\sum B_i) \neq 0$, then $\text{Cap}(T) \gtrsim (\text{Bits}(B) \cdot n)^{-O(n)}$.
- If $\text{Cap}(T) > 0$, there is $d > 0$ and $d \times d$ matrices C_i s.t.
$$\det(\sum C_i \otimes B_i) \neq 0 \quad \text{and} \quad \text{Bits}(C) \leq \text{Poly}(d, \text{Bits}(B)).$$
- 'Fairly simple' to show that if $S(A) = \sum C_i A C_i^*$, and $(S \otimes T)(A) = \sum (C_i \otimes B_i) A (C_i \otimes B_i)^*$, then
$$\text{Cap}(S \otimes T) \leq \text{Cap}(S)^n \text{Cap}(T)^d.$$

Upper Bounds For Capacity

- To guarantee efficiency, we need to show that for partially normalized T , $\text{Cap}(T) \geq 1/e^{\text{Poly}(\text{Bits}(B))}$.
- (Gurvits, 2004) If $T(A) = \sum B_i A B_i^*$, and $\det(\sum B_i) \neq 0$, then $\text{Cap}(T) \gtrsim (\text{Bits}(B) \cdot n)^{-O(n)}$.
- If $\text{Cap}(T) > 0$, there is $d > 0$ and $d \times d$ matrices C_i s.t.
$$\det(\sum C_i \otimes B_i) \neq 0 \quad \text{and} \quad \text{Bits}(C) \leq \text{Poly}(d, \text{Bits}(B)).$$
- 'Fairly simple' to show that if $S(A) = \sum C_i A C_i^*$, and $(S \otimes T)(A) = \sum (C_i \otimes B_i) A (C_i \otimes B_i)^*$, then
$$\text{Cap}(S \otimes T) \leq \text{Cap}(S)^n \text{Cap}(T)^d.$$
- Since $\text{Cap}(L) \leq 1$, it follows that
$$\text{Cap}(T) \geq \text{Cap}(S \otimes T)^{1/d} \gtrsim (\text{Poly}(d, \text{Bits}(B))n)^{-O(n)}.$$

Upper Bounds For Capacity

- To guarantee efficiency, we need to show that for partially normalized T , $\text{Cap}(T) \geq 1/e^{\text{Poly}(\text{Bits}(B))}$.
- (Gurvits, 2004) If $T(A) = \sum B_i A B_i^*$, and $\det(\sum B_i) \neq 0$, then $\text{Cap}(T) \gtrsim (\text{Bits}(B) \cdot n)^{-O(n)}$.
- If $\text{Cap}(T) > 0$, there is $d > 0$ and $d \times d$ matrices C_i s.t.
$$\det(\sum C_i \otimes B_i) \neq 0 \quad \text{and} \quad \text{Bits}(C) \leq \text{Poly}(d, \text{Bits}(B)).$$
- 'Fairly simple' to show that if $S(A) = \sum C_i A C_i^*$, and $(S \otimes T)(A) = \sum (C_i \otimes B_i) A (C_i \otimes B_i)^*$, then
$$\text{Cap}(S \otimes T) \leq \text{Cap}(S)^n \text{Cap}(T)^d.$$
- Since $\text{Cap}(L) \leq 1$, it follows that
$$\text{Cap}(T) \geq \text{Cap}(S \otimes T)^{1/d} \gtrsim (\text{Poly}(d, \text{Bits}(B))n)^{-O(n)}.$$
- Invariant theory shows we can choose $d \leq n^4[(n+1)!]^2$.

The Invariant Theory

The Invariant Theory

- We have a group action of $SL_n \times SL_n$ on tuples $B = (B_1, \dots, B_m)$, such that

$$(M, N) \circ B = (MB_1N, \dots, MB_mN).$$

The Invariant Theory

- We have a group action of $SL_n \times SL_n$ on tuples $B = (B_1, \dots, B_m)$, such that

$$(M, N) \circ B = (MB_1N, \dots, MB_mN).$$

- Invariant Theory: Find the ring R of all 'invariant polynomials' $f(B)$ such that

$$f((M, N) \circ B) = f(B)$$

for all $(M, N) \in SL_n \times SL_n$ and all tuples B .

The Invariant Theory

- We have a group action of $SL_n \times SL_n$ on tuples $B = (B_1, \dots, B_m)$, such that

$$(M, N) \circ B = (MB_1N, \dots, MB_mN).$$

- Invariant Theory: Find the ring R of all 'invariant polynomials' $f(B)$ such that

$$f((M, N) \circ B) = f(B)$$

for all $(M, N) \in SL_n \times SL_n$ and all tuples B .

- Note: $f_C(B) = \det(\sum C_i \otimes B_i)$ is an invariant homogeneous polynomial under this action for any C_i .

The Invariant Theory

- (Nayak and Subrahmanyam, 2010) R is a ring generated by the homogeneous polynomials $f_C(A) = \det(\sum C_i \circ A_i)$, for $d \times d$ matrices C_i , for all $d > 0$.

The Invariant Theory

- (Nayak and Subrahmanyam, 2010) R is a ring generated by the homogeneous polynomials $f_C(A) = \det(\sum C_i \circ A_i)$, for $d \times d$ matrices C_i , for all $d > 0$.
 - (1890) Hilbert showed that for a fairly general family of group actions, R is finitely generated, so we should expect there is some d_0 such that R is generated by the polynomials above for $d \leq d_0$. Killed the field for 100 years.

The Invariant Theory

- (Nayak and Subrahmanyam, 2010) R is a ring generated by the homogeneous polynomials $f_C(A) = \det(\sum C_i \circ A_i)$, for $d \times d$ matrices C_i , for all $d > 0$.
 - (1890) Hilbert showed that for a fairly general family of group actions, R is finitely generated, so we should expect there is some d_0 such that R is generated by the polynomials above for $d \leq d_0$. Killed the field for 100 years.
 - (Ivanyos, Qiao, Subrahmanyam, 2015)
 $d_0 \lesssim n^4[(n+1)!]^2$.

The Invariant Theory

- (Nayak and Subrahmanyam, 2010) R is a ring generated by the homogeneous polynomials $f_C(A) = \det(\sum C_i \circ A_i)$, for $d \times d$ matrices C_i , for all $d > 0$.
 - (1890) Hilbert showed that for a fairly general family of group actions, R is finitely generated, so we should expect there is some d_0 such that R is generated by the polynomials above for $d \leq d_0$. Killed the field for 100 years.
 - (Ivanyos, Qiao, Subrahmanyam, 2015)
 $d_0 \lesssim n^4[(n+1)!]^2$.
 - Suppose there are $d \times d$ matrices C_i (with d minimal) such that $f_C(B) \neq 0$.

The Invariant Theory


- (Nayak and Subrahmanyam, 2010) R is a ring generated by the homogeneous polynomials $f_C(A) = \det(\sum C_i \circ A_i)$, for $d \times d$ matrices C_i , for all $d > 0$.
 - (1890) Hilbert showed that for a fairly general family of group actions, R is finitely generated, so we should expect there is some d_0 such that R is generated by the polynomials above for $d \leq d_0$. Killed the field for 100 years.
 - (Ivanyos, Qiao, Subrahmanyam, 2015)
 $d_0 \lesssim n^4[(n+1)!]^2$.
 - Suppose there are $d \times d$ matrices C_i (with d minimal) such that $f_C(B) \neq 0$.
 - Find families of matrices $C(1), \dots, C(n)$ of dimension at most d_0 such that

$$f_C(B) = \sum c_\alpha f_{C(1)}(B)^{\alpha_1} \dots f_{C(n)}(B)^{\alpha_n}.$$

The Invariant Theory

- (Nayak and Subrahmanyam, 2010) R is a ring generated by the homogeneous polynomials $f_C(A) = \det(\sum C_i \circ A_i)$, for $d \times d$ matrices C_i , for all $d > 0$.
 - (1890) Hilbert showed that for a fairly general family of group actions, R is finitely generated, so we should expect there is some d_0 such that R is generated by the polynomials above for $d \leq d_0$. Killed the field for 100 years.
 - (Ivanyos, Qiao, Subrahmanyam, 2015)
 $d_0 \lesssim n^4[(n+1)!]^2$.
 - Suppose there are $d \times d$ matrices C_i (with d minimal) such that $f_C(B) \neq 0$.
 - Find families of matrices $C(1), \dots, C(n)$ of dimension at most d_0 such that

$$f_C(B) = \sum c_\alpha f_{C(1)}(B)^{\alpha_1} \dots f_{C(n)}(B)^{\alpha_n}.$$

- Since $f_C(B) \neq 0$, there must exist i with $f_{C(i)}(B) \neq 0$. 

The Invariant Theory

- (Nayak and Subrahmanyam, 2010) R is a ring generated by the homogeneous polynomials $f_C(A) = \det(\sum C_i \circ A_i)$, for $d \times d$ matrices C_i , for all $d > 0$.
 - (1890) Hilbert showed that for a fairly general family of group actions, R is finitely generated, so we should expect there is some d_0 such that R is generated by the polynomials above for $d \leq d_0$. Killed the field for 100 years.
 - (Ivanyos, Qiao, Subrahmanyam, 2015)
 $d_0 \lesssim n^4[(n+1)!]^2$.
 - Suppose there are $d \times d$ matrices C_i (with d minimal) such that $f_C(B) \neq 0$.
 - Find families of matrices $C(1), \dots, C(n)$ of dimension at most d_0 such that

$$f_C(B) = \sum c_\alpha f_{C(1)}(B)^{\alpha_1} \dots f_{C(n)}(B)^{\alpha_n}.$$

- Since $f_C(B) \neq 0$, there must exist i with $f_{C(i)}(B) \neq 0$.

• Thus $d \leq d_0$.

Rank Decreasing Operators

$$\int_{\mathbb{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbb{R}^{n_i})}^{p_i}.$$

Rank Decreasing Operators

$$\int_{\mathbb{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbb{R}^{n_i})}^{p_i}.$$

- (Bennett et al, 2008) implies that $\text{BL}(B, p) < \infty$ if and only if $\sum p_i n_i = n$, and for any subspace $V \subset \mathbb{R}^n$,

$$\dim(V) \leq \sum p_i \dim(B_i V).$$

Rank Decreasing Operators

$$\int_{\mathbb{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbb{R}^{n_i})}^{p_i}.$$

- (Bennett et al, 2008) implies that $\text{BL}(B, p) < \infty$ if and only if $\sum p_i n_i = n$, and for any subspace $V \subset \mathbb{R}^n$,

$$\dim(V) \leq \sum p_i \dim(B_i V).$$

- An operator $T : M_n \rightarrow M_n$ is *rank non-decreasing* if for any $A \succeq 0$, $\text{Rank}(TA) \geq \text{Rank}(A)$.

Rank Decreasing Operators

$$\int_{\mathbb{R}^n} \prod_{i=1}^m |f_i(B_i x)|^{p_i} dx \leq \text{BL}(B, p) \cdot \prod_{i=1}^m \|f_i\|_{L^1(\mathbb{R}^{n_i})}^{p_i}.$$

- (Bennett et al, 2008) implies that $\text{BL}(B, p) < \infty$ if and only if $\sum p_i n_i = n$, and for any subspace $V \subset \mathbb{R}^n$,

$$\dim(V) \leq \sum p_i \dim(B_i V).$$

- An operator $T : M_n \rightarrow M_n$ is *rank non-decreasing* if for any $A \succeq 0$, $\text{Rank}(TA) \geq \text{Rank}(A)$.
- (Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

- Results from (Gurvits and Samorodnitsky, 2002) show the result is true if $T(X) = \sum X_{ij} A_i$, where $A_i \succeq 0$. The general case can be reduced to this case.

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

- Results from (Gurvits and Samorodnitsky, 2002) show the result is true if $T(X) = \sum X_{ii} A_i$, where $A_i \succeq 0$. The general case can be reduced to this case.
- Given an orthonormal basis $U = \{u_1, \dots, u_N\}$, we define the *decoherence operator* $D_U(A) = \sum \langle Au_i, u_i \rangle \cdot u_i u_i^*$.

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

- Results from (Gurvits and Samorodnitsky, 2002) show the result is true if $T(X) = \sum X_{ii} A_i$, where $A_i \succeq 0$. The general case can be reduced to this case.
- Given an orthonormal basis $U = \{u_1, \dots, u_N\}$, we define the *decoherence operator* $D_U(A) = \sum \langle Au_i, u_i \rangle \cdot u_i u_i^*$.
- Define $T_U = D_U \circ T$.

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

- Results from (Gurvits and Samorodnitsky, 2002) show the result is true if $T(X) = \sum X_{ii} A_i$, where $A_i \succeq 0$. The general case can be reduced to this case.
- Given an orthonormal basis $U = \{u_1, \dots, u_N\}$, we define the *decoherence operator* $D_U(A) = \sum \langle Au_i, u_i \rangle \cdot u_i u_i^*$.
- Define $T_U = D_U \circ T$.
- Result follows from the following two facts:

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

- Results from (Gurvits and Samorodnitsky, 2002) show the result is true if $T(X) = \sum X_{ii}A_i$, where $A_i \succeq 0$. The general case can be reduced to this case.
- Given an orthonormal basis $U = \{u_1, \dots, u_N\}$, we define the *decoherence operator* $D_U(A) = \sum \langle Au_i, u_i \rangle \cdot u_i u_i^*$.
- Define $T_U = D_U \circ T$.
- Result follows from the following two facts:
 - (1) T is rank non-decreasing if and only if T_U is rank non-decreasing for all U .

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

- Results from (Gurvits and Samorodnitsky, 2002) show the result is true if $T(X) = \sum X_{ii} A_i$, where $A_i \succeq 0$. The general case can be reduced to this case.
- Given an orthonormal basis $U = \{u_1, \dots, u_N\}$, we define the *decoherence operator* $D_U(A) = \sum \langle Au_i, u_i \rangle \cdot u_i u_i^*$.
- Define $T_U = D_U \circ T$.
- Result follows from the following two facts:
 - (1) T is rank non-decreasing if and only if T_U is rank non-decreasing for all U .
 - (2) $\text{Cap}(T) = \inf_U \text{Cap}(T_U)$.

Proof Idea

(Gurvits, 2004) $T : M_n \rightarrow M_n$ is rank non-decreasing if and only if $\text{Cap}(T) > 0$.

- Results from (Gurvits and Samorodnitsky, 2002) show the result is true if $T(X) = \sum X_{ii} A_i$, where $A_i \succeq 0$. The general case can be reduced to this case.
- Given an orthonormal basis $U = \{u_1, \dots, u_N\}$, we define the *decoherence operator* $D_U(A) = \sum \langle Au_i, u_i \rangle \cdot u_i u_i^*$.
- Define $T_U = D_U \circ T$.
- Result follows from the following two facts:
 - (1) T is rank non-decreasing if and only if T_U is rank non-decreasing for all U .
 - (2) $\text{Cap}(T) = \inf_U \text{Cap}(T_U)$.
- To prove (1) and (2), use a simple trick: Given $A \succeq 0$, find U diagonalizing $T(A)$. Then $T(A) = T_U(A)$.

Thanks For Listening!