


OPEN



No link found between psychedelics and psychosis


In large US survey, users of LSD and similar drugs were no more likely to have mentalhealth conditions than other respondents.

Zoe Cormier

NATURE | NEWS

04 March 2015 | Corrected: 26 May 2015

Rights & Permissions



An image inspired by the reported experiences of psychedelic drug users. Anecdotal evidence of 'acid casualties' — users who develop chronic psychoses —is not supported by data from large populations, researchers say.

Data from population surveys in the United States challenge public fears that psychedelic drugs such as LSD can lead to psychosis and other mental-health conditions and to increased risk of suicide, two studies have found^{1, 2}.

In the first study, clinical psychologists Pål-Ørjan Johansen of EmmaSofia, a non-profit advocacy group based in Oslo, and Teri Suzanne Krebs of the Norwegian University of Science and Technology in Trondheim, scoured data from the US National Survey on Drug Use and Health (NSDUH), an annual random sample of the general population, and analysed answers from more than 135,000 people who took part in surveys from 2008 to 2011.

Of those, 14% described themselves as having used at any point in their lives any of the three 'classic' psychedelics: LSD, psilocybin (the active ingredient in so-called magic mushrooms) and mescaline (found in the peyote and San Pedro cacti). The researchers found that individuals in this group were not at increased risk of developing 11 indicators of mental-health problems such as schizophrenia, psychosis, depression, anxiety disorders and suicide attempts. Their paper appears in the March issue of the *Journal of Psychopharmacology* ¹.

Related stories

Rave drug holds promise

for treating depression

Psychosis risk syndrome

excluded from DSM-5

More related stories

• This Is Your Brain on

Drugs

The findings are likely to raise eyebrows. Fears that psychedelics can lead to psychosis date to the 1960s, with widespread reports of "acid casualties" in the mainstream news. But Krebs says that because psychotic disorders are relatively prevalent, affecting about one in 50 people, correlations can often be mistaken for causations. "Psychedelics are psychologically intense, and many people will blame anything that happens for the rest of their lives on a psychedelic experience."

The three substances Johansen and Krebs looked at all act through the brain's serotonin 2A receptor. The authors did not

include ketamine, PCP, MDMA, fly agaric mushrooms, DMT or other drugs that fall broadly into the category of hallucinogens, because they act on other receptors and have different modes of biochemical action. Ketamine and PCP, for example, act on the NMDA receptor and are both known to be addictive and to cause severe physical harms, such

"Absolutely, people can become addicted to drugs like ketamine or PCP, and the effects can be very destructive. We restricted our study to the 'classic psychedelics' to clarify the findings," says Johansen.

The 'acid casualty' myth

as damage to the bladder³.

"This study assures us that there were not widespread 'acid casualties' in the 1960s," says Charles Grob, a paediatric psychiatrist at the University of California, Los Angeles. He has long has advocated the therapeutic use of psychedelics, such as administering psilocybin to treat anxiety in terminal-stage cancer⁴. But he has concerns about Krebs and Johansen's overall conclusions, he says, because individual cases of adverse effects use can and do occur.

For example, people may develop hallucinogen persisting perception disorder (HPPD), a 'trip' that never seems to end, involving incessant distortions in the visual field, shimmering lights and coloured dots. "I've seen a number of people with these symptoms following a psychedelic experience, and it can be a very serious condition," says Grob.

Krebs and Johansen, however, point to studies that have found symptoms of HPPD in people who have never used psychedelics⁵.

The second of the new two studies, also published in the *Journal of Psychopharmacology*², looked at 190,000 NSDUH respondents from 2008 to 2012. It also found that the classic psychedelics were not associated with adverse mental-health outcomes. In addition, it found that people who had used LSD and psilocybin had lower lifetime rates of suicidal thoughts and attempts.

"We are not claiming that no individuals have ever been harmed by psychedelics," says author Matthew Johnson, an associate professor in the Behavioral Pharmacology Research Unit at Johns Hopkins University in Baltimore, Maryland. "Anecdotes about acid casualties can be very powerful — but these instances are rare," he says. At the population level, he says, the data suggest that the harms of psychedelics "have been overstated".

Nature | doi:10.1038/nature.2015.16968

Corrections

Corrected: An earlier version of this article stated that Johansen was affiliated with the Norwegian University of Science and Technology.

References

- 1. Johansen, P. Ø. & Krebs, T. S. J. Psychopharmacol.
- http://dx.doi.org/10.1177/0269881114568039 (2015). + Show context
- 2. Hendricks, P. S., Thorne, C. B., Clark, C. B., Coombs, D. W. & Johnson, M. W. J. Psychopharmacol. http://dx.doi.org/10.1177/0269881114565653 (2015).

+ Show context

3. Morgan, C. J. et al. Addiction 107, 27–38 (2012).

+ Show context

Article PubMed ISI

4. Grob, C. S. et al. Arch. Gen. Psychiatry 68, 71–78 (2011). + Show context

5. Schankin, C. J. et al. Brain 137, 1419–1428 (2014).

Article PubMed ISI ChemPort

+ Show context

Article PubMed ISI

Related stories and links

From nature.com Rave drug holds promise for treating depression fast

- 07 January 2015 Psychosis risk syndrome excluded from DSM-5
- 09 May 2012
- This Is Your Brain on Drugs 16 April 2012
- LSD helps to treat alcoholism 09 March 2012
- Benefits of 'magic mushroom' therapy long lasting 03 July 2008

From elsewhere


- Charles Grob Matthew Johnson
- For the best commenting experience, please login or register as a user and agree to our Community Guidelines. You will be re-directed back to this page where you will see comments updating in real-time and have the ability to recommend comments to other users.


Commenting is currently unavailable.

Nature ISSN 0028-0836 EISSN 1476-4687

About us Privacy policy Naturejobs About Nature Contact us Use of cookies Nature Asia Contact Nature Accessibility statement Nature Education About the Editors Legal notice Help Terms RSS web feeds Nature awards

Search Search go








Nature Podcast

Our award-winning show features highlights from the week's edition of *Nature*, interviews with the people behind the science, and in-depth commentary and analysis from journalists around the world.



Science jobs from naturejobs

South China Normal University sincerely invite oversea talented scholars to apply for the **Recruitment Program for Young Professionals** South China Normal University

Postdoctoral Research Associate The Scripps Research Institute - Florida

Worldwide Search for Talent at City University of Hong Kong City University of Hong Kong

Assistant Professor - Immunobioengineering

University of Toronto **Associate / Full Professor -Immunobioengineering**

University of Toronto

Post a Job 🕨 More Science Jobs

SPRINGER NATURE

© 2019 Macmillan Publishers Limited, part of Springer Nature. All Rights Reserved. partner of AGORA, HINARI, OARE, INASP, CrossRef and COUNTER