

Predictors of Stroke Mortality in Elderly People from the General Population: The Cardiovascular Study in the Elderly

Author(s): A. Mazza, A. C. Pessina, A. Pavei, R. Scarpa, V. Tikhonoff and E. Casiglia

Source: *European Journal of Epidemiology*, 2001, Vol. 17, No. 12 (2001), pp. 1097-1104

Published by: Springer

Stable URL: <https://www.jstor.org/stable/3582740>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



JSTOR

Springer is collaborating with JSTOR to digitize, preserve and extend access to *European Journal of Epidemiology*

Predictors of stroke mortality in elderly people from the general population

The CARDiovascular STudy in the ELderly

A. Mazza, A.C. Pessina, A. Pavei, R. Scarpa, V. Tikhonoff & E. Casiglia

Department of Clinical and Experimental Medicine, Laboratory of Epidemiology, University of Padova, Italy

Accepted in revised form 30 May 2002

Abstract. Stroke occurs particularly frequently in elderly people and, being more often disabling than fatal, entails a high social burden. The predictors of stroke mortality have been identified in 3282 subjects aged ≥ 65 years, taking part in the CARDiovascular STudy in the ELderly (CASTEL), a population-based study performed in Northeast Italy. Historical and clinical data, blood tests and 14-year fatal events were recorded. Continuous items were divided into quintiles and, for each quintile, adjusted relative risk (RR) with 95% confidence intervals [CI] was derived from multivariate Cox analysis. Age, historical stroke (RR: 5.2; 95% CI: 3.18–8.6) and coronary artery disease (RR: 1.38; CI: 1.18–2.1), atrial fibrillation (RR: 2.40; CI: 1.42–4.0), arterial hypertension (RR:

1.33; CI: 1.15–1.76), systolic blood pressure ≥ 163 mmHg (RR: 1.84; CI: 1.20–2.59), pulse pressure ≥ 74 mmHg (RR: 1.50; CI: 1.13–2.40), cigarette smoking (RR: 1.60; CI: 1.03–2.47), electrocardiographic left ventricular hypertrophy (RR: 1.72; CI: 1.10–2.61), impaired glucose tolerance (IGT, RR: 1.83; CI: 1.10–3.0), uric acid (UA) >0.38 mmol/l (RR: 1.61; CI: 1.14–2.10), serum potassium ≥ 5 mEq/l (RR: 1.70; CI: 1.24–2.50) and serum sodium ≤ 139 mEq/l (RR: 1.34; CI: 1.10–2.10) increased the risk of stroke. In the CASTEL, stroke was the first cardiovascular cause of death. Some independent predictors usually unrelated to stroke mortality (namely pulse pressure, pre-diabetic IGT, UA and blood electrolytes disorders) have been identified.

Key words: Elderly, Epidemiology, Risk factors, Stroke mortality

Abbreviations: BL – borderline; BMI – body mass index; BP – blood pressure; CAD – coronary artery disease; CI – confidence intervals; EKG – electrocardiogram; FEV₁ – maximal expiratory flow in 1 sec; HT – sustained hypertensive; IGT – impaired glucose tolerance; LVH – left ventricular hypertrophy; LVH_{EKG} – electrocardiographic left ventricular hypertrophy; LVMI – left ventricular mass index; NT – normotensive; RR – relative risk; SD – standard deviation; TC – total cholesterol; UA – uric acid

Introduction

Stroke is the primary cause of cardiovascular death in Eastern and the secondary one in Western societies [1]. Due to its high morbidity and mortality, stroke is a worldwide health care problem [2] exacting an enormous financial toll [3], as more than 50% of survivors have severe and permanent disability [4].

Most epidemiological studies identified the key risk factors for stroke, and have provided an estimate of their relative weight in middle-aged subjects [5–7]. Although the incidence of stroke rises sharply with advancing age, studies assessing the risk of stroke in the elderly are sparse, and almost limited to men [8–13]. A part from the recent data derived from the cardiovascular health study [14], the risk pattern of stroke in the elderly from the general population are therefore not well defined.

In this paper, we identified in a cohort of elderly subjects from Northeast Italy the factors able to increase the risk of fatal stroke. This study is part of the

CARDiovascular STudy in the ELderly (CASTEL), a population-based prospective study aimed at identifying cardiovascular risk factors and, more in general, the predictors of mortality in the elderly. The 14-year results are described herein.

Materials and methods

General protocol

The CASTEL enrolled 3282 subjects aged 65 years or above, representing 73% of elderly subjects from the Northern Italian towns of Castelfranco and Chioggia. Previous surveys [15, 16] showed for this cohort a cardiovascular risk pattern comparable to that of the Italian general population [17, 18]. The general characteristics of the two cohorts are summarised in Table 1.

Briefly, an initial survey was organised, and mortality was monitored yearly for 14 years. Historical

Table 1. Baseline characteristics of two elderly populations of the CASTEL

Items	Chioggia (N = 1028)	Castelfranco (N = 2254)	<i>p</i> -Value between two towns
Age (years)	74.3 ± 5.3	73.5 ± 5.2	<0.002
Males:females	426:602	855:1399	NS
Systolic BP (mmHg)	155.7 ± 23.6	162.2 ± 24.7	<0.0001
Diastolic BP (mmHg)	85.7 ± 10.8	90.3 ± 12.0	<0.0001
Heart rate (bpm)	75.7 ± 11.9	76.1 ± 11.2	NS
BMI (kg/m ²)	27.3 ± 4.7	26.6 ± 4.5	NS
TC (mmol/l)	5.28 ± 1.15	5.85 ± 1.12	<0.0001
Triglycerides (mmol/l)	1.3 ± 0.49	1.43 ± 0.54	<0.0001
Creatinine (μmol/l)	79.6 ± 35.4	79.6 ± 26.5	NS
UA (mmol/l)	0.32 ± 0.09	0.31 ± 0.08	NS
Frank diabetes (%)	27.3	17.3	<0.001
Hystorical CAD (%)	9.6	11.4	NS
Hystorical stroke (%)	4.6	8.6	<0.0001
Smoking habits			
Never smokers (%)	62.7	69.3	
Former smokers (%)	19.1	17.5	<0.002
Current smokers (%)	18.2	13.2	
No of cigarettes (daily)	6.7 ± 0.5	5.7 ± 0.4	NS

data were recorded by means of Rose’s questionnaire [19]. During the initial survey, trained doctors measured BP in triplicate with a sphygmomanometer at 15-min intervals, together with pulse rate. This procedure was repeated thrice at 1-month intervals. To minimise the effects of the alarm reaction, only the average of the last two measurements was taken into consideration. Subjects were considered to be sustained hypertensive when having a systolic blood pressure (BP) ≥ 160 mmHg and/or a diastolic BP ≥ 95 mmHg and/or a history of hypertension. Those with systolic BP values ranging between 140 and 159 mmHg and diastolic BP between 90 and 94 mmHg were labelled as borderline (BL). The difference between systolic and diastolic BP was the pulse pressure.

At the beginning of the screening, subjects were categorised into those who had a normal glucose tolerance (blood glucose <6.10 mmol/l without any antidiabetic treatment or any history of diabetes) and those with impaired glucose tolerance (IGT). The latter were considered as diabetic if taking antidiabetic drugs or having a fasting blood glucose level ≥ 6.99 mmol/l in at least two separate samples, without therapy [20].

Subjects were classified into never, current (≥ 1 cigarette daily) or former smokers (having given up their smoking habit for at least 1 year prior to the screening). Current smokers were subsequently segregated into quintiles based on number of cigarettes smoked daily.

Body mass index (BMI) was calculated as the ratio between weight and squared height. ‘Overweight’ was defined as an index >25 kg/m² [21]. Body surface area was calculated in m² with the formula of Du Bois and Du Bois [22]. All subjects were screened for albuminuria with a dipstick using a random morning urine sample, as 24-hour samples are impractical for

population screenings due to inconvenience and frequent collection errors [23]; urine infection was previously excluded.

Subjects having serum uric acid (UA) levels >0.38 mmol/l (last quintile of UA) were labelled as hyperuricemic, those with serum potassium levels >5 mmol/l (last quintile) as hyperkaliemic, and those having serum sodium levels <139 mmol/l (first quintile) as hyponatremic. Maximal expiratory flow in 1 sec (FEV₁) was measured with a dry spirometer, and reference values were individually calculated [24].

In a randomly chosen subset of 504 subjects, left ventricular mass was calculated from the following algorithm [25]: 1.04 [(end-diastolic diameter + end-diastolic posterior wall thickness + end-diastolic septum thickness)³ – end-diastolic diameter³] – 13.6. Left ventricular mass index (LVMI, in g/m²) was calculated by dividing left ventricular mass by body surface area [25]. According to Hammond [26], echocardiographic diagnosis of left ventricular hypertrophy was based on an LVMI >134 g/m² in males and >110 g/m² in females.

Electrocardiogram (EKG) was analysed on the basis of the Minnesota Code [27] by an expert who did not know the aim of the study. Diagnosis of atrial fibrillation was based on the code 8-3 and that of electrocardiographic left ventricular hypertrophy (LVH_{EKG}) on the code 3-1 or 3-3.

Coronary artery disease (CAD) at entry was defined as a history of myocardial infarction and/or angina and/or electrocardiographic signs of ischemia (Minnesota code 4-1, 4-3, 5-1 or 5-3, if codes 6-4, 7-1 and 7-2 were absent) and/or echocardiographic hypokinesia.

Annual mortality obtained from the registrar’s office was double-checked for causes of death by re-

ferring to hospitals, retirement homes or physicians' files. Causes of death were codified according to the International Classification of Disease. In 85% of the cases, autopsy or computerised tomography was performed in order to distinguish haemorrhagic from ischaemic events. Annual cumulative survival curves were generated with the life-table method and compared with the Mantel-Haenszel approach. Continuous variables were expressed as mean \pm standard deviation (SD) and divided into quintiles. For each quintile the RR (95% CI) adjusted for confounders was calculated from logistic models [28, 29].

Results

Univariate analysis

The mean age at entry was 73.8 ± 5.3 years (range: 65–91). In the five quintiles of age (656 subjects each) the mean age was 67.5 ± 1.2 , 70.4 ± 0.8 , 73 ± 1.1 , 76 ± 1.4 and 83 ± 3.2 years, respectively (all quintiles $p < 0.0001$ vs. each others). The last quintile, ranging from 80 to 95 years of age, corresponded to the current definition of very old. At the 14th year, 1616 subjects were dead and 1666 were alive (mortality rate: 49.2%).

Historical stroke was reported in 126 cases (3.8%), 68 women and 58 men, respectively (insignificant difference between genders). During the follow-up period, there were 170 new cases of fatal stroke (stroke mortality rate: 5.2%, insignificant difference between genders). Cumulative curves of survival showed that stroke was the first cause of cardiovascular mortality (Figure 1).

Multivariate prediction of stroke mortality

In the whole population, 13 items (age, historical stroke and CAD, BL IGT, arterial hypertension, systolic BP, pulse pressure, atrial fibrillation, LVH_{EKG} , UA, status of current smoker and serum potassium and sodium) turned out to be independent predictors of stroke mortality in multivariate Cox

analysis (Table 2). Systolic and pulse pressure as continuous variables had the same predictive value as the label of hypertension.

Role of clinical history

Historical stroke (RR: 5.2, [CI: 3.18–8.6]) and historical CAD (RR: 1.38, [CI: 1.18–2.1]) were the most important risk factors of stroke mortality.

Role of hypertension

Arterial hypertension was found to be an independent predictor of fatal stroke, mainly among sustained hypertensive (HT) (RR: 1.33, [CI: 1.15–1.76]) and BL hypertensive subjects (RR: 1.18, [CI: 1.02–1.63]). Besides, stroke mortality progressively increased from first to last quintile of systolic BP (3.7, 4.4, 4.6, 6.6, 6.7%, respectively; each quintile $p < 0.005$ vs. the previous one). It also increased sharply when pulse BP was > 74 mmHg (Table 2). Based on multivariate analysis, pulse pressure appeared to be a better predictor of fatal stroke than systolic BP (Table 3).

Role of atrial fibrillation

Atrial fibrillation was diagnosed in 145 subjects, without any difference between males and females (4.1 vs. 4.6%). Stroke mortality was higher in subjects with (11.7%) than without atrial fibrillation (4.9%; $p < 0.0005$); among the former, it was significantly higher in females than males (16.1 vs. 3.8%; $p < 0.02$) and increased with increasing age until 79 years (RR: 2.3, 6.4 and 7.8 in the age classes < 73 , 74–76 and 77–79 years, respectively; all differences $p < 0.001$). In multivariate analysis also, atrial fibrillation strongly predicted stroke mortality ($\chi^2 = 9.9$, $p < 0.002$, Cox coefficient: 0.93 ± 0.27) until 79 years of age.

Role of IGT

Stroke mortality rate was 6.6% in diabetic patients and 7.4% in those with pre-diabetic IGT; both rates were significantly higher ($p < 0.03$) than that of normoglycemic subjects (4.6%). In multivariate analysis, pre-diabetic IGT predicted fatal stroke ($\chi^2 = 4.29$, [CI: 1.10–3], Cox coefficient 0.57 ± 0.26) while frank diabetes was always rejected from the Cox model. When compared to frankly diabetic subjects, those with pre-diabetic IGT had significantly higher levels of diastolic BP (90.2 ± 11.7 vs. 88.5 ± 11.5 mmHg; $p < 0.05$), UA (0.35 ± 0.08 vs. 0.31 ± 0.08 ; $p < 0.01$) and total cholesterol (5.76 ± 1.09 vs. 5.53 ± 1.16 ; $p < 0.01$).

Role of LVH_{EKG}

Women with LVH_{EKG} had higher stroke mortality than those without this item (8.1 vs. 5%, RR: 1.72, [CI:




Figure 1. 14-years cumulative survival for overall (---), coronaric (---) and stroke (—) mortality in 3282 elderly subjects from the CASTEL.

Table 2. Stroke mortality rate, adjusted RR and 95% CI in 3282 elderly subjects of the CASTEL

Items	Mortality (%)	Relative risk	CI (95%)
Cardiovascular risk factors			
Hystorical stroke	12.4	5.2*	(3.18–8.6)
Hystorical CAD	13.4	1.38*	(1.18–2.1)
Pre-diabetic IGT	7.4	1.83**	(1.10–3.0)
Sustained hypertension	5.9	1.33**	(1.19–1.83)
Borderline hypertension	4.6	1.18**	(1.02–1.63)
Systolic BP (mmHg)	4.4	1.20	(0.62–1.88)
2nd quintile (141–150)	4.6	1.23	(0.68–2.0)
3rd quintile (151–162)	6.6	1.84**	(1.20–2.59)
4th quintile (163–180)	6.7	1.88*	(1.27–2.75)
5th quintile (≥181)			
Pulse pressure (mmHg)			
2nd quintile (55–64)	5.1	1.26	(1.01–2.08)
3rd quintile (65–73)	4.8	1.37	(1.09–2.12)
4th quintile (74–87)	6.3	1.50**	(1.13–2.40)
5th quintile (≥88)	7.0	1.94*	(1.24–2.80)
Atrial fibrillation	11.7	2.40**	(1.42–4.0)
Left ventricular hypertrophy	8.1	1.72**	(1.10–2.61)
Serum uric acid (mmol/l)			
1st quintile (≤0.23)	4.8	1.22	(1.02–1.65)
2nd quintile (0.24–0.28)	4.7	1.03	(0.67–1.70)
4th quintile (0.33–0.37)	5.1	1.09	(0.78–1.75)
5th quintile (≥0.38)	7	1.61	(1.14–2.10)**
Smoking habits			
Former	3.2	1.22	(0.67–1.75)
Current	6.4	1.60*	(1.03–2.47)
Blood electrolytes disorders			
Potassium (mEq/L)			
1st quintile (≤3.5)	4.1	0.57	(0.34–0.96)
2nd quintile (3.6–4.1)	4.9	0.72	(0.44–1.17)
4th quintile (4.5–4.9)	5.5	1.01	(0.69–1.21)
5th quintile (≥5)	6.7	1.70**	(1.24–2.50)
Sodium (mEq/L)			
1st quintile (≤139)	6.5	1.34**	(1.10–2.10)
2nd quintile (140–142)	5.4	1.02	(0.73–1.31)
4th quintile (143–144)	4.2	0.76	(0.52–1.01)
5th quintile (≥145)	3.9	0.98	(0.61–1.13)

For categorical variables, the opposite or missing category served as reference groups with RR = 1 (e.g., no historical stroke, never smoked, 1st quintile).
* $p < 0.001$; ** $p < 0.01$.

1.74–3.37; ($p < 0.004$), while no difference was found in men. No relation was found between echocardiographic left ventricular hypertrophy and stroke mortality in both genders.

Role of UA

A U-shaped trend of stroke mortality rate was found in relation to UA (in the five quintiles: 4.8, 4.7, 4, 5.1, and 7%).¹ In the Cox analysis, hyperuricemia predicted fatal strokes with an RR of 1.61, [CI: 1.04–2.61].

¹ $p < 0.04$ last vs. first and second, $p < 0.001$ last vs. third.

Role of cigarette smoking

Current smokers had a higher risk of fatal stroke (RR: 1.60, [CI: 1.03–2.60]) than non-smokers, whereas no difference was found between former- and never-smokers (Table 2). Excess mortality was particularly high in subjects who had smoked ≥40 years (7.2 vs. 1.8%, $p < 0.01$). No significant relationship was shown between fatal stroke and number of cigarettes smoked daily.

Role of blood electrolytes

The risk of fatal stroke was significantly higher when potassium levels were ≥5 mEq/l (RR: 1.70,

Table 3. Results of the multivariate Cox analysis of fatal stroke in 3282 elderly subjects from the CASTEL

	Improve- ment χ^2	p-Value	Cox-coefficient ± SE
Age	4.10	<0.04	0.08 ± 0.03
History of stroke	23.29	<0.0001	1.64 ± 0.31
Hystory of CAD	13.86	<0.04	1.19 ± 0.24
Arterial hypertension	12.22	<0.05	0.39 ± 0.11
Atrial fibrillation	11.01	<0.0001	1.26 ± 0.31
Systolic BP	10.15	<0.001	0.01 ± 0.003
Pulse pressure	9.97	<0.002	0.01 ± 0.004
Pre-diabetic IGT	3.17	<0.05	0.62 ± 0.31
LVH _{EKG}	3.10	<0.05	0.56 ± 0.31
Current smokers	6.34	<0.01	0.05 ± 0.01
Serum UA	0.43	<0.007	0.19 ± 0.07
Serum potassium	3.29	<0.05	0.43 ± 0.23
Serum sodium	3.00	<0.05	−0.04 ± 0.02

Only the items that were accepted by the equation are summarised.

[CI: 1.24–2.50]) and when serum sodium levels were ≤139 mEq/l (RR: 1.34, [CI: 1.10–2.10]). When both the above-mentioned disorders were simultaneously present, RR was 3.03, [CI: 1.67–5.46].

Very old subjects

In very old subjects, only three items (age, historical stroke and LVH_{EKG}) predicted stroke mortality, and increased pulse pressure (> 74 mmHg) had the same meaning as hypertension.

Discussion

In this study, historical stroke was the best predictor of stroke mortality. Many evidences show that survivors from an acute stroke are pre-disposed to other life-threatening diseases [30–32]. For example, in the Dubbo Study [33], patients with a history of stroke had a 27% risk excess of a new stroke compared to those free from prior cerebrovascular events.

Atrial fibrillation, pulse pressure, history of CAD, LVH_{EKG}, IGT, smoking, UA and electrolytes were other predictors. Atrial fibrillation, a common clinical problem in the elderly [34, 35], notoriously predisposes stroke [34, 36, 37]. Approximately 17% of all autopsied elderly subjects have atrial fibrillation [37], and emboli from the heart are responsible for up to 20% of cases of stroke [38]. In the CASTEL, however, differently from Framingham [39], atrial fibrillation had no prognostic value after the age of 80 years. This could be of interest when evaluating the opportunity of treating very old subjects with warfarin.

Hypertension is known to be a predictor of stroke [40], and practically all the studies have demonstrated a relationship between BP levels and stroke [41–43].

This was confirmed in the present study, where RR of stroke mortality was 1.18, [CI: 1.02–1.63] in BL and 1.33, [CI: 1.15–1.76] in HT, as compared to normotensive (NT) subjects. The pulse component of BP was of the greatest importance even after the age of 80 years, with a progressive increase of RR with increasing pulse pressure and a sharp rise above the value of 74 mmHg. It is difficult to discuss this point, as only systolic and diastolic BP have been systematically taken into consideration in the literature [44, 45], whereas pulse BP has been neglected in this respect, particularly in the elderly. High pulse pressure is known to be associated with an increase of the carotid intima-media thickness [46]; nevertheless, in the CASTEL, stroke mortality was not higher in subjects with carotid murmurs in the neck, the only index of carotid damage available on a large scale in 1983–1985.

Based on the results of many previous studies [41, 47], a history of myocardial infarction or angina pectoris is also considered to be strongly associated with an increased risk of stroke. This was also true in the CASTEL, where subjects with a history of CAD were at an increased risk of stroke. It is therefore a good practice to make all efforts to identify and to treat subjects with CAD, with the dual aim to prevent both coronary and cerebrovascular events.

Although not as sensitive as the echocardiogram, EKG remains an acceptable index of LVH, having a specificity around 12% [48]. In population studies, and in the CASTEL study too, LVH_{EKG} was a predictor of stroke [49–51]. The simplistic conclusion that this is due to coexistent arterial hypertension must be rejected, as LVH_{EKG} independently contributed to cerebrovascular risk.

Diabetes mellitus, a strong risk factor for atherosclerosis [52, 53], seems to have a less consistent role in the pathogenesis of cerebrovascular disease. In fact, some studies have indicated that subjects with IGT have an increased risk of stroke [52, 54, 55], whereas other investigations [33, 54] have denied this possibility. In the CASTEL, subjects with pre-diabetic IGT had an increased risk of stroke, independently from other cardiovascular risk factors. On the contrary, frank diabetes had no prognostic role, being probably overwhelmed, in the multivariate analysis, by other variables such as hypertension [52] and history of CAD [40]. IGT is one of the aspects of the ‘X syndrome’ [56], whose role in generating stroke has been clearly demonstrated [57].

Smoking acts as a risk factor for stroke in a dose-dependent manner and may also interact with hypertension, making the identification of its specific role difficult. Tobacco has short- as well as long-term effects on vessels [58] and contributes to the development of atherosclerosis and its complications [59]. In the CASTEL, only current smokers experienced an increased stroke mortality. This is in agreement with the belief that a reduction of risk of stroke follows

smoking cessation [60] and confirms the opportunity to promote community-based educational programs against tobacco not only among young and middle-aged subjects, but also among elderly people.

A possible prognostic role of UA has been emphasised in recent years [61, 62], but its meaning as an independent risk factor for cardiovascular disease and in particular for stroke remains uncertain. In the ARIC study [63], UA did not result to be *per se* a risk factor for atherosclerosis, since, when the known risk factors for cardiovascular diseases were included in multivariate analysis, UA lost any prognostic role. Nevertheless, in a recent study on middle-aged diabetic patients [64], hyperuricemia was a strong predictor of stroke events, independently of other cardiovascular risk factors such as hypertension [65, 66], nephropathy [61], overweight [62] and diuretic therapy [66] (all conditions accompanied by higher UA levels). In the CASTEL, hyperuricemia (last quintile of UA) was found to be a strong and independent predictor of stroke mortality. The mechanism of this association remains unexplained, but it has been recently shown that human atherosclerotic plaque contains a considerable amount of UA [67], and hyperuricemia – via purine metabolism – may promote thrombus formation [68]. Finally, the reason why the curve of stroke mortality was patently J-shaped remains unknown, although it can be postulated that subjects with the lowest UA levels have a muscular mass below the standard and are consequently disadvantaged in survival.

A positive relationship between dietary sodium/potassium intake and stroke has been reported [69]. Electrolyte disorders are common in the elderly, particularly in diabetic [70] and nephropathic [71] subjects, probably due to both hypervolaemia and renal decay. In the CASTEL, hyperkalaemia and hyponatremia were independent predictors of stroke, and the RR was particularly high (3.03, [CI: 1.67–5.46]) when the above-mentioned disorders were simultaneously present. Animal studies have shown that local cerebral ischaemia produces electrolyte disturbances via lactic acidosis [72], and a recent report in humans has also shown a significant association between cerebral injuries and hyponatremia [73].

Conclusions

The CASTEL has confirmed the important role of age, hypertension, atrial fibrillation, cigarette smoking, LVH and history of stroke and CAD as risk factors for stroke, while it has failed to confirm that of gender, diabetes, serum lipids, alcohol intake, BMI and carotid murmurs. Instead, the CASTEL has identified additional risk factors such as pulse BP, IGT, UA and the blood electrolyte disorders. Finally, in our study, the risk pattern appears to be quite different in subjects who are just elderly and in those who are very old.

Further data are needed to generate strategies for treatment and programs of public health policy related to prevention of stroke in elderly people.

References

1. Bonita R. Epidemiology of stroke. *Lancet* 1992; 339: 342–344.
2. Bonita R, Beaglehole R, Asplund K. The worldwide problem of stroke. *Cur Opin Neurol Neurosurg* 1994; 7: 5–10.
3. Taylor TN, Davis PH, Torner JC, Holmes J, Meyer JW, Jacobson MF. Lifetime cost of stroke in the United States. *Stroke* 1996; 27: 1459–1466.
4. Simons LA, McCallum J, Friedlander Y, Simons J, Powell I, Heller R. Dubbo Study of the elderly: Sociological and cardiovascular risk factors at entry. *Aust N Z J Med* 1991; 21: 701–709.
5. Davis PH, Dambrosia JM, Schoenberg BS, et al. Risk factors for ischemic stroke: A prospective study in Rochester, Minnesota. *Ann Neurol* 1978; 22: 319–327.
6. Kagan A, Poppers JS, Rhoads GG. Factors related to stroke incidence in Hawaii Japanese men: The Honolulu Heart Study. *Stroke* 1980; 11: 14–21.
7. Wolf PA. Risk factors for stroke. *Stroke* 1985; 16: 359–360.
8. Lechner H, Schmidt R, Reinhart B, et al. Cerebrovascular risk factors in an elderly Austrian population: First year results of the Austrian Stroke Prevention Study (ASPS). *Wien Klin Wochenschr* 1993; 105: 398–403.
9. Manolio TA, Kronmal RA, Burke GL, O'Leary DH, Price TR. Short-term predictors of incident stroke in older adults: The Cardiovascular Health Study. *Stroke* 1996; 27: 1479–1486.
10. Wolf PA, Abbot RD, Kannel WB. Atrial fibrillation: A major contributor to stroke in the elderly. *Arch Intern Med* 1987; 147: 1561–1564.
11. Welin L, Svardsudd K, Wilhelmsen L, Larsson B, Tibblin G. Analysis of risk factors for stroke in a cohort of men born in 1913. *N Engl J Med* 1987; 317: 521–526.
12. Zeiler K, Siostrzonek P, Lang W, et al. Different risk factor profiles in young and elderly stroke patients with special reference to cardiac disorders. *J Clin Epidemiol* 1992; 45: 1383–1389.
13. Woo J, Lau EM. Risk factors predisposing to stroke in an elderly Chinese population a longitudinal study. *Neuroepidemiology* 1990; 9: 131–134.
14. Longstreth WT, Bernick C, Fitzpatrick A, et al. Frequency and predictors of stroke death in 5888 participants in the Cardiovascular Health Study. *Neurology* 2001; 56: 368–375.
15. Casiglia E, Spolaore P, Ginocchio G, et al. Predictors of mortality in very old subjects aged 80 years or over. *Eur J Epidemiol* 1993; 9: 577–586.
16. Casiglia E, Spolaore P, Ginocchio G, et al. Proteinuria predicts mortality among elderly but not among 'very old' subjects. *J Nephrol* 1992; 5: 89–93.
17. Descovich GC, Aluigi L, Benassi MS, et al. The Brighella investigation. Results of the observational study 1972–1980. *Minerva Cardioangiolog* 1985; 33: 565–574.

18. Giampaoli S, Vanuzzo D. Cardiovascular risk factors in Italy: An interpretation with reference to the national health plan 1998–2000. *G Ital Cardiol* 1999; 29: 1463–1471.
19. Rose GA, Blackburn H, Gillum RF, Prineas RJ (eds), *Cardiovascular Survey Methods*, Geneva: WHO, 1982.
20. Drouin P, Blicke JF, Charbonnel B, et al. Diagnosis and classification of diabetes mellitus: The new criteria. *Diabetes Metab* 1999; 25: 72–83.
21. Lean MEJ, Han TS, Seidell JC. Impairment of health and quality of life using new US Federal Guidelines for the identification of obesity. *Arch Int Med* 1999; 159: 837–843.
22. Du Bois D, Du Bois F. A formula to estimate the approximate surface area if height and weight are known. *Arch Intern Med* 1916; 17: 863–871.
23. Blainey JD, Brewer DB, Hardwicke J. Proteinuric glomerular disease in adults: Cumulative life tables over twenty years. *Quart J Med* 1986; 230: 557–567.
24. Cherniack RM, Raber MB. Normal standard for ventilatory function using an automated wedge spirometer. *Ann Rev Resp Dis* 1972; 106: 38–46.
25. Devereux RB. Evaluation of cardiac structure and dysfunction by echocardiography and other noninvasive techniques. In: Laragh JH, Brenner BM (eds), *Hypertension: Pathophysiology, Diagnosis, and Management*, New York: Raven Press, 1990.
26. Hammond IW, Devereux RB, Alderman MH, et al. The prevalence and correlates of echocardiographic left ventricular hypertrophy among patients with uncomplicated hypertension. *J Am Coll Cardiol* 1986; 7: 639–643.
27. Prineas RJ, Crow RS, Blackburn H. The Minnesota code manual of electrocardiographic findings: Standards and procedures for measurement and classification. Littleton, Mass: John Wright-PSG Inc, 1982.
28. Cox DR. Regression models and life tables. *J Roy Statist Soc* 34 Series B, 1972.
29. Cox DR, Oakes D. *Analysis of Survival Data*. London: Chapman and Hall, 1984.
30. Hanger HC, Walker G, Paterson LA, McBride S, Sainsbury R. What do patients and their carers want to know about stroke? A two-year follow-up study. *Clin Rehabil* 1998; 12: 45–52.
31. Paolucci S, Antonucci G, Gialloreti LE, et al. Predicting stroke inpatient rehabilitation outcome: The prominent role of neuropsychological disorders. *Eur Neurol* 1996; 36: 385–390.
32. Viitanen M, Fugl Meyer KS, Bernspang B, Fugl Meyer AR. Life satisfaction in long-term survivors after stroke. *Scand J Rehabil Med* 1988; 20: 17–24.
33. Simons LA, McCallum JM, Friedlander Y, Simons J. Risk factors for ischemic stroke. Dubbo study of the elderly. *Stroke* 1998; 29: 1341–1346.
34. Yamanouchi H, Tomonaga M, Shimada H, Matsushita S, Kuramoto K, Toyokura Y. Nonvalvular atrial fibrillation as a cause of fatal massive cerebral infarction in the elderly. *Stroke* 1989; 20: 1653–1656.
35. Kannel WB, Abbot RD, Savage DD, McNamara PM. Epidemiologic features of chronic atrial fibrillation. The Framingham study. *N Engl J Med* 1982; 306: 1018–1022.
36. Wolf PA, Abbot RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: The Framingham study. *Stroke* 1991; 22: 983–988.
37. Kuramoto K, Matsushita S, Yamanouchi H. Atrial Fibrillation as a cause of myocardial and cerebral infarction. *Jpn Circ J* 1984; 48: 67–74.
38. Martin R, Bogousslavsky J. Embolic versus nonembolic causes of ischemic stroke. *Cerebrovasc Dis* 1995; 5: 70–74.
39. Benjamin EJ, Wolf PA, D'Agostino RB, Silbershatz H, Kannel WB, Levy D. Impact of atrial fibrillation on the risk of death: The Framingham heart study. *Circulation* 1998; 98: 946–952.
40. Lavy S, Melamed E, Cahane E, Carmon A. Hypertension and diabetes as risk factors in stroke patients. *Stroke* 1973; 4: 751–759.
41. Wolf PA, D'Agostino RB, Belanger AJ, Kannel WB. Probability of stroke: A risk profile from the Framingham study. *Stroke* 1991; 22: 312–318.
42. Hypertension Detection and Follow-up Program Cooperative Group: Five-year findings of the Hypertension Detection and Follow up Program: III. Reduction in stroke incidence among persons with high blood pressure. *JAMA* 1982; 247: 633–638.
43. Perry HM Jr, Smith WM, Mc Donald RH, et al. Morbidity and mortality in the systolic hypertension in the elderly program (SHEP) Pilot Study. *Stroke* 1989; 20: 4–13.
44. Kannel WB, Dawber TR, Sorlie P, Wolf PA. Components of blood pressure and risk of Atherothrombotic brain infarction: The Framingham study. *Stroke* 1976; 7: 327–331.
45. Kannel WB, Wolf PA, McGee DL, Dawber TR, McNamara PM, Castelli WP. Systolic blood pressure, arterial rigidity and risk of stroke: The Framingham study. *JAMA* 1981; 245: 1225–1229.
46. Zakopoulos NA, Lekakis JP, Papamichael CM, et al. Pulse pressure in normotensives: A marker of cardiovascular disease. *Am J Hypertens* 2001; 14: 195–199.
47. Aronow WS. Prevalence of heart disease in older women in a nursing home. *J Womens Health* 1998; 7: 1105–1112.
48. Casiglia E, Maniati G, Daskalakis C, et al. Left-ventricular hypertrophy in the elderly: Unreliability of ECG criteria in 477 subjects aged 65 years or more. The cardiovascular study in the elderly (CASTEL). *Cardiology* 1996; 87: 429–435.
49. Kahn S, Frishman WH, Weissman S, Ooi WL, Aronson M. Left ventricular Hypertrophy on electrocardiogram: Prognostic implications from a 10-year cohort study of older subjects: A report from the Bronx longitudinal aging study. *J Am Geriatr Soc* 1996; 44: 524–529.
50. Levy D, Garrison RJ, Savage-DD, Kannel WB, Castelli WP. Prognostic implications of echocardiographically determined left ventricular mass in the Framingham heart study. *N Engl J Med* 1990; 322: 1561–1566.
51. Furberg CD, Manolio TA, Psaty BM. Major electrocardiographic abnormalities in persons aged 65 years and older (the cardiovascular health study). *AM J Cardiol* 1992; 69: 1329–1335.
52. Kannel WB, McGee DL. Diabetes and cardiovascular disease: The Framingham study. *JAMA* 1979; 241: 2035–2038.
53. Fuller JH, Shipley MJ, Rose G, Jarret RJ, Keen H. Mortality from coronary heart disease and stroke in

- relation to degree of glycemia: The Whitehall study. *Br Med J* 1983; 287: 867–870.
54. Qureshi AI, Giles WH, Croft JB. Impaired glucose tolerance and the likelihood of nonfatal stroke and myocardial infarction: The third national health and nutrition examination survey. *Stroke* 1998; 29: 1329–1332.
 55. Abbott RD, Donahue RP, MacMahon SW, Reed DM, Yano K. Diabetes and the risk of stroke: The Honolulu Heart Program. *JAMA* 1987; 257: 949–952.
 56. Suba I, Halmos T, Kautzky L. The value of certain parameters in the diagnosis and detection of metabolic X syndrome. *Orv Hetil* 1997; 138: 2407–2411.
 57. Vavrkova H, Ficker L, Vlachova I, Chudackova J, Novotny D, Budikova M. Reaven's metabolic syndrome X in the families of individuals with premature cerebrovascular attacks. *Vnitr Lek* 1993; 39: 745–754.
 58. Taylor BV, Oudit GY, Kalman PG, Liu-P. Clinical and pathophysiological effects of active and passive smoking on the cardiovascular system. *Can J Cardiol* 1998; 14: 1129–1139.
 59. Mast H, Thompson JL, Lin IF, et al. Cigarette smoking as a determinant of high-grade carotid artery stenosis in Hispanic, black, and white patients with stroke or transient ischemic attack. *Stroke* 1998; 29: 908–912.
 60. Fiore MC, Jorenby DE, Baker TB. Smoking cessation: Principles and practice based upon the AHCPR Guideline, 1996. Agency for health care policy and research. *Ann Behav Med* 1997; 19: 213–219.
 61. Cannon PJ, Stason WB, De Martini FE, Sommers SC, Laragh JH. Hyperuricemia in primary and renal hypertension. *N Engl J Med* 1996; 275: 457–464.
 62. Brand FN, McGee DL, Kannel WB, Stokes III J, Castelli WP. Hyperuricemia as a risk factor of coronary heart disease: The Framingham study. *Am J Epidemiol* 1985; 121: 11–18.
 63. Iribarren C, Folsom AR, Eckfeldt JH, McGovern PG, Nieto FJ. Correlates of uric acid and its association with asymptomatic carotid atherosclerosis: The ARIC study. *Am Epidemiol* 1996; 6: 331–340.
 64. Letho S, Niskanen L, Rönnemaa T, Laakso M. Serum uric acid is a strong predictor of stroke in patients with non-insulin-dependent diabetes mellitus. *Stroke* 1998; 29: 635–639.
 65. Staessen J. The determinants and prognostic significance of serum uric acid in elderly patients of the European working party on high blood pressure in the elderly trial. *Am J Med* 1991; 90(Suppl 3A): 50–54.
 66. Coronary Drug Project Research Group. Serum uric Acid: Its association with other risk factors and with mortality in coronary heart disease. *J Chron Dis* 1976; 29: 557–569.
 67. Suarna C, Dean RT, May J, Stocker R. Human atherosclerotic plaque contains both oxidized lipids and relatively large amounts of alpha-tocopherol and ascorbate. *Arterioscler Thromb Vasc Biol* 1995; 15: 1616–1624.
 68. Visy J, Le Coz P, Chadeaux B, et al. Homocystinuria due to 5,10-methylenetetrahydrofolate reductase deficiency revealed by stroke in adult siblings. *Neurology* 1991; 41: 1313–1315.
 69. Sasaki S, Zhang XH, Kesteloot H. Dietary sodium, potassium, saturated fat, alcohol, and stroke mortality. *Stroke* 1995; 26: 783–789.
 70. Elisaf S, Tsatsoulis AA, Katopodis KP, Siamopoulos KC. Acid-base and electrolyte disturbances in patients with diabetic ketoacidosis. *Diabetes Res Clin Pract* 1996; 34: 23–27.
 71. Brown WW, Schmitz PG. Acute and chronic kidney disease. *Clin Geriatr Med* 1998; 14: 211–236.
 72. Rudolf U, Zoltan N, Zsuzsa J. Ion and metabolic disturbances after global and focal cerebral ischemia. *Acta Biomed Ateneo Parmense* 1995; 66: 75–82.
 73. Bacic A, Gluncic I, Gluncic V. Disturbances in plasma sodium in patients with war head injuries. *Mil Med* 1999; 164: 214–217.
 74. Wolf PA, D'Agostino RB, Kannel WB, Bonita R, Belanger AJ. Cigarette smoking as a risk factor for stroke: The Framingham study. *JAMA* 1998; 259: 1025–1029.
 75. Shinton R, Beevers G. Meta-analysis of relation between cigarette smoking and stroke. *Br Med J* 1989; 298: 789–794.

Address for correspondence: E. Casiglia, Department of Clinical and Experimental Medicine, University of Padova, Via Giustiniani 2, 35128 Padova, Italy
 Phone: +39-49-8212278/8212275; Fax: +39-49-8754179/9754179
 E-mail: edoardo.casiglia@unipd.it