

# Mortality Risk in Spanish Adults With Diagnosed Diabetes, Undiagnosed Diabetes, or Pre-Diabetes. The Asturias Study 1998-2004

Sergio Valdés,<sup>a,b,c</sup> Patricia Botas,<sup>d</sup> Elías Delgado,<sup>a</sup> and Francisco Díaz Cadórniga<sup>a</sup>

<sup>a</sup>Servicio de Endocrinología y Nutrición, Hospital Universitario Central de Asturias, Oviedo, Asturias, Spain

<sup>b</sup>Servicio de Endocrinología y Nutrición, Hospital Universitario Carlos Haya, Málaga, Spain

<sup>c</sup>CIBER de Diabetes y Enfermedades Metabólicas Asociadas, CIBERDEM, Málaga, Spain

<sup>d</sup>Servicio de Medicina Interna, Sección de Endocrinología, Hospital San Agustín, Avilés, Asturias, Spain

**Introduction and objectives.** Although type-2 diabetes is a well-known cause of death, the mortality associated with undiagnosed diabetes and early-stage dysglycemia has not been clearly determined.

**Methods.** This study included 1015 individuals aged 30-75 years who took part in the first phase of the Asturias study (1998-1999). Participants completed a questionnaire and underwent a physical examination and an oral glucose tolerance test (OGTT). All deaths that occurred in the cohort within 6 years of follow-up (ie December 1998 to December 2004) were recorded.

**Results.** Participants were divided into four groups according to the condition indicated by their OGTT result in the first phase of the study: normoglycemia, pre-diabetes, undiagnosed diabetes, or diagnosed diabetes (World Health Organization 1999 criteria). A total of 42 deaths were recorded during follow-up. With normoglycemic individuals acting as a control group, multivariate analysis showed that the relative risk of mortality was 2.5 (95% CI, 1-6.3) in the group with diagnosed diabetes, 2.7 (95% CI, 1.1-6.7) in the group with undiagnosed diabetes, and 1.6 (95% CI, 0.7-4) in the group with pre-diabetes.

**Conclusions.** Both individuals with diagnosed diabetes and those with undiagnosed diabetes had a risk of mortality around 2.5-3 times greater than individuals with normoglycemia. Those with pre-diabetes also had increased mortality relative to the control group, though the difference was not significant.

**Key words:** Type-2 diabetes. Mortality. Undiagnosed diabetes. Pre-diabetes. Spain.

**Riesgo de mortalidad en diabetes diagnosticada, diabetes no diagnosticada y prediabetes en población adulta española. Estudio Asturias 1998-2004**

**Introducción y objetivos.** Aunque la diabetes mellitus DM tipo 2 es una causa establecida de mortalidad, el riesgo de mortalidad asociado a DM no diagnosticada y a estadios previos de disglucemia no está claramente definido.

**Métodos.** El estudio incluyó a 1.015 individuos de 30-75 años de edad que participaron en la primera fase del Estudio Asturias (1998-1999), realizando encuesta, exploración física y SOG. Se registraron los fallecimientos en la cohorte durante 6 años de seguimiento (diciembre de 1998 a diciembre de 2004).

**Resultados.** Se clasificó a los sujetos en cuatro grupos según el resultado de la SOG en la primera fase del estudio: normoglucemia, prediabetes, DM ignorada y DM conocida (criterios de la OMS 1999). Se registraron 42 muertes durante el seguimiento. Respecto al grupo control con normoglucemia, el riesgo relativo (RR) de mortalidad en el modelo multivariable fue 2,5 (intervalo de confianza [IC] del 95%, 1-6,3) en el grupo con DM conocida, RR = 2,7 (IC del 95%, 1,1-6,7) en el grupo con DM ignorada y RR = 1,6 (IC del 95%, 0,7-4) en el grupo con prediabetes.

**Conclusiones.** Tanto los individuos con DM conocida como los que tenían DM no diagnosticada presentaron un riesgo de mortalidad alrededor de 2,5-3 veces superior al de los individuos con normoglucemia. En individuos con prediabetes también se encontró un incremento de mortalidad frente al grupo control, aunque no estadísticamente significativo.

**Palabras clave:** Diabetes mellitus tipo 2. Mortalidad. Diabetes no diagnosticada. Prediabetes. España.

The Asturias Study was undertaken with finance from the grant FIS 98/0462, a grant from the Consejería de Salud of the Principado de Asturias and a Pfizer grant. The CIBER de Diabetes y Enfermedades Metabólicas Asociadas (CB07/08/0019) is an initiative of the Instituto de Salud Carlos III.

Correspondence: Dr. S. Valdés.  
Servicio de Endocrinología y Nutrición. Hospital Universitario Carlos Haya.  
Pl. del Hospital Civil. 29009 Málaga. España.  
E-mail: Sergio.valdes@hotmail.es

Received November 17, 2008.

Accepted for publication February 11, 2009.

## INTRODUCTION

Type 2 diabetes mellitus (DM) has become one of the most severe health problems of our time. The situation in Spain is worrying: the various

## ABBREVIATIONS

CVD: cardiovascular disease

DM: diabetes mellitus

OGTT: oral glucose tolerance test

cross-sectional studies carried out in this country show that the prevalence of DM has increased over the last decade. Currently, the estimated prevalence of type 2 DM in the adult population is 10%-15%.<sup>1,2</sup> More worrying still is the fact that at least half of these persons are unaware that they have the disease (undiagnosed or unknown DM).<sup>1,2</sup> This is due to the fact that this form of diabetes often remains undiagnosed for many years because the hyperglycemia develops gradually, with a long asymptomatic, preclinical phase. Nevertheless, chronic hyperglycemia, even in the absence of symptoms, is associated with an increased risk of diabetic microangiopathy and cardiovascular disease (CVD), which can even start with degrees of dysglycemia below current diagnostic levels of DM.<sup>3</sup> Most studies on CVD, death and health costs related with DM are based solely on the proportion of diagnosed or known DM. Less is known about the impact of persons with unknown or undiagnosed DM or of the early stages of carbohydrate metabolism disorders. The demonstration of an increase in mortality in these stages could reinforce the importance of screening programs and early intensive treatment in these persons.

The aim of this study was to evaluate the risk of death in persons with diagnosed DM, undiagnosed DM, and prediabetes as compared with persons with normoglycemia in a representative cohort of the general population from the province of Asturias, in the north of Spain.

## METHODS

### Asturias Study

The Asturias Study is a population-based prospective cohort study of DM and cardiovascular risk factors that is taking place among the whole population of the province of Asturias, in northern Spain.<sup>4,7</sup> The first phase took place over the years 1998-1999 and was designed to determine the prevalence of type 2 DM, both diagnosed and undiagnosed, and of prediabetes in the population of Asturias, which is 1 073 761 inhabitants, mostly Caucasian. About half the population live in urban areas. The sample was selected using a 2-stage cluster sampling technique: 15 basic health areas were randomly selected from among the 76 in Asturias, with a probability

proportional to the number of health cards from persons aged 30 to 75 years in each area. A computer program was then used for the random selection of 125 persons in each basic health area. The final sample selected was 1875 persons. A total of 87 persons were excluded for various reasons (type 1 DM, pregnancy, severe disease, hospitalization, treatment with hyperglycemic drugs). Another 162 were excluded because their contact data were not complete. The final sample was 1626 persons, of whom 1034 (63.6%) participated in the study.

The study was approved by an Ethics Committee of the Principality of Asturias Health Service and all the participants gave their informed consent.

All the participants completed a health questionnaire, which included data on demographics, smoking, physical activity, socioeconomic status, and a family history of DM.

The height and weight were measured with the participant in light clothing and without shoes, and the BMI was calculated (weight in kilograms divided by the square of the height in meters). The blood pressure (BP) was measured using a digital sphygmomanometer (OMROM MX3, OMROM Healthcare, Tokyo, Japan) with the person seated and at rest. The mean of 2 BP measurements, taken 1-2 min apart, was used for this analysis.

All the participants, except those with diagnosed DM, underwent an oral glucose tolerance test (OGTT), with extraction of venous blood at baseline and after 2 h, in accordance with the recommendations of the World Health Organization<sup>8</sup>; 15 min after each extraction the blood was centrifuged in situ using a portable centrifuge. The samples were transported daily in a portable fridge (4-6°C) for processing at the Clinical Biochemistry laboratory of the Hospital Universitario Central de Asturias. Measurements were made of blood glucose fasting and after the OGTT (glucose-hexokinase enzyme method, Hitachi 747 analyzer, Roche Diagnostics, Mannheim, Germany). Measurements were also made of concentrations of total cholesterol, high-density lipoprotein cholesterol (HDL-C), triglycerides (colorimetric enzyme method, Hitachi 747 analyzer, Roche Diagnostics, Mannheim, Germany), low-density lipoprotein cholesterol (LDL-C) (Friedewald equation),<sup>9</sup> and glycohemoglobin (HbA<sub>1c</sub>) (high power liquid chromatography [HPLC], Jokoh HS-10 analyzer). The participants were notified by letter of the results of their OGTT, and those who had undiagnosed DM or glucose intolerance were advised to contact their primary care physician for follow-up and control.

### Follow-up and Identification of Fatal Events

In December 2004, prior to starting the second phase of the Asturias Study, the vital status was

verified of the whole cohort that participated in the first phase of the study (1998-1999). Data obtained from the health card provided by the central services of the Principality of Asturias Health Service were used to record all deaths and their dates. This information was verified by checking the medical registries of the whole cohort; 19 persons who had moved their residence away from Asturias before the start of the second phase were excluded from this re-evaluation, as their vital status could not be verified. In total, the registry included 1015 persons, with a follow-up time of six years (December 1998 to December 2004).

The cause of death was determined by systematic review of the clinical records, both at their health centers and in the corresponding referral hospital. If necessary, the cases were commented on in person with the primary care physician. Additional data from death certificates were also available for 93% of the cases. The cause of death was coded according to the IDC-10 of the World Health Organization.<sup>10</sup> Cardiovascular death was defined with codes I00-I99 ("diseases of the circulatory system") or R96 ("other sudden death, cause unknown"), because sudden death is generally of cardiovascular origin.<sup>11</sup> Death due to cancer was defined with the codes C00-D48 ("neoplasms").

### Statistical Study

All the statistical analyses were done with SPSS 12.0 (SPSS, Chicago, IL) and Epibasic 1.0 (University of Aarhus, Nordre Ringgade, Denmark). The reported *P* values are based on a 2-tailed test with a limit of statistical significance of  $P < .05$ .

Evaluation of the risk for death in the different categories of dysglycemia was done by classifying the persons included in the registry of deaths into four groups, according to the results of the OGTT during the first study (World Health Organization 1999 criteria)<sup>9</sup>: normoglycemia (fasting glycemia  $< 110$  mg/dL and 2 h post OGTT glycemia  $< 140$  mg/dL), prediabetes (fasting glycemia 110-126 mg/dL and/or 2 h post OGTT glycemia 140-200 mg/dL), undiagnosed DM (fasting glycemia  $\geq 126$  mg/dL and/or 2 h post OGTT glycemia  $\geq 200$  mg/dL), and diagnosed DM.

Comparisons between groups for quantitative variables were done with an analysis of variance (ANOVA). Multiple comparisons between pairs were adjusted with the Bonferroni test. The  $\chi^2$  test was used for comparison of proportions.

The person-years of follow-up were estimated for each group, as was the number of events (deaths), calculating the mortality rates for each 1000 inhabitant-years (95% confidence interval [CI]), which was adjusted for age and sex by the direct

method, using the age and sex structure of the normoglycemic group as the reference.

Cox regression analysis was used to analyze the accumulated impact curves and the corresponding relative risks (RR) of death, adjusted for age and sex and multivariable (adjusted for age, sex, BMI, systolic BP, diastolic BP, smoking, LDL-C and the prior presence of CVD) in the different groups.

### RESULTS

A total of 42 deaths were recorded during the follow-up period. The causes of death were: cardiovascular origin, 17 (40.5%); cancer, 19 (45.2%); other causes, 6 (14.3%).

Table 1 shows a comparison of the metabolic and cardiovascular risk parameters according to the different clinical categories of dysglycemia during the first phase of the study (1998-1999). The age increased with the category of dysglycemia, and was maximum in the group with diagnosed DM. The percentage of men was higher in all the groups as compared with the reference group that had normoglycemia. The cardiovascular risk profile worsened progressively from normoglycemia to prediabetes to DM in most of the study parameters, including systolic BP, diastolic BP, BMI, HDL-C, and triglycerides. Those persons with undiagnosed DM had similar figures for systolic BP, diastolic BP, HDL-C, and triglycerides to the persons with diagnosed DM. However, persons with diagnosed DM had similar, or even lower figures for total cholesterol and LDL-C to the general population without dysglycemia, values that were maximum in the group with prediabetes and undiagnosed DM. A possible explanation for this is the greater use of lipid-lowering drugs in persons with diagnosed DM. Likewise, there was a progressive increase across the groups for the figures of fasting glycemia, 2 h post OGTT glycemia and HbA<sub>1c</sub>. Of note was the fact that the mean levels of HbA<sub>1c</sub> in the group with undiagnosed DM were almost within the range of normality.

Around 10% of the persons with prediabetes or undiagnosed DM already had some symptom of CVD at the start of the study, which was only present in 2.1% of the general population with normoglycemia ( $P < .01$ ). In the group with diagnosed DM, 17.4% had CVD at the start of the study.

Only 39.1% of the diagnosed diabetic patients were taking treatment with hypertension drugs and 15.6% with lipid-lowering drugs.

Table 2 shows the number of deaths, the mortality rates for each 1000 inhabitant-years adjusted for age and sex and the risk of death adjusted for age and sex and multivariable in the different groups.

Both those with diagnosed DM and those who had undiagnosed DM had a risk of mortality almost

**TABLE 1. Initial Characteristics of the Persons Included in the Mortality Registry According to Category of Dysglycemia: 1999 WHO Classification**

	Normoglycemia	Prediabetes	Undiagnosed DM	Diagnosed DM	P (trend)
Patients, n	712	178	79	45	—
Age, y	50 (12.9) <sup>a</sup>	59 (11.6) <sup>b</sup>	61 (11.2) <sup>b</sup>	66.2 (8.3) <sup>c</sup>	<.001
Men, %	41.6 <sup>a</sup>	55.1 <sup>b</sup>	55.7 <sup>b</sup>	58.7 <sup>b</sup>	<.001
SBP, mm Hg	128.7 (19.4) <sup>a</sup>	143.6 (22.1) <sup>b</sup>	151.7 (21.6) <sup>c</sup>	156.2 (22.8) <sup>c</sup>	<.001
DBP, mm Hg	81.2 (12.7) <sup>a</sup>	88.4 (14.4) <sup>b</sup>	91.1 (13.3) <sup>b</sup>	91.8 (11.8) <sup>b</sup>	<.001
BMI	26.8 (4.3) <sup>a</sup>	29.3 (4.6) <sup>b</sup>	30.6 (4.9) <sup>b</sup>	28.6 (5.4) <sup>b</sup>	<.001
Total cholesterol, mg/dL	225.4 (41.5) <sup>a</sup>	238.5 (42.2) <sup>b</sup>	239.8 (38.8) <sup>b</sup>	221.9 (33.2) <sup>a</sup>	<.001
LDL-C, mg/dL	147.3 (37.4) <sup>a</sup>	156.9 (37) <sup>b</sup>	157.7 (33.7) <sup>b</sup>	140.6 (31.3) <sup>a</sup>	<.01
HDL-C, mg/dL	57.6 (14.3) <sup>a</sup>	54.4 (13.3) <sup>b</sup>	51.9 (14.5) <sup>b</sup>	51.8 (12.9) <sup>b</sup>	<.001
Triglycerides, mg/dL	106.2 (65.9) <sup>a</sup>	134.8 (68.8) <sup>b</sup>	154.2 (76.3) <sup>b</sup>	151 (79) <sup>b</sup>	<.001
HbA <sub>1c</sub> , %	4.7 (0.5) <sup>a</sup>	5 (0.5) <sup>b</sup>	5.6 (1) <sup>c</sup>	7.2 (1.8) <sup>d</sup>	<.001
Baseline glycemia, mg/dL	92 (7.8) <sup>a</sup>	105.1 (11.4) <sup>b</sup>	124 (23.6) <sup>c</sup>	169.1 (57.2) <sup>d</sup>	<.001
2 h post-OGTT glycemia, mg/dL	95.4 (22.3) <sup>a</sup>	149.5 (30.7) <sup>b</sup>	232.1 (63.2) <sup>c</sup>	—	<.001
Smokers, %	28.2	22.5	24.1	22.7	NS
On treatment for hypertension, %	9.4	25.3	27.8	39.1	<.001
On treatment for dyslipidemia, %	2.8	6.2	8.9	15.6	<.001
Prior CVD, %	2.1	9	10.1	17.4	<.001

BMI indicates body mass index; CVD, cardiovascular disease; DBP, diastolic blood pressure; DM, diabetes mellitus; HDL, high-density lipoprotein; LDL, low-density lipoprotein; SBP, systolic blood pressure.

The means labeled with the same letter are not statistically different with a  $P < .05$ .

**TABLE 2. All-Cause Mortality According to Category of Dysglycemia (1999 WHO Classification)**

	Normoglycemia	Prediabetes	Undiagnosed DM	Diagnosed DM
Patients, n	712	178	79	45
Deaths, n	14	10	10	8
Mortality/1000 person-years, <sup>a</sup> mean (95% CI)	3.3 (2-5.6)	4.9 (1.6-11.3)	10.8 (3.5-25.1)	13.8 (3.8-35.3)
Age and sex, RRA (95% CI)	1	1.5 (0.7-3.5)	2.8 (1.2-6.5) <sup>b</sup>	2.9 (1.2-7) <sup>b</sup>
Multivariate <sup>c</sup>	1	1.6 (0.7-4)	2.7 (1.1-6.7) <sup>b</sup>	2.5 (1-6.3) <sup>b</sup>

BMI indicates body mass index; CVD, cardiovascular disease; DBP, diastolic blood pressure; DM, diabetes mellitus; HDL, high-density lipoprotein; LDL, low-density lipoprotein; RRA, adjusted relative risk; SBP, systolic blood pressure.

<sup>a</sup>Mortality rates adjusted for age and sex.

<sup>b</sup>Statistically significant ( $P < .05$ ).

<sup>c</sup>The Cox multivariate regression model was adjusted for age, sex, BMI, SBP, DBP, smoking, LDL-C, and previous diagnosis of CVD.


3-fold greater than persons with normoglycemia in the analysis adjusted for age and sex; these levels remained just slightly lower after the multivariate analysis. The group with prediabetes also had an increase in mortality as compared with the control group, although the difference was not statistically significant. Using the American Diabetes Association 2003 criteria for the diagnosis of prediabetes showed the relative risk of death to be similar in the model adjusted for age and sex: RR=1.5 (0.7-3.3), and slightly lower in the multivariate model: RR=1.3 (0.5-3.0) as compared with the normoglycemic group. The risk of death in those with DM, both diagnosed and undiagnosed, versus those with normoglycemia was 2.9 (1.4-5.9) in the model adjusted for age and sex, and 2.6 (1.2-5.7) in the multivariate model.

Figure 1 shows the accumulated mortality curves for the four groups. The mortality curves in the groups with diagnosed DM and undiagnosed DM were practically the same.

The specific analysis for cardiovascular death showed the same trend, although this analysis was rejected because of the low number of cases available.

## DISCUSSION

The main finding of this study was that both persons with diagnosed DM and those with undiagnosed DM had a risk of death 2.5 to 3 times greater than those with normoglycemia, a risk that was similar in both groups. We also found an excess mortality in persons with prediabetes, though the difference was not statistically significant.



**Figure 1.** Accumulated all-cause mortality curves for the groups with normoglycemia, prediabetes, undiagnosed diabetes mellitus, and diagnosed diabetes mellitus (model adjusted for age, sex, body mass index, diastolic blood pressure, systolic blood pressure, smoking, low-density lipoprotein cholesterol, and presence of previous cardiovascular disease).

The data from this registry are alarming, as the different cross-sectional studies that have been carried out in Spain (see the “Introduction”) suggest that at least half those persons with DM remain undiagnosed whilst the disease remains asymptomatic, even over many years. Diagnosis during this preclinical phase is only possible by screening or incidentally. The fact that those persons with undiagnosed DM have a significantly higher risk of death than the control group with normoglycemia and a similar risk to those with diagnosed DM underlines the importance of detection in these persons.

Of note, though, is the fact that, unlike other diseases that require a screening test and a diagnostic test, for persons with DM the same test can be used for both screening and diagnosis of the disease, ie, the OGTT. Furthermore, this screening has the added advantage of being able to diagnose persons who have prediabetes, and who could thus benefit from possible preventive strategies.

Admittedly, it is true that no direct evidence exists that early detection and multifactorial treatment of persons with asymptomatic DM results in an improvement in prognosis.<sup>12</sup> However, it is also true that the lack of evidence for this is due to the absence of studies investigating this hypothesis and, for obvious ethical reasons, it is unlikely that any rigorously controlled studies will be carried out to provide any direct proof. What we do have, though, is the indirect evidence that aggressive multifactorial

treatment aimed not only at reducing the HbA<sub>1c</sub>, but also at lowering LDL-C, BP, and anti-platelet aggregation, produces effective reductions in vascular complications and the risk of death in persons with type 2 DM.<sup>13,14</sup> Thus, there appears to exist no reason for supposing that early detection and aggressive intensive therapy of asymptomatic persons should not produce similar results.

This intensive therapy was not optimal in our cohort of patients with diabetes, not even in the case of the group with diagnosed DM. As can be seen from Table 1, the mean values of the LDL, as well as the systolic BP and the diastolic BP and the percentage of persons who were receiving treatment with hypertension drugs and lipid-lowering drugs were all well below the optimal objectives for control in patients with type 2 DM according to current clinical practice guidelines.<sup>15</sup> It is possible that aggressive treatment of cardiovascular risk factors during the follow-up might have led to a lower risk of death. In the group of persons with undiagnosed DM, intensive treatment of cardiovascular risk factors was even lower; in fact, during the re-evaluation of the cohort, we found that a high number of these participants (around 40%) had not even told their primary care physicians about the results of the OGTT, and there was no record in their clinical histories. At the end of the follow-up period, the percentages of patients receiving hypertension drugs, lipid-lowering drugs and anti-platelet aggregators were 64%, 36%, and



44% in the group with diagnosed DM, as compared to 46.3%, 22%, and 7.3%, respectively, in the group with undiagnosed DM (data corresponding to 700 participants in the second phase of the Asturias Study, 2004-2005). This could account for the fact that the excess mortality in this group was similar to that of the group with diagnosed DM, even though their age, HbA<sub>1c</sub> levels and the initial presence of CVD were all lower. This study was not designed as an intervention study, but rather an observation study, and what we saw may reflect the true situation facing us: it is estimated that fewer than half of those persons who have a cardiovascular risk factor are aware of their diagnosis, that fewer than half of these are receiving non-specific treatment and, that fewer than half of those who are receiving treatment reach the therapeutic aims recommended by the clinical practice guidelines.<sup>16</sup>

These data have important public health implications. The impact of persons with undiagnosed hyperglycemia on CVD and overall risk of death in our populations is an undervalued problem, the dimensions of which could surpass any estimate. The Euro Heart Survey on Diabetes and the Heart, in which 4196 patients with coronary disease were recruited from 110 centers in 25 countries, produced convincing findings: 22% of patients seen for an acute coronary syndrome were found to have undiagnosed DM after undergoing an OGTT and 36% had impaired fasting glucose (IFG) and/or impaired glucose tolerance (IGT); normoglycemia is less common in these patients than dysglycemia. The overall proportion of persons with DM, both diagnosed and undiagnosed, was estimated to be around 45%.<sup>17</sup> Studies carried out in stroke patients have found similar results, with just a minority of patients showing normoglycemia after undergoing an OGTT.<sup>18</sup> The impact on overall mortality is unknown.

In Spain, the only previous reference to the evaluation of the risk of death in persons with undiagnosed DM and prediabetic dysglycemia is the Lejona Study. Ten years after the initial study, the vital status of the cohort was evaluated. An excess all-cause mortality was found in patients with diabetes that was almost twice that of persons with normal glucose tolerance, an excess risk that proved to be mainly due to those persons with undiagnosed DM.<sup>19</sup>

Outside Spain, we can find several studies. Shaw et al<sup>20</sup> analyzed longitudinal studies from the islands of Mauritius, Fiji, and Nauru, and found a risk for cardiovascular death in persons with undiagnosed DM that was 2-fold greater than in persons without DM only in those who had 2 h post-OGTT glycemic levels  $\geq 200$  mg/dL but not for persons with a fasting glucose of  $\geq 126$  mg/dL and 2 h post-OGTT  $< 200$  mg/dL.

A review of the second national NHANES survey in the United States, which included 3092 persons aged 30 to 74 years, found a relative risk of death of 2.26 (1.78-2.87) in persons with diagnosed DM, 1.76 (1.17-2.66) in persons with undiagnosed DM and 1.37 (1.05-1.79) in persons with IGT, compared with the control group with normoglycemia. The relative risks for cardiovascular death were similar.

The most conclusive data can be found in the DECODE study, which included 29 714 persons aged 30-89 years from 22 European cohorts. The study found an increase in the risk for all-cause mortality and cardiovascular mortality in persons with undiagnosed DM, especially for persons with 2 h post-OGTT glycemic levels  $\geq 200$  mg/dL, who had a relative risk of death that was similar to those persons with diagnosed DM.<sup>3</sup>

Our study has certain limitations: the main one is the low number of fatal events available for the analysis. The Asturias Study was designed as a cross-sectional study to estimate the prevalence of type 2 DM and IGT, as well as the proportion of persons with undiagnosed DM in the population of Asturias, and the sample size was sufficient for these estimations.<sup>4</sup> Estimation of the risk of death in the different groups would probably have required a larger cohort or longer follow-up period in order to obtain greater precision in the results. Additionally, participation in the initial field study was not complete (64% of the persons initially selected). This rate of participation might affect the representativity of the sample and the extrapolation to the general population of the data obtained. Nevertheless, in spite of these limitations, inherent to the study design itself, we consider that the study presents the important advantage of analyzing a representative sample of the general population from the whole province, including both rural and urban areas, with a complete initial evaluation by means of an OGTT, which gave us the unusual opportunity of being able to evaluate mortality in a number of persons with prediabetes and undiagnosed DM, a fact that was almost unprecedented in Spain. Despite the relatively low number of fatal events studied, the data are concordant with earlier studies, which reinforces our results.

## CONCLUSIONS

In summary, this study found a risk of mortality in persons with diabetes that was 2.5-3 times higher than in persons with normoglycemia, with a similar risk in both persons with diagnosed and undiagnosed diabetes. These results, which agree with those of previous studies, reflect the importance for public health of detection and intensive treatment of these persons.

## ACKNOWLEDGMENTS

The authors wish to express special thanks to all those Asturian persons who participated in the 2 studies, without whose collaboration it would have been impossible to carry out. Also, to the primary care teams of the Basic Health Areas selected. To Ismael Huerta, Shura Rozada, and Enrique Bustillo. To the Servicio de Bioquímica Clínica of the Hospital Universitario Central de Asturias and to the Endocrinology group of Asturias.

## REFERENCES

1. Goday A. Epidemiología de la diabetes y sus complicaciones no coronarias. *Rev Esp Cardiol*. 2002;55:657-70.
2. Valdés S, Rojo-Martínez G, Soriguer F. Evolución de la prevalencia de la diabetes tipo 2 en población adulta española. *Med Clin (Barc)*. 2007;129:352-5.
3. DECODE Study Group. Is the current definition for diabetes relevant to mortality risk from all causes and cardiovascular and noncardiovascular diseases? *Diabetes Care*. 2003;26:688-96.
4. Botas P, Delgado E, Castaño G, Díaz de Greñu C, Prieto J, Díaz-Cadorniga FJ. Prevalencia de diabetes mellitus e intolerancia a la glucosa en población entre 30 y 75 años en Asturias, España. *Rev Clin Esp*. 2002;202:423-915.
5. Botas P, Delgado E, Castaño G, Díaz de Greñu C, Prieto J, Díaz-Cadorniga FJ. Comparison of the diagnostic criteria for diabetes mellitus, WHO-1985, ADA-1997 and WHO-1999 in the adult population of Asturias (Spain). *Diabet Med*. 2003;20:904-8.
6. Valdés S, Botas P, Delgado E, Álvarez F, Díaz-Cadorniga F. Population-Based incidence of type 2 diabetes in Asturias, Spain. "The Asturias Study". *Diabetes Care*. 2007;30:2258-63.
7. Valdés S, Botas P, Delgado E, Álvarez F, Díaz-Cadorniga F. Does the new American Diabetes Association definition for impaired fasting glucose improve its ability to predict type 2 diabetes mellitus in Spanish persons? "The Asturias Study". *Metabolism*. 2008;57:399-403.
8. World Health Organization: Definition, Diagnosis and Classification of Diabetes Mellitus and its Complications: Report of a WHO Consultation. Part 1: Diagnosis and Classification of Diabetes Mellitus. Geneva: WHO; 1999.
9. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem*. 1972;18:499-502.
10. World Health Organization. International Statistical Classification of Diseases and Related Health Problems 10th Revision. Version for 2007 [citado 8 Oct 2008]. Disponible en: <http://www.who.int/classifications/apps/icd/icd10online/>
11. Kannel WB, Plehn JF, Cupples LA. Cardiac failure and sudden death in the Framingham Study. *Am Heart J*. 1988;115:869-75.
12. Engellau MM, Narayan KM, Herman WH. Screening for type 2 diabetes. *Diabetes Care*. 2000;23:1563-80.
13. Gaede P, Vedel P, Larsen N, Jensen GVH, Parving H-H, Pedersen O. Multifactorial intervention and cardiovascular disease in patients with type 2 diabetes. *N Engl J Med*. 2003;348:383-93.
14. Gaede P, Lund-Andersen H, Parving HH, Pedersen O. Effect of a multifactorial intervention on mortality in type 2 diabetes. *N Engl J Med*. 2008;358:580-91.
15. American Diabetes Association: Standards of medical care in diabetes—2008. *Diabetes Care*. 2008;31 Suppl 1:12-54.
16. Alfonso F, Segovia J, Heras M, Bermejo J. Prevención cardiovascular: ¿siempre demasiado tarde? *Rev Esp Cardiol*. 2008;61:291-8.
17. Bartnik M, Ryden L, Ferrari R, Malmberg K, Pyorala K, Simoons M, et al; on behalf of the Euro Heart Survey Investigators. The prevalence of abnormal glucose regulation in patients with coronary artery disease across Europe. *Eur Heart J*. 2004;25:1880-90.
18. Matz K, Keresztes K, Tatschl C, Nowotny M, Dachenhausenm A, Brainin M, et al. Disorders of glucose metabolism in acute stroke patients. *Diabetes Care*. 2006;29:792-7.
19. Vázquez JA, Gaztambide S, Soto-Pedre E. Estudio prospectivo a 10 años sobre la incidencia y factores de riesgo de diabetes mellitus tipo 2. *Med Clin (Barc)*. 2000;115:534-9.
20. Shaw JE, Hodge AM, De courten M, Chitson P, Zimmet PZ. Isolated post-challenge hyperglycaemia confirmed as a risk factor for mortality. *Diabetologia*. 1999;42:1050-4.
21. Saydah S, Loria CM, Eberhardt MS, Brancati FL. Subclinical states of glucose intolerance and risk of death in the U.S. *Diabetes Care*. 2001;24:447-53.