

Simulation Process

- Sim true pop from thornthall density
- calc $R2 \dots O2 / P2$ estimators ** N not factor into true SE*
- fit $dsm()$, calc striplet/boxlet
- draw from $dsm()$ to create new pop
- N-bet from each pop forms MC est

100x
200x

Vary # Detections

- Options:
 - ① change total N
 - ② change # of transects
 - ③ change σ or w


Vary density grid spacing

- proportional to survey area?
- one smaller than $\frac{10,000 \text{ m}^2}{5,000,000 \text{ m}^2} = \text{base}$
- one larger than
- Since density grid spacing dictates the simulated transect size, this tests that change too

Any Others?

Base Set-up

- 4 triangles w/ Fowler-like densities
- 100 population re-generations
- 200 MC reps
- Once w/ points, once w/ lines
- Once code is finalized, I can sort out how long this sim will take



Resulting SEs:

	R1	R2	S1	S2	O1	O2	P2	P3	Striplet	Boxlet	MC Est.	"True SE"
Points	NA	NA	NA	NA	NA	NA	#	#	NA	#	#	#
Lines	#	#	#	#	#	#	NA	NA	#	NA	#	#

Density #1



Same table for all four densities } probably better to do two tables, all for the points, then all for the lines