

Maximum Likelihood Estimation

Given a collection of data, what is the underlying distribution

→ Assuming a distribution, what are the likeliest parameters for that distribution.

Suppose we've conducted an experiment and our collected data is a list of values $x_1, x_2, x_3, \dots, x_n$.

Suppose there is a function f for the probability of these data to be measured. This function is parametrized by θ :

$$\text{Prob}(\text{data}) = f(x_1, x_2, x_3, \dots, x_n | \theta) \leftarrow \begin{array}{l} \text{joint prob of all} \\ \text{the data simultaneously} \\ \text{given a (set of)} \\ \text{parameters } \theta. \end{array}$$

Now let's make things easier: Assume each data point can occur independently from all others and each follows the same distribution (independent, identically distributed or "iid").

$$\begin{aligned} \hookrightarrow f(x_1, x_2, \dots, x_n | \theta) &= f(x_1 | \theta) \cdot f(x_2 | \theta) \cdot \dots \cdot f(x_n | \theta) \\ &= \prod_{i=1}^n f(x_i | \theta) \end{aligned}$$

OK, our goal is to find θ , we're given the data. We need to flip this around: what is the likelihood of θ , given the data? No problem, flip!

$$\mathcal{L}(\theta | x_1, x_2, \dots, x_n) = f(x_1, x_2, \dots, x_n | \theta) = \prod_{i=1}^n f(x_i | \theta)$$

(whoa are you allowed to just do that?)

Moving forward, we want to find the θ that maximizes $\mathcal{L}(\theta | \text{data})$. (called the maximum likelihood estimator, $\hat{\theta}$)

Often it's easier to maximize the log-likelihood which is maximized at the same place: $\ell(\theta | \text{data}) = \sum_{i=1}^n \ln f(x_i | \theta)$ □

MLE cont

Sometimes we can find the $\hat{\theta}$ easily with calculus, other times we need a numerical method.

Using calculus, compute the derivative $\frac{\partial \ell}{\partial \theta}$, set equal to 0, solve.

Let's do a specific example:

Poisson distribution:

Prob for a certain number of events to occur if the average rate of events is known and the prob. for the next event to occur does not depend on the time since the previous event.

$$P(X=k; \lambda) = P(k; \lambda) = \frac{\lambda^k e^{-\lambda}}{k!}$$

Let's assume the x_i 's are drawn from a poisson. Then we need to find $\hat{\lambda}$:

$$f(x_i | \theta) \Rightarrow f(x_i | \lambda) = \frac{\lambda^{x_i} e^{-\lambda}}{x_i!}$$

Compute the (log)-likelihood:

$$\mathcal{L}(\theta | \text{data}) = \prod_{i=1}^n f(x_i | \lambda) = \prod_{i=1}^n \frac{\lambda^{x_i} e^{-\lambda}}{x_i!} = e^{-n\lambda} \prod_{i=1}^n \frac{\lambda^{x_i}}{x_i!}$$

$$\begin{aligned} \ell(\theta | \text{data}) &= \ln \mathcal{L}(\theta | \text{data}) = \ln e^{-n\lambda} + \sum_{i=1}^n \ln \lambda^{x_i} - \sum_{i=1}^n \ln(x_i!) \\ &= -n\lambda + \sum_{i=1}^n x_i \ln \lambda - () \\ &= -n\lambda + \ln \lambda \sum_{i=1}^n x_i - () \end{aligned}$$

Now we find an equation for the maximum as a function of λ , and solve it for $\hat{\lambda}$:

$$\frac{\partial \ell}{\partial \lambda} = \frac{\partial}{\partial \lambda} \left(-n\lambda + \ln \lambda \sum x_i - () \right) = -n + \underbrace{\ln \lambda}_{0} \underbrace{\frac{\partial}{\partial \lambda} \sum x_i}_{0} + \sum x_i \cdot \frac{1}{\lambda} - 0$$

↓ 2

Solving this for $\hat{\lambda}$

$$-n + \frac{1}{\hat{\lambda}} \sum_{i=1}^n x_i = 0 \Rightarrow \hat{\lambda} = \frac{1}{n} \sum_{i=1}^n x_i$$

The MLE $\hat{\lambda}$ for poisson is the average of the data.

So if we are given a bunch of integer x -data, we can compute the mean, and draw $P(X=k, \hat{\lambda})$ on top of the histogram and see if it's close.
