8. How many different three-letter initials with none of the letters repeated can people have?

With no repeated letters there will be 26 \* 25 \* 24 = 15600 different three letter initials.

12. How many bit strings are there of length six or less?

There will be 6 separate cases where each case is  $2^n$ . The total will be  $\sum_{n=0}^{6} 2^n = 127$ 

16. How many strings are there of four lowercase letters that have the letter x in the term?

If there were no X's then the highest number would be 26 \* 26 \* 26 \* 26 \* 26 \* 26 \* 26. Since there needs to be an at least one X there will be 4 cases:

Case 1 (one x): With one X there are 4 combinations of where the letter can be placed in the string. That makes the total 4\*25\*25\*25=62500

Case 2 (two x): There will now be 6 combinations of placements for the letter in the string but it also takes up two positions. That makes the total 6 \* 25 \* 25 = 3750

Case 3 (three x): There are 4 combinations of placements for three x's. That makes the total 2 \* 25 = 100

Case 4 (four x): There is only one string with 4 x's.

Adding all of the cases up gives a total of 66351 four letter strings.

- 24. How many strings of four decimal digits
- a) do not contain the same digit twice?
- b) end with an even digit?
- c) have exactly three digits that are 9s?

A: Since the string can't contain the same digit twice it can have 10 choices for the first digit, 9 for the second, 8 for the third, and 7 for the last. Giving a total of 10 \* 9 \* 8 \* 7 = 5040

B: Half of the strings will end in an even digit. The total of all four digit strings is  $10^4$  and the even strings will be  $\frac{10^4}{2} = 5000$ 

C: There are 4 combinations of three digits (9) in a 4 digit string. That makes the total 4\*9=36

26. How many license plates can be made using either three digits followed by three letters or three letters followed by three digits?

There are 10 \* 10 \* 10 number combinations and 26 \* 26 \* 26 letter combinations. The total for one type of plate is  $10^3 * 26^3$ . The other plate will give the same number so the overall total is  $2 * (10^3 * 26^3) = 35152000$ 

42. How many bit strings of length seven either begin with two 0s or end with three 1s?

There will be  $2^5$  strings that begin with two 0s and  $2^4$  strings that end with three 1s. The overlap between the two sets is  $2^2$  so the total is  $(2^5 + 2^4) - 2^2 = 44$ .

46. Every student in a discrete mathematics class is either a computer science or a mathematics major or is a joint major in these two subjects. How many students are in the class if there are 38 computer science majors(including joint majors), 23 mathematics majors(including joint majors), and 7 joint majors?

There are 38compsci + 23math - 7joint = 54 students in the class.

60. Use mathematical induction to prove the product rule for m tasks from the product rule for two tasks.

P(m) is the product rule for m tasks.

Basis step:

m = 2, P(2) is true since there are  $n_1$  ways to do the first task and  $n_2$  ways to do the second one. The total being  $n_1n_2$ .

Inductive Step:

Prove P(k) is true for  $k \geq 2$ .

For a k+1 tasks we have  $T_1T_2 \cdot T_{k+1}$ . Each of which can be done  $n_1n_2 \cdot n_{k+1}$  ways. That means that for the next k+1 task we will have  $(n_1n_2 \cdot n_k) * n_{k+1}$  ways which proves P(k+1).