


Stellar mass and sSFR · CIGALE 15-band · full range


4,989,187 galaxies; same sample in each panel, weighted by current environment probabilities
Each class is normalized by its total probability weight; shared density scale.



Provisional LoA environments; real-DESI coverage unverified. High- z population mismatch; trends are not physical evolution.
CIGALE SFRs may be affected by Tractor modelling, especially at low z . Descriptive probability-weighted relationships.

Stellar mass and sSFR · CIGALE 15-band · lowz full sky


2,327,331 galaxies; same sample in each panel, weighted by current environment probabilities
Each class is normalized by its total probability weight; shared density scale.



Provisional LoA environments; real-DESI coverage unverified. High- z population mismatch; trends are not physical evolution.
CIGALE SFRs may be affected by Tractor modelling, especially at low z . Descriptive probability-weighted relationships.

Stellar mass and sSFR · CIGALE 5-band · lowz full sky




2,327,332 galaxies; same sample in each panel, weighted by current environment probabilities
Each class is normalized by its total probability weight; shared density scale.



Provisional LoA environments; real-DESI coverage unverified. High-z population mismatch; trends are not physical evolution.
CIGALE SFRs may be affected by Tractor modelling, especially at low z . Descriptive probability-weighted relationships.

Galaxy properties versus environment · full range




CG15 and CG5 use identical galaxies and control cells. Mass: common redshift mix.
sSFR / low-sSFR: common CG15-mass and redshift mix. Error bars: 16–84% sky-block bootstrap.



Provisional Lo α environments; real-DESI coverage unverified. High- z population mismatch; trends are not physical evolution.
CIGALE SFRs may be affected by Tractor modelling, especially at low z . Descriptive probability-weighted relationships.

Galaxy properties versus environment · lowz full sky



CG15 and CG5 use identical galaxies and control cells. Mass: common redshift mix.
sSFR / low-sSFR: common CG15-mass and redshift mix. Error bars: 16–84% sky-block bootstrap.



Provisional Lo α environments; real-DESI coverage unverified. High- z population mismatch; trends are not physical evolution.
CIGALE SFRs may be affected by Tractor modelling, especially at low z . Descriptive probability-weighted relationships.

Fixed stellar mass · CG15 · $0.20 \leq z < 0.30$, full sky



Each mass bin shares a redshift distribution across environments; shading: 16-84% spatial bootstrap.



Provisional LoA environments; real-DESI coverage unverified. High- z population mismatch; trends are not physical evolution.
CIGALE SFRs may be affected by Tractor modelling, especially at low z . Descriptive probability-weighted relationships.

Environment contrasts by redshift · selection-sensitive diagnostics



Knot minus void, with common CG15-mass / redshift controls. Same galaxies for CG15 and CG5.
These are not measurements of physical evolution; model-transfer and CIGALE systematics are not in the intervals.



Provisional Lo_a environments; real-DESI coverage unverified. High-z population mismatch; trends are not physical evolution.
CIGALE SFRs may be affected by Tractor modelling, especially at low z. Descriptive probability-weighted relationships.

CIGALE matching and property availability

Denominator: all supported VAC galaxies in each redshift bin, weighted by environment probability.



Provisional Lo_a environments; real-DESI coverage unverified. High-*z* population mismatch; trends are not physical evolution.
CIGALE SFRs may be affected by Tractor modelling, especially at low *z*. Descriptive probability-weighted relationships.