Loa colour–mass diagrams by inferred environment
Observed colour · original selections · Highest-probability class

Void · 0.15 ≤ z < 0.20
N = 101,865
log M★ ≥ 10.60
Outside axes: 0.0

Wall · 0.15 ≤ z < 0.20
N = 229,478
log M★ ≥ 10.60
Outside axes: 1.0

Filament · 0.15 ≤ z < 0.20
N = 171,474
log M★ ≥ 10.60
Outside axes: 0.0

Cluster · 0.15 ≤ z < 0.20
N = 31,253
log M★ ≥ 10.60
Outside axes: 0.0

Void · 0.20 ≤ z < 0.22
N = 44,019
log M★ ≥ 10.69
Outside axes: 0.0

Wall · 0.20 ≤ z < 0.22
N = 99,223
log M★ ≥ 10.69
Outside axes: 1.0

Filament · 0.20 ≤ z < 0.22
N = 76,222
log M★ ≥ 10.69
Outside axes: 0.0

Cluster · 0.20 ≤ z < 0.22
N = 12,560
log M★ ≥ 10.69
Outside axes: 0.0

Void · 0.20 ≤ z < 0.30
N = 104,151
log M★ ≥ 10.96
Outside axes: 2.0

Wall · 0.20 ≤ z < 0.30
N = 262,655
log M★ ≥ 10.96
Outside axes: 3.0

Filament · 0.20 ≤ z < 0.30
N = 210,268
log M★ ≥ 10.96
Outside axes: 0.0

Cluster · 0.20 ≤ z < 0.30
N = 34,123
log M★ ≥ 10.96
Outside axes: 2.0

Observed g − r [mag]

log₁₀ stellar mass / M⊙ (CIGALE)

Count / probability weight per bin (log scale)

Horizontal cuts: 0.75 and 1.5. Shared axes and count scale. Raw distributions; no mass/redshift matching or volume normalization.
Environment predictions are held fixed. Rest-frame colours retain internal attenuation; these are not dust-free or quenching labels.