Loa colour–mass diagrams by inferred environment
Observed colour · original selections · Probability-weighted inferred classes

Void · 0.15 ≤ z < 0.20
Σ p = 106,223
log M★ ≥ 10.60
Outside axes: 0.2

Wall · 0.15 ≤ z < 0.20
Σ p = 220,417
log M★ ≥ 10.60
Outside axes: 0.8

Filament · 0.15 ≤ z < 0.20
Σ p = 165,179
log M★ ≥ 10.60
Outside axes: 0.0

Cluster · 0.15 ≤ z < 0.20
Σ p = 42,251
log M★ ≥ 10.60
Outside axes: 0.0

Void · 0.20 ≤ z < 0.22
Σ p = 45,874
log M★ ≥ 10.69
Outside axes: 0.0

Wall · 0.20 ≤ z < 0.22
Σ p = 95,532
log M★ ≥ 10.69
Outside axes: 1.0

Filament · 0.20 ≤ z < 0.22
Σ p = 72,822
log M★ ≥ 10.69
Outside axes: 0.0

Cluster · 0.20 ≤ z < 0.22
Σ p = 17,797
log M★ ≥ 10.69
Outside axes: 0.0

Void · 0.20 ≤ z < 0.30
Σ p = 109,538
log M★ ≥ 10.96
Outside axes: 1.5

Wall · 0.20 ≤ z < 0.30
Σ p = 251,421
log M★ ≥ 10.96
Outside axes: 2.9

Filament · 0.20 ≤ z < 0.30
Σ p = 200,220
log M★ ≥ 10.96
Outside axes: 1.2

Cluster · 0.20 ≤ z < 0.30
Σ p = 50,019
log M★ ≥ 10.96
Outside axes: 1.3

Observed g − r [mag]
log₁₀ stellar mass / M⊙ (CIGALE)
Count / probability weight per bin (log scale)

Horizontal cuts: 0.75 and 1.5. Shared axes and count scale. Raw distributions; no mass/redshift matching or volume normalization.
Environment predictions are held fixed. Rest-frame colours retain internal attenuation; these are not dust-free or quenching labels.