Loa colour–mass diagrams by inferred environment
Rest-frame SDSS colour at zref=0.1 · audited selections · Highest-probability class

Void · 0.15 ≤ z < 0.20
N = 89,515
log M★ ≥ 10.65
Outside axes: 0.0

Wall · 0.15 ≤ z < 0.20
N = 204,244
log M★ ≥ 10.65
Outside axes: 0.0

Filament · 0.15 ≤ z < 0.20
N = 153,682
log M★ ≥ 10.65
Outside axes: 0.0

Cluster · 0.15 ≤ z < 0.20
N = 28,120
log M★ ≥ 10.65
Outside axes: 0.0

Void · 0.20 ≤ z < 0.22
N = 38,185
log M★ ≥ 10.74
Outside axes: 0.0

Wall · 0.20 ≤ z < 0.22
N = 87,065
log M★ ≥ 10.74
Outside axes: 0.0

Filament · 0.20 ≤ z < 0.22
N = 67,441
log M★ ≥ 10.74
Outside axes: 0.0

Cluster · 0.20 ≤ z < 0.22
N = 11,094
log M★ ≥ 10.74
Outside axes: 0.0

Void · 0.20 ≤ z < 0.30
N = 95,317
log M★ ≥ 10.98
Outside axes: 2.0

Wall · 0.20 ≤ z < 0.30
N = 242,144
log M★ ≥ 10.98
Outside axes: 1.0

Filament · 0.20 ≤ z < 0.30
N = 195,318
log M★ ≥ 10.98
Outside axes: 0.0

Cluster · 0.20 ≤ z < 0.30
N = 31,686
log M★ ≥ 10.98
Outside axes: 1.0

Rest-frame $^{0.1}(g - r)$ [mag]

log₁₀ stellar mass / M⊙ (CIGALE)

Count / probability weight per bin (log scale)

Horizontal cuts: 0.75 and 1.5. Shared axes and count scale. Raw distributions; no mass/redshift matching or volume normalization.
Environment predictions are held fixed. Rest-frame colours retain internal attenuation; these are not dust-free or quenching labels.