Loa colour–mass diagrams by inferred environment
Rest-frame SDSS colour at zref=0.1 · audited selections · Probability-weighted inferred classes

Void · 0.15 ≤ z < 0.20
Σ p = 93,492
log M★ ≥ 10.65
Outside axes: 0.0

Wall · 0.15 ≤ z < 0.20
Σ p = 196,174
log M★ ≥ 10.65
Outside axes: 0.0

Filament · 0.15 ≤ z < 0.20
Σ p = 147,939
log M★ ≥ 10.65
Outside axes: 0.0

Cluster · 0.15 ≤ z < 0.20
Σ p = 37,956
log M★ ≥ 10.65
Outside axes: 0.0

Void · 0.20 ≤ z < 0.22
Σ p = 39,831
log M★ ≥ 10.74
Outside axes: 0.0

Wall · 0.20 ≤ z < 0.22
Σ p = 83,827
log M★ ≥ 10.74
Outside axes: 0.0

Filament · 0.20 ≤ z < 0.22
Σ p = 64,373
log M★ ≥ 10.74
Outside axes: 0.0

Cluster · 0.20 ≤ z < 0.22
Σ p = 15,754
log M★ ≥ 10.74
Outside axes: 0.0

Void · 0.20 ≤ z < 0.30
Σ p = 100,291
log M★ ≥ 10.98
Outside axes: 1.5

Wall · 0.20 ≤ z < 0.30
Σ p = 231,867
log M★ ≥ 10.98
Outside axes: 1.1

Filament · 0.20 ≤ z < 0.30
Σ p = 185,821
log M★ ≥ 10.98
Outside axes: 0.8

Cluster · 0.20 ≤ z < 0.30
Σ p = 46,485
log M★ ≥ 10.98
Outside axes: 0.5

Rest-frame $^{0.1}(g - r)$ [mag]
log₁₀ stellar mass / M⊙ (CIGALE)
Count / probability weight per bin (log scale)

Horizontal cuts: 0.75 and 1.5. Shared axes and count scale. Raw distributions; no mass/redshift matching or volume normalization.
Environment predictions are held fixed. Rest-frame colours retain internal attenuation; these are not dust-free or quenching labels.