

# MNRAS L<sup>A</sup>T<sub>E</sub>X 2<sub>ε</sub> template – title goes here

Keith T. Smith,<sup>1\*</sup> A. N. Other,<sup>2</sup> Third Author<sup>2,3</sup> and Fourth Author<sup>3</sup>

<sup>1</sup>*Royal Astronomical Society, Burlington House, Piccadilly, London W1J 0BQ, UK*

<sup>2</sup>*Department, Institution, Street Address, City Postal Code, Country*

<sup>3</sup>*Another Department, Different Institution, Street Address, City Postal Code, Country*

Accepted XXX. Received YYY; in original form ZZZ

## ABSTRACT

This is a simple template for authors to write new MNRAS papers. The abstract should briefly describe the aims, methods, and main results of the paper. It should be a single paragraph not more than 250 words (200 words for Letters). No references should appear in the abstract.

**Key words:** keyword1 – keyword2 – keyword3

## 1 INTRODUCTION

This is a simple template for authors to write new MNRAS papers. See `mnras_sample.tex` for a more complex example, and `mnras_guide.tex` for a full user guide.

?

## 2 ILLUSTRIS SIMULATION

Illustris-1, also known simply as Illustris, is a highly resolved cosmological simulation. It reproduces large-scale statistical features of the Universe, such as the galaxy population of massive clusters, as well as small-scale properties such as the morphology of galaxies and detailed values for their stellar and gas content.

This was achieved by following the evolution of  $2 \times 1820^3$  elements with dark mass resolution of  $m_{\text{DM}} = 6.26 \cdot 10^6 M_{\odot}$  and initial baryonic mass resolution of  $m_{\text{b}} = 1.26 \cdot 10^6 M_{\odot}$  from a glass-like configuration in a periodic box of 106.5Mpc. The  $\Lambda$ CDM cosmology of this run follows:  $\Omega_{\Lambda} = 0.7274$ ,  $\Omega_{\text{m}} = 0.2726$ ,  $\Omega_{\text{b}} = 0.0456$ ,  $\sigma_8 = 0.0809$ ,  $n_s = 0.963$  &  $H_0 = 70.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$  which is consistent with the (last) Anisotropy Probe (WMAP)-9 ?. Illustris has a constant spatial resolution of 1.4kpc for DM particles in comoving units, and for baryonic particles it has the same spatial resolution of DM for  $z \geq 1$ , which is later modified to 0.7kpc in physical units for the rest of the simulation.

The evolution in time was performed with the hydrodynamic code AREPO ?, which combines a moving Voronoi tessellation with the finite volume approach. Included in the evolution algorithm, there are galaxy formation models

which account for the evolution of stars and SMBHs. Specifically, the physics followed by this model includes energetic feedback from supermassive black holes and supernovae, as well as stellar evolution and chemical enrichment. This level of detail in Illustris is advantageous for the analysis of the effect of the environment in barionic properties of galaxies. This level of resolution is hard to find among other simulations, i.e. post-processed runs with semi-analytical models which do not directly simulate baryons.(Expand) Describe Arepo more

The output of this simultaion consists of 136 snapshots. 61 of them were taken at  $z < 3$  spaced with a cosmological scale factor  $\Delta a \approx 0.02$ . The remaining 75 were taken at  $z > 3$  with spacing  $\Delta a \approx 0.01$ . Each snapshot was post-processed with a modified version of FOF ? to identify DM haloes with more than 32 particles using a linking length of 0.2 times the mean particle separation. Each output group from the 7,713,601 post-processed halos (FOF) is analyzed with the SUBFIND algorithm ? to generate, at  $z=0$ , and 4,366,546 (sub)halo catalogues with their respective characterization properties.

(HERE TALK ABOUT ILLUSTRIS-2,3 FOR RESOLUTION PURPOSES)

## 3 THE METHOD

The principal purpose of this work is to verify if the results obtained observationally by Jones et al. ? are supported by a similar analysis in the cosmological simulation Illustris. Specifically, we focus on the (small-scale) environment definition of the 3<sup>rd</sup> nearest neighbor, which we want to adapt as closely as possible to our computational application.

\* E-mail: mn@ras.org.uk (KTS)

### 3.1 The 3<sup>rd</sup> nearest neighbor definition of environment

Jones et al. ? analyses the dependence of the Schechter parameters on the cosmological environment. To obtain a quantification of said cosmological environment, they calculated for each galaxy in the studied survey the projected distance in the sky to the third nearest neighbor, filtrating candidate neighbors by their radial velocity relative to the reference galaxy. According to Hubble's law, this effectively restricts the candidate galaxies to a smaller radial range, reducing apparent closer galaxies arising from the projection method. The radial relative velocity cut was chosen to be  $450\text{km/s}$ . In addition to this, a photometrics pre-selection was performed, selecting galaxies whose r-filter magnitude was above  $-19$  (VERIFY).

Once quantified the environment with the distance to the 3rd nearest neighbor definition they divided the group in four quartiles according this quantity. For each quartile, the respective HIMF was calculated, and with this, a Schechter fit was obtained. The analysis was specifically performed over the Schechter parameters of the Knee-mass and the faint-end slope. (Some observational adjustments for the Schechter functions were performed)

As we want to replicate as closely as possible the method effectuated by Jones et. al ?, we use the same specifications to calculate the environment. We perform the analysis of the Schechter functions to gas, DM, stellar and BH mass functions. However, we do not perform it directly on HIMFs because this would require some extense calculations. We take the gas mass function as a first approximation. As Illustris is a cosmological simulation, it is subject to resolution biases. The most important bias in this study is related to the minimum mass of discrete elements in the simulation. Each mass considered here is subject directly or indirectly this resolution restriction, and the effect it causes is specially evident in the produced mass functions. We used Illustris-2,3, which are less resolute copies of Illustris-1, to show this effect on the mass functions resolution effects. Taking this into account, mass functions are heavily affected for masses below a certain cut and for this reason, only galaxies above said cut are considered in this study. This mass cut has no aparent systematical way to be calculated and therefore, we performed the cut having into account the performance of the respective Schechter fit.

### 3.2 The cosmic web method

To complement and verify the analysis performed with the third nearest neighbor method, we chose a complementary method of completely different nature. The chosen method was developed by Forero et al. ?. It uses the Hessian 1 of the gravitational potential obtained from the DM density field to classify galaxies into four morphological environments.

**Table 1.** This is an example table. Captions appear above each table. Remember to define the quantities, symbols and units used.

A	B	C	D
1	2	3	4
2	4	6	8
3	5	7	9

Specifically, we diagonalize the Hessian matrix obtaining three eigenvalues. These eigenvalues determine the concavity of the gravitational potential on the principal axes, at each point. We use these eigenvalues to interpret the tendence of mass to cluster with reference with some given threshold  $\lambda$ . Eigenvalues bigger than  $\lambda$  are interpreted as a local tendence of mass to expand? in the respective eigenaxes, and viceversa. With this characterization, clusters, sheets, fillaments and voids can be defined with the number of axes above  $\lambda$ . Forero et. al. found in ? that the critical value that better reproduces the correct morphological pulations was  $\lambda = 0.2$ .

With these specifications, we could obtain mass functions for galaxies belonging to clusters, fillaments and sheets. The amount of galaxies belonging to voids was strongly affected by the mentioned mass cuts and could not produce reasonable mass functions to analyze.

## 4 RESULTS

### 4.1 Preliminaries

### 4.2 Mass functions comparison

Figures and tables should be placed at logical positions in the text. Don't worry about the exact layout, which will be handled by the publishers.

Figures are referred to as e.g. Fig. ??, and tables as e.g. Table 1.

## 5 CONCLUSIONS

The last numbered section should briefly summarise what has been done, and describe the final conclusions which the authors draw from their work.

## ACKNOWLEDGEMENTS


The Acknowledgements section is not numbered. Here you can thank helpful colleagues, acknowledge funding agencies, telescopes and facilities used etc. Try to keep it short.

## APPENDIX A: SOME EXTRA MATERIAL


If you want to present additional material which would interrupt the flow of the main paper, it can be placed in an Appendix which appears after the list of references.

This paper has been typeset from a  $\text{\LaTeX}$  file prepared by the author.


$$E = mc^2 \quad (1)$$




**Figure 1. Upper side:** Schechter fits of each gas mass function for each environmental quartile (tendence) using the 3rd nearest neighbor definition. **Down side:** Knee-mass  $m_*$  and the faint end slope  $\alpha$  values obtained from the Schechter fits for each curve shown above.




**Figure 2. Upper side:** Schechter fits of each gas mass function for each morphological environment characterized by the T-Web algorithm proposed by Forero et al. **Down side:** Knee-mass  $m_*$  and the faint end slope  $\alpha$  values obtained from the Schechter fits for each curve shown above.




**Figure 3.** Similar to figure 1 using DM.




**Figure 4.** Similar to figure 2 using DM.




**Figure 5.** Similar to figure 1 using stellar mass



**Figure 6.** Similar to figure 2 using stellar mass



**Figure 7.** Similar to figure 1 using BHs mass.



**Figure 8.** Similar to figure 2 using BHs mass.