

Environmental dependence of HI-Mass functions in cosmological simulations

Jesus Prada

Advisor: PhD Jaime Forero-Romero
Universidad de los Andes, Physics Department

In collaboration with:

PhD Michael Jones - Cornell University
PhD Martha Haynes - Cornell University


October, 2016

Outline

- 1 Introduction
 - Mass functions and Press-Schechter formalism
 - Environment
 - 3rd Nearest neighbour method
 - T-Web structures
- 2 Motivation
- 3 Our Work
 - Illustris run
 - Results
 - Conclusions
 - Further Work
- 4 References


- Histogram of masses of galaxies and halos
- Logarithmic scales are needed for better interpretation
- Press-Schechter adjusts surprisingly well to observations

$$n(M) = \phi_* \left(\frac{M}{M_*} \right)^{\alpha+1} e^{\left(\frac{M}{M_*} \right)}$$




- Related to the mass density
- Useful to verify its effect on the physics of galaxy formation
- Great variety of methods to quantify environment
- Nearest neighbour, control volume [2].
- 3rd Nearest neighbour (Observational) [1]
- T-Web structures (Theoretical) Forero-Romero et al. [3]


- Projected distance on sky.
- Velocity cut is needed.
- Distance to 3rd (not 1st) neighbour is required to avoid noise.



- Hessian of gravitational potential.
- 3 eigenvalues can identify 4 structures
- Clusters, sheets, filaments and voids.




Forero-Romero et al. [3]



Forero-Romero et al. [1]


- HIPASS and ALFALFA big surveys allow the study of HI-Mass functions.
- The effect of environment on the physics of galaxy formation.
- Mass tendency to cluster.
- Voids: slow evolution. Sample HIMF at earlier times.
- Studies are so recent and sometimes contradict each others (Springob, Haynes & Giovanelli) [4] and Stierwalt et al. [5].



- Illustris run
- $106.5 Mpc$
- 1820^3 particles
- $6.3 \cdot 10^6 M_{\odot}$ DM resolution
- $1.3 \cdot 10^6 M_{\odot}$ Gas resolution
- Detailed galaxy characteristics




<http://www.illustris-project.org/media/>



- Distance to 3rd nearest neighbour for galaxies with gas.
- Divided group in quartiles and obtained Schechter fits.





- Divided galaxies into Cluster, Sheet, filament and void groups.
- Obtained Schechter fits.





- We obtained a clear general tendency for the knee-mass and the faint-end slope in both methods.
- For high density environments from T-web, both tendencies becomes more diffuse.
- For low density neighbouring environments, the faint-end slope relation is diffuse.

- The knee-mass increases with neighbouring density in a range from 10.00 ± 0.03 to 10.62 ± 0.02 .
- For Cosmic web environment the range is from 9.90 ± 0.03 to 10.41 ± 0.04 .
- Faint-end ranges are $[-0.80 \pm 0.04, -1.05 \pm 0.02]$ and $[-0.76 \pm 0.05, 1.04 \pm 0.03]$.

- Detailed analysis of effects for big-scale environments using both methods.
- Relation between of different mass functions (DM-SMBH-gas).
- Incorporate a reasonable approximation for HI mass in terms of gas mass.

Referencias I

- [1] M. G. Jones, E. Papastergis, M. P. Haynes, and R. Giovanelli.
Environmental dependence of the H I mass function in the ALFALFA 70% catalogue.
MNRAS, 457:4393–4405, April 2016.
- [2] S. I. Muldrew, D. J. Croton, R. A. Skibba, F. R. Pearce, H. B. Ann, I. K. Baldry, S. Brough, Y.-Y. Choi, C. J. Conselice, N. B. Cowan, A. Gallazzi, M. E. Gray, R. Grützbauch, I.-H. Li, C. Park, S. V. Pilipenko, B. J. Podgorzec, A. S. G. Robotham, D. J. Wilman, X. Yang, Y. Zhang, and S. Zibetti.
Measures of galaxy environment - I. What is 'environment'?
MNRAS, 419:2670–2682, January 2012.
- [3] J. E. Forero-Romero, Y. Hoffman, S. Gottlöber, A. Klypin, and G. Yepes.
A dynamical classification of the cosmic web.
MNRAS, 396:1815–1824, July 2009.
- [4] C. M. Springob, M. P. Haynes, and R. Giovanelli.
Morphology, Environment, and the H I Mass Function.
APJ, 621:215–226, March 2005.
- [5] S. Stierwalt, M. P. Haynes, R. Giovanelli, B. R. Kent, A. M. Martin, A. Saintonge, I. D. Karachentsev, and V. E. Karachentseva.
The Arecibo Legacy Fast Alfa Survey. IX. The Leo Region H I Catalog, Group Membership, and the H I Mass Function for the Leo I Group.
AJ, 138:338–361, August 2009.