

Dark matter halo shapes at $z = 0$ in the Auriga simulations

Jesus Prada,^{1*} Jaime E. Forero-Romero,¹ Volker Springel²

¹*Departamento de Física, Universidad de los Andes, Cra. 1 No. 18A-10, Edificio 1p, Bogotá, Colombia*

²*Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, D-85741 Garching, Germany*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We present shape measurements of Milky Way sized dark matter halos at redshift $z = 0$ in a suite of 30 zoom simulations from the Auriga project. We compare the results in full magnetohydrodynamics against dark matter only simulations. We find a strong influence of baryons in making dark matter halos rounder at all radii compared to its dark matter only counterparts. At distances $\lesssim 30$ kpc, rounder dark matter distributions correlate with extended massive stellar discs and low core gas densities. We measure the alignment between the halo and the stellar disc at different radii and find almost perfect alignment at $0.25R_{200} \sim 56$ kpc. In some cases the alignment significantly changes as a function of radius implying that the halo shape twists; this effect correlates with extended stellar discs and low gas densities and is almost absent in the dark matter only simulations. In a comparison against observational constraints we find that 20% of halos in our sample are consistent with observational results derived from the Pal 5 stream that favor an almost spherical shape. Including baryons is a required element to achieve this level of agreement. In contrast, none of the simulations (dark matter only nor with baryons) match the constraints derived from the Sagittarius stream favoring an oblate dark matter halo.

Key words: galaxies: evolution — galaxies: formation — galaxies: haloes — dark matter

1 INTRODUCTION

Our physical picture of the Universe as a whole has been shaped by accurate observations and modeling of our own Galaxy. Explaining the matter budget and kinematical state of the Milky Way (MW) is equivalent to figuring out its formation process in a cosmological context. As a first approximation, the MW morphology and separation into global kinematically coherent components, such as its disc and bulge, can be used to support the existence of a Dark Matter (DM) component to explain its dynamics around the solar neighborhood (Olling & Merrifield 2000; Sofue et al. 2009; Catena & Ullio 2010; Bovy & Rix 2013; Iocco et al. 2015).

A more detailed description of the full three-dimensional MW gravitational potential is possible through the interpretation of fossil records of stellar streams. These remnants, resulting from infalling globular clusters or satellite galaxies that got tidally disrupted by the gravitational potential of the Milky Way, allow in principle tighter constraints on the shape of the dark matter halo in outer regions of our Galaxy (Johnston 1998; Helmi & White 1999;

Tremaine 1999) than dynamical measurements close to the solar neighborhood.

The observational constraints on the gravitational potential shape can then be confronted against the expectations from different galaxy formation models in an explicit cosmological context. For instance, in the current dominant paradigm of Cold Dark Matter (CDM) dominated Universe galaxies are expected to be hosted by triaxial DM halos. To what extent the CDM expectations are born out by observations in our Galaxy? Can the MW’s DM halo shape be considered typical or atypical in a cosmological context?

These two questions have been challenging to address because for a long time it was difficult to produce realistic galactic discs within the CDM context (Navarro & Steinmetz 1997). The success of such enterprise has been the result of high numerical resolution and the understanding that baryonic effects such as stellar feedback and black hole feedback play an important role in forming a galactic disc resembling the Milky Way. Along the way, many numerical experiments have also shown that the baryonic effects also impact the dark matter halo shape making the halo rounder than it otherwise be in a dark matter only simulation (Dubinski 1994; Debattista et al. 2008; Kazantzidis et al. 2010; Abadi et al. 2010; Bryan et al. 2013; Chua et al. 2019; Artale et al. 2019). But

* E-mail: jd.prada1760@uniandes.edu.co

only few simulations have presented detailed results on another important aspect of the disc-halo relationship, namely, the radial evolution of the alignment between the stellar disc and the dark matter halo (Bailin et al. 2005; Debattista et al. 2013).

Observational constraints of the dark matter halo have also been in tension and rapidly evolving during the last decade. One can find studies favoring prolate (Banerjee & Jog 2011; Bowden et al. 2016), oblate (Law & Majewski 2010; Deg & Widrow 2013; Vera-Ciro & Helmi 2013) and spherical configurations (Bovy et al. 2016). Constraints from modeling of stellar streams discard the prolate configuration (Law & Majewski 2010; Pearson et al. 2015; Bovy et al. 2016) although some other studies still doubt that stellar streams can be used to constrain the halo shape once other assumptions, such as the density profile, are relaxed (Ibata et al. 2013).

In this paper we analyze the internal DM halo shape in thirty MW type galaxies from state-of-the-art simulations from the Auriga project. These simulations have large enough numerical resolution, and explicit cosmological context and an appropriate feedback physics to produce realistic MW discs. Furthermore, the relatively large number of simulated systems allows us to have a handle on the statistical significance of our results. This paper is structured as follows. In Section 2 we present the most relevant details of the simulations, in Section 3 we present the method that we use to measure the DM halo shape. In Section 4 we present our results focusing on the radial shape trends at $z = 0$ and the alignments with the stellar disc. We discuss our results in Section 5 to place them into the context of other numerical work, explore correlations of the shape with baryonic properties in the disc and finally make a direct comparison against observational constraints for the MW's dark matter halo shape. We finalize with our conclusions in Section 6.

2 NUMERICAL SIMULATIONS

The Auriga project offers cosmological zoom in simulations of MW-sized dark matter halos in a Λ CDM cosmology. This simulations come in two versions: dark matter only and baryonic physics including magnetohydrodynamics. A detailed description of the simulations and their disc properties can be found in (Grand et al. 2017). Here we summarize its main features.

The objects in the simulations were selected from a set of 30 isolated halos in the Evolution and Assembly of GaLaxies and their Environments (EAGLE) project (Schaye et al. 2015). These halos were randomly selected from a sample of the most isolated halos at $z = 0$ whose virial mass M_{200} was between $10^{12} M_{\odot}$ and $2 \times 10^{12} M_{\odot}$. The cosmological parameters in these simulations correspond to $\Omega_m = 0.307$, $\Omega_b = 0.048$, $\Omega_{\Lambda} = 0.693$ and a dimensionless Hubble parameter $h = 0.6777$ (Planck Collaboration et al. 2014).

The selected halos were re-simulated at higher resolution by applying a zoom-in technique with varying physical realism using the moving-mesh AREPO code that includes gravity, ideal magnetohydrodynamics, phenomenological descriptions for star formation, chemical enrichment from supernovae and its stellar feedback. (Springel 2010; Pakmor & Springel 2013). The simulation also follows the formation

and evolution of black holes together with the Active Galactic Nuclei feedback

The 30 zoom-in halos have a dark matter particle mass of $\sim 3 \times 10^5 M_{\odot}$ while the baryonic mass resolution is $\sim 5 \times 10^4 M_{\odot}$. The softening length for gravitational force computation for stellar particles and high-resolution dark matter particles is fixed to be $500 h^{-1}$ pc in comoving coordinates up to $z = 1$, and 396 pc in physical coordinates afterwards. The gravitational softening length for gas cells changes with the mean cell radius but is limited to be larger than the stellar softening length and 1850 pc physical. From now on we refer to the haloes simulated with baryonic physics as the MHD sample and to the haloes simulated with dark matter only as the DMO sample.

3 HALO SHAPE MEASUREMENT

The DM halo shape is an estimate of either the isopotential or isodensity surfaces. Observational inference models usually estimate the isopotential contours which are probed by tracers (gas, stars), while simulations work with the isodensity contours which can be directly calculated from particle positions. Furthermore, the density contours in thin shells are very sensitive to the presence of small satellites. For this reason we measure the shape by taking volume-enclosed particles, rather than shell-enclosed. This method yields results in good agreement to the isodensity contours for radii ≤ 140 kpc as explored by (Vera-Ciro et al. 2011).

In particular, we measure the shape using the reduced inertia tensor (Allgood et al. 2006),

$$I_{ij} = \sum_k \frac{x_k^{(i)} x_k^{(j)}}{d_k^2}, \quad (1)$$

where the particle positions are measured from the minimum of the gravitational potential in each halo and each is weighted by the k -th particle distance $d_k^2 = x_k^2 + y_k^2 + z_k^2$.

The diagonalization of this tensor yields the eigenvectors and eigenvalues that represent an ellipsoidal dark matter halo. The axis lengths of this ellipsoid $a \geq b \geq c$ are the square root of the \mathbf{I} eigenvalues and the direction of the principal axis are the corresponding eigenvectors

We start the calculations taking into account particles within a sphere of radius R_{initial} and then recharacterize the triaxial parameters by taking into account particles within an ellipsoid of semi-axes $r, r/q, r/s$ and re-scaled distance $d^2 = x^2 + (y/q)^2 + (z/s)^2$, where $q = b/a$ and $s = c/a$ are the previously calculated axial ratios. We repeat this process until the average deviation of semi-axes is less than 10^{-6} . After convergence we define a unique radius R as the geometrical mean of the axial lengths $R = (abc)^{1/3}$. We use this radial coordinate R to parameterize the spatial changes in halo shape we report in the following sections. This is the same method used to estimate the halo shape in the DM-only Aquarius simulations (Vera-Ciro et al. 2011).

Following the convergence criterion by Vera-Ciro et al. (2011) in DM only simulations we restrict the measurement of the ellipsoidal parameters to radii between 2kpc and R_{200} where R_{200} corresponds to the radius computed enclosing a sphere with 200 times the critical density of the Universe. This scale is similar to the gravitational smoothing length




Figure 1. DM density in logarithmic scale within a slice of one tenth of the virial radius. The cut is perpendicular to the short axis of the inertia tensor ellipsoid. The black ellipses show the results of the fitting procedure described in Section 3. Upper/lower panels correspond to DMO/MHD simulations, respectively. Left/right panels show data at small/large radii, respectively. This plot showcases the most noticeable effect in all halos across the Auriga simulations: haloes are rounder at all radii after baryonic physics is included.

for stellar particles. For this reason, in order to make the DM halo shape results comparable against stellar disc shape measurements, we report our DM halo shape results above 10 kpc. For the shape of the stellar disc we include all stellar particles within a radius of $0.01R_{200}$, that is ≈ 23 kpc.

In our results we express the radius as a fraction of R_{200} in order to have a self-consistent dimensionless radius scale across all halos. Given the small variance across halos, each dimensionless radius is associated with a physical value with a mean value and a standard deviation. For instance, over the 30 halos in the sample, the dimensionless radius R_{200} corresponds to a physical distance of 230 ± 16 kpc.

4 RESULTS

4.1 Triaxiality

In the DMO sample we find that halos are rounder with increasing radius. The upper panels in Figure 1 illustrate this effect. The contours show a projected DM slice while the ellipsoid corresponds to the full 3D shape determination. There we see a highly ellipsoidal halo shape at radii ~ 3 kpc that becomes less triaxial at ~ 50 kpc.

We summarize this trend in Figure 2 by plotting the results of all the 30 halos in the DMO sample. The left panel shows every halo in the $c/a-b/a$ plane at two different radii




Figure 2. Axial ratios for all simulated halos. Left/right panel corresponds to DMO/MHD simulations, respectively. Triangles (squares) represent the measurements at $R_{200}/16$ (R_{200}) that correspond to physical distances of 14 ± 1 kpc (230 ± 15 kpc). Here we visualize three main population trends. First, in DMO simulations halos are rounder in the outskirts than in the inner part. Second, halos in MHD are rounder than its DMO counterparts. Third, halos in MHD are rounder in the inner regions than in the outskirts (opposite to the DMO trend).




Figure 3. Triaxiality cumulative distribution at five different radii. Right/left panel corresponds to DMO/MHD simulations, respectively. In DMO simulations the median triaxiality at all radii is larger than $2/3$. Furthermore, the triaxiality increases as one moves towards the inner part of the halo. In MHD simulations these trend reverses. The median triaxiality at all radii is smaller than $1/3$ and the halo is less triaxial as one moves towards the stellar disc.

$R_{200}/8$ (~ 20 kpc) and R_{200} . The outer part of the halo is systematically rounder than its inner region. Nevertheless the halo shape can still be considered to be prolate at all radii.

A different picture presents itself in the MHD sample. There all halos become rounder at all radii than its DMO version. The lower panel in Figure 1 can be directly compared to its MHD counterpart; there we observe how at large radii the halo becomes oblate and almost spherical. The right panel in Figure 2 shows the results for the 30 halos in the MHD sample.

In Figure 3 we summarize the results at different radii using the cumulative distributions for the triaxiality parameter T defined as

$$T = \frac{a^2 - b^2}{a^2 - c^2}. \quad (2)$$

The left panel of Figure 3 shows that in the DMO sample the triaxiality has a median larger than $2/3$ (a typical value that marks the transition from triaxiality to prolateness) at all radii. Furthermore this median value increases as




Figure 4. Cumulative distribution for the changes in triaxiality in simulations with different physics, $\Delta T = T_{\text{MHD}} - T_{\text{DMO}}$. At the virial radius all halos become less triaxial in the MHD simulations, the change is stronger towards the halo center. Only a small fraction of the halos ($\sim 1/15$) presents the opposite trend of becoming more triaxial towards the center.

we move towards the inner part of the halo. The right panel shows how the correlations in the MHD simulations go in the opposite direction with respect to DMO results. There the median triaxiality is always smaller than $1/3$ (a typical value that marks the transition from triaxiality to prolateness) and this median value decreases as we move towards the inner part of the halo.

We now quantify this change in triaxiality at the level of individual halos. We compute $\Delta T \equiv T_{\text{MHD}} - T_{\text{DMO}}$ to quantify the change between the triaxiality in the MHD and the DMO simulation. Figure shows the cumulative distribution at the same radii as in Figure . From this Figure we observe that at R_{200} all the halos have reduced its triaxiality include baryonic physics. At smaller radii this general trend continues, although there is a small fraction ($\approx 1/15$) of halos that slightly increased their triaxiality with MHD physics. However, this small fraction correspond to halos that already were outliers in the DMO sample and had triaxialities around $1/3$, considerably lower than its parent sample.

4.2 Halo-disc alignments

A common assumption in dynamical models of the MW DM halo is that its minor axis is perfectly aligned with the stellar disc minor axis. Although it is a reasonable assumption to guarantee the stability of the galactic disc in simplified models of isolated galaxies, this might not perfectly hold in an explicit cosmological context. To examine the degree of validity of this assumption we study in this subsection the alignment between the eigenvectors of the inertia tensor of stellar particles within $0.1R_{200}$ ($\sim 23 \pm 2$ kpc) and the eigenvectors of the dark matter halo shape.

In Figure 5 we summarize our main results regarding these alignments with the halo shape measured at five different radii. The upper row shows the alignment of the halos

in the DMO simulations with the stellar disc in the MHD simulations. The main objective of this measurement is to calibrate the radial evolution of the DM halo shape. We find that the DM shape remains constant with radius.

The upper row in Figure 5 shows the alignments between the major/intermediate/minor axis of the disk and the halo as a function of the dimensionless radius at which the halo shape was measured. The upper row shows the angle between the DMO halo and the stellar disk in the MHD simulation. This comparison help us to measure how the shape directions in the dark matter shells change with radius. For the bulk of the sample the lines are horizontal. Only two halos in the sample show a significant change in alignment in on the three shape axis.

The lower row in Figure 5 shows the results for the MHD simulations. The most striking feature of this plot is that around the radius of $0.25R_{200}$ there is an almost perfect halo-disc alignment measured in three shape axis. Above and below this radius the degree of alignment changes. We find the strongest misalignments at a radius of $0.06R_{200}$ when the median and the standard deviation of the alignment angles between the minor/median/major axis are 18 ± 18 , 25 ± 22 and 7 ± 21 degrees, respectively; at $0.25R_{200}$ these values drop to 2 ± 3 , 2 ± 4 and 1 ± 3 degrees, respectively.

This changing alignment as a function of radius can be seen as a twisting in the shape ellipsoids. As we are dealing with oblate halos $b/a \approx 1$ and the directions corresponding to the medium and major axis could be subject to noise. For this reason, in order to better quantify the twisting for each halo we take the standard deviation of the angle between the two minor axis, which should be less susceptible to noise. With this measurement, twisting becomes a global property of the halo, as the standard deviation is measured over the different radii at which we have quantified the alignment.

Figure 8 shows the correlation between twisting at the four different radii at which we have measured the alignments. The median, mean and standard deviation of the triaxiality are 3, 7 and 8 degrees, respectively. The correlation between twisting and triaxiality also changes with radius as measured by the Spearman's rank correlation coefficient. The largest correlation shows with the halo triaxiality measured at $0.12R_{200}$. The most important consequence of this changing degree in correlation is that twisting cannot only be explained by large triaxiality values and a possibly noisy determination of the shape axis.

4.3 Correlation with baryonic disc properties

The presence of baryons produce rounder dark matter halos than its DMO counterpart. To better understand this relationship we quantify the correlation between halo shape with baryonic disc properties. Looking into the measurements already reported by Grand et al. (2017) and Pakmor et al. (2017) we find three baryonic quantities that have the strongest correlation with DM halo triaxiality: the central gas density in a sphere of radius 1kpc, the disc to total mass ratio and the optical radius.

Figure 7 shows the correlations of these quantities with the triaxiality at five different radii. We use the Spearman's rank correlation coefficient to quantify the correlation strength. We find that the strongest correlations are found with the halo shape measured at radii smaller than $0.12R_{200}$.




Figure 5. Angles between the principal axis of the dark matter halo shape and the stellar disc shape as a function of the radius at which the halo shape directions are measured. Each panel compares the alignment of the corresponding major/middle/minor axis. Thin lines correspond to each one of the thirty halos in the sample while the thick line traces the median value at every radius. In the upper row the haloes come from the DMO simulation, showing that the directions in the ellipsoids describing the shape are constant as a function of radius for the most part of the sample. In the lower row the haloes come from the MHD simulation providing a self-consistent comparison with the stellar discs. In this case the dark matter shells twists significantly in a noticeable fraction of the haloes. Interestingly, an almost perfect halo-disc alignment happens across the sample at an intermediate radius of $0.25R_{200}$ (56 ± 4 kpc).




Figure 6. Twisting in the halo-disc alignment (measured as the standard deviation of the alignment angles between two minor shape axis) as a function of the halo triaxiality at the radius indicated in each panel's title. The label with the ρ value corresponds to the Spearman's rank correlation coefficient while the line shows the result of the minimum squares fit to the data. Twisting does not correlate with triaxiality in the same way at every radii. The strongest correlation appears around $0.12R_{200}$ (28 ± 2 kpc).




Figure 7. Correlations between the halo triaxiality at different radii and baryonic disc properties. The label with the ρ value corresponds to the Spearman's rank correlation coefficient. The line is the best minimum squares fit to a line. The x-axis in the first column is the gas density at the center of the galaxy within a sphere of radius 1 kpc (Pakmor et al. 2017); the second column shows the disc to total mass ratio and the last column includes the disc optical radius defined to be the radius at which the B -band surface brightness drops below 25 mag arcsec $^{-2}$ (Grand et al. 2017). The largest correlations are found for the two smaller radii ($0.06R_{200}$ and $0.12R_{200}$). Extended and massive stellar discs with a low gas content at its core are correlated with low dark matter triaxialities. The correlation decreases as one approaches larger radii.




Figure 8. Twisting in the halo-disc alignment as a function of the same baryonic disc properties in Figure 7. The label with the ρ value corresponds to the Spearman's rank correlation coefficient. The line is the best minimum squares fit to a line. In this case the disc to total mass ratio does not correlate strongly with the alignment twisting. The gas density at the center and the disc optical radius still show a strong correlation.

The trend is such that halos with large triaxiality correlate with high gas density and stellar discs with low mass and small size. In turn, massive and large stellar discs within with low density of gas in its core correlate with low halo triaxiality.

The halo shape also twists in the presence of baryons, as opposed to the situation in DMO simulations where the shape is coherent as a function of radius. In Figure 8 we show the correlation between twisting and the same baryonic properties that show a high degree of correlation with the triaxiality. In this case the disc to total ratio does not have a significant degree of correlation. Instead, the disc optical radius and the gas density at the core are the most correlated. The correlation is such that extended stellar discs with a very low gas density at its core show the most coherent halo shapes.

5 DISCUSSION

The first effect that we put in evidence in this paper is the effect of baryons in producing rounder DM halos. This effect has been already widely reported in the literature. It has been found that the strength of the change depends on the numerical resolution, the gas cooling implementation and the models describing star formation and stellar feedback (Bailin et al. 2005; Debattista et al. 2008; Bryan et al. 2013; Butsky et al. 2016; Chua et al. 2019; Artale et al. 2019). The key concept unifying these results is that the baryon distribution influences and correlates with the dark matter halo shape. Indeed, as we presented in the previous section on the correlations with disc properties, we find that the broad tendency is that extended massive stellar discs correlate with spherical dark matter distributions.

In contrast, there are fewer studies reporting on the halo-disc alignment as a function of radius. Jing & Suto (2002) reported a shell alignment twisting in their dark matter only simulations as measured in the direction changes of large and medium shape axis. They only had three high resolution MW-like halos (with $\sim 10^6$ inside the virial radius) and could not make a statistical statement about the impact of this effect. However, they see the twisting as an artifact

resulting from high values of the b/a ratios that make the determination of the medium axis noisy. For that reason we measured the twisting using the minor axis alignments, however they do not report these alignments.

Bailin et al. (2005) used seven hydrodynamical simulations studied the disc halo alignment as a function of radius. For radii smaller than $0.1R_{200}$ they find a strong alignment between the two minor axis of the halo and the disc. For larger radii the angle between the components changes showing a weak correlation between the two components. Debattista et al. (2013) performed a study on the disc-halo alignment performing controlled experiments on three different halos to study changes the evolutionary change that drive the minor axis in the disc and the halo to be aligned, their study by construction focuses only on radii around $0.1R_{200}$. Nevertheless, similarly to Bailin et al. (2005), they also report a strong alignment in the region where the disc dominated. In conclusion, the twisting in the dark matter halo shapes seems to be a feature already reported in the literature with smaller samples than the one we use here. The strongest point of depart is that in our case the alignments do not grow stronger as one approached the disc. It is very likely that the detailed radial dependence of the twisting depends on the hydrodynamics and feedback implementations.

We now use the results reported by Law & Majewski (2010) and Bovy et al. (2016) to place our results in an observational context. Law & Majewski (2010) used observations of the Sagittarius tidal stream to constrain the shape of the gravitational potential. Their point of depart is that previous studies that assumed an axisymmetric galactic potential were not able to fit all the available dynamic constraints for the Sagittarius stream, therefore making necessary the use of a rigid triaxial potential with coaxial potential ellipsoids for the dark matter component. Their results constrain the triaxiality of this potential component. They also translate their results into a triaxiality of the density contours (that could be compared against our results) to be $(c/a) = 0.44$ and $(b/a) = 0.97$ at a radius of $\sim 40\text{kpc}$. They do not report any uncertainties for these two values. Looking at their plots for the quality of fit criterion as a function of dark halo axial scales (their Figure 5), we choose a conservative 10%

relative uncertainty. One surprising element in their results is that the major axis of the halo shape is perpendicular to the stellar disc plane.

The results by Bovy et al. (2016) have the same general approach but use instead the GD-1 (Grillmair & Dionatos 2006) and Pal 5 (Odenkirchen et al. 2009) streams to constrain the shape of the dark matter component of the galactic halo potential. They use general models with many degrees of freedom for the galactic potential in order to measure to what extent these two streams are sensitive to the triaxiality of the dark matter halo component. The DM component is written directly as a triaxial density profile with coaxial ellipsoids and the corresponding potential is found by numerical integration. They find that the width of the Pal 5 stream constraints $b/a \approx 1$ and therefore fix it to be $b/a = 1$ exactly. Using that value they report their most stringent constrain of $c/a = 0.93 \pm 0.16$ at a radius of ≈ 19 kpc from the galactic center.

Figure 9 shows an explicit comparison in the c/a - b/a plane of our results in MHD simulations against the results by Law & Majewski (2010) and Bovy et al. (2016). We find six MHD halos with $b/a < 0.93$ and $c/a > 0.77$ that could be considered consistent with their shape constraints by Bovy et al. (2016), while only one outlier DMO halo is consistent with those constraints. In contrast, none of the simulated halos (MHD nor DMO) is consistent with the results Law & Majewski (2010). The change of triaxiality with radius in our simulations cannot account for these two extremely different shape constraints at different radii.

The results by could then place the dark matter halo of our Milky Way as an extreme outlier in the Λ CDM model. This extreme prolateness also correlates with the extreme triaxiality of the 11 classical satellites of the MW ($c/a \approx 0.2$, and $b/a \approx 0.9$) measured at larger radii, with an spatial distribution also oriented perpendicular to the MW plane, another highly unusual feature in the Λ CDM (Forero-Romero & Arias 2018).

6 CONCLUSIONS

In this paper we measured the shape of 30 isolated Milky Way sized dark matter haloes simulated in the Auriga project using the zoom-in technique. The halos were with two different setups: dark matter only (DMO) simulation and full magnetohydrodynamics (MHD) including star formation and feedback. We used the shape measurement algorithm by Allgood et al. (2006) on the dark matter halos of these simulations to quantify the halo shape as a function of radius and the degree of alignment between the stellar disc and the halo.

We find that MHD halos are rounder than DMO halos at every radius. MHD halos tend towards more oblate, sometimes almost spherical, shapes ($T < 1/3$), while DMO halos tend towards more prolate shapes ($T > 2/3$). The rounding effect is more noticeable as one moves closer to galactic disc and strongly correlates with baryonic properties of the disc. More precisely, the triaxiality is smaller for extended and massive stellar discs with low gas densities at its core.

We also measured the shape alignment of the halo with the stellar disc at different radii. At a radius of $0.25R_{200}$ the alignment is almost perfect and degrades at smaller and

larger distances. In some halos the alignment between the stellar disc and dark matter halo changes noticeably with radius. This alignment evolution implies a radial twisting between the ellipsoids describing the halo shape. We quantify this twist with the standard deviation of the angle between the two minor axis at radii below $\leq 0.5R_{200}$ and find that the twisting strongly correlates with the disc size and the core gas mass density, there is only a weak correlation with the disc to total mass ratio.

We compared our results against two observational constraints for the dark matter halo shape of the Milky Way. The constraints are at two different radii and come from different observational tracers. We find that 20% halos in the MHD simulations are consistent with the constraints by Bovy et al. (2016) at ≈ 19 kpc that correspond to an almost spherical halo, while none of the halos, either in MHD or DMO, has some overlap with the shape constraints by Law & Majewski (2010) at ≈ 40 kpc that is closer to oblate.

Using dark matter only simulations Vera-Ciro et al. (2011) suggested that the current dark matter halo shape strongly correlates with the time evolution of the halo as traced by the shape measured at the virial radius. This effect might no longer hold once baryons are included. The opposed trend of triaxiality in MHD simulation as a function of radius to the DMO simulations together with the twisting effect in some of the halos seems to suggest that that historical buildup memory is not stored in the halo at $z = 0$.

A more complete understanding of the influence of baryons on the different properties we have measured will require at least two more elements. First, a study on how the halo and the disc co-evolved as a function of time. Second, a calibrating the effect of the cosmic web (both dark matter and gaseous) on any evolutionary trend (Forero-Romero et al. 2014; Borzyszkowski et al. 2017; Ganeshaiah Veena et al. 2019) We finalize by suggesting that the twisting density shells we find in some of the halos is feature that deserves to be explored in the process of constraining shape parameters from tidal stream data. The inclusion of a parameterization describing this degree of twisting might relax the conflict between the observational constraints and the numerical results.

ACKNOWLEDGEMENTS

This project has received funding from the European Union's Horizon 2020 Research and Innovation Programme under the Marie Skłodowska-Curie grant agreement No 734374.

REFERENCES

- Abadi M. G., Navarro J. F., Fardal M., Babul A., Steinmetz M., 2010, *MNRAS*, **407**, 435
- Allgood B., Flores R. A., Primack J. R., Kravtsov A. V., Wechsler R. H., Faltenbacher A., Bullock J. S., 2006, *MNRAS*, **367**, 1781
- Artale M. C., Pedrosa S. E., Tissera P. B., Cataldi P., Di Cintio A., 2019, *A&A*, **622**, A197
- Bailin J., et al., 2005, *ApJ*, **627**, L17
- Banerjee A., Jog C. J., 2011, *ApJ*, **732**, L8




Figure 9. Comparison of our results against observational constraints for the dark matter halo shape in the Milky Way by [Law & Majewski \(2010\)](#) (LM10) and [Bovy et al. \(2016\)](#) (BBFK16). We find that 1/5 of the halos in the MHD sample are consistent with the constraints by [Bovy et al. \(2016\)](#). In contrast, none of the halos in the MHD and DMO seems to be consistent with the results by [Law & Majewski \(2010\)](#).

Borzyszkowski M., Porciani C., Romano-Díaz E., Garaldi E., 2017, *MNRAS*, **469**, 594
 Bovy J., Rix H.-W., 2013, *ApJ*, **779**, 115
 Bovy J., Bahmanyar A., Fritz T. K., Kallivayalil N., 2016, *ApJ*, **833**, 31
 Bowden A., Evans N. W., Williams A. A., 2016, *MNRAS*, **460**, 329
 Bryan S. E., Kay S. T., Duffy A. R., Schaye J., Dalla Vecchia C., Booth C. M., 2013, *MNRAS*, **429**, 3316
 Butsky I., et al., 2016, *MNRAS*, **462**, 663
 Catena R., Ullio P., 2010, *J. Cosmology Astropart. Phys.*, **8**, 004
 Chua K. T. E., Pillepich A., Vogelsberger M., Hernquist L., 2019, *MNRAS*, **484**, 476
 Debattista V. P., Moore B., Quinn T., Kazantzidis S., Maas R., Mayer L., Read J., Stadel J., 2008, *The Astrophysical Journal*, **681**, 1076
 Debattista V. P., Roškar R., Valluri M., Quinn T., Moore B., Wadsley J., 2013, *MNRAS*, **434**, 2971
 Deg N., Widrow L., 2013, *MNRAS*, **428**, 912
 Dubinski J., 1994, *ApJ*, **431**, 617
 Forero-Romero J. E., Arias V., 2018, *MNRAS*, **478**, 5533
 Forero-Romero J. E., Contreras S., Padilla N., 2014, *MNRAS*, **443**, 1090
 Ganeshaiah Veena P., Cautun M., Tempel E., van de Weygaert R., Frenk C. S., 2019, *MNRAS*, **487**, 1607
 Grand R. J. J., et al., 2017, *Monthly Notices of the Royal Astro-*

nomical Society, **467**, 179
 Grillmair C. J., Dionatos O., 2006, *ApJ*, **641**, L37
 Helmi A., White S. D. M., 1999, *MNRAS*, **307**, 495
 Ibata R., Lewis G. F., Martin N. F., Bellazzini M., Correnti M., 2013, *ApJ*, **765**, L15
 Iocco F., Pato M., Bertone G., 2015, *Nature Physics*, **11**, 245
 Jing Y. P., Suto Y., 2002, *ApJ*, **574**, 538
 Johnston K. V., 1998, *ApJ*, **495**, 297
 Kazantzidis S., Abadi M. G., Navarro J. F., 2010, *ApJ*, **720**, L62
 Law D. R., Majewski S. R., 2010, *The Astrophysics Journal*, **714**, 229
 Navarro J. F., Steinmetz M., 1997, *ApJ*, **478**, 13
 Odenkirchen M., Grebel E. K., Kayser A., Rix H.-W., Dehnen W., 2009, *AJ*, **137**, 3378
 Olling R. P., Merrifield M. R., 2000, *MNRAS*, **311**, 361
 Pakmor R., Springel V., 2013, *MNRAS*, **432**, 176
 Pakmor R., et al., 2017, *MNRAS*, **469**, 3185
 Pearson S., Küpper A. H. W., Johnston K. V., Price-Whelan A. M., 2015, *ApJ*, **799**, 28
 Planck Collaboration et al., 2014, *A&A*, **571**, A16
 Schaye J., et al., 2015, *Monthly Notices of the Royal Astronomical Society*, **446**, 521
 Sofue Y., Honma M., Omodaka T., 2009, *PASJ*, **61**, 227
 Springel V., 2010, *Monthly Notices of the Royal Astronomical Society*, **401**, 791
 Tremaine S., 1999, *MNRAS*, **307**, 877

Vera-Ciro C., Helmi A., 2013, [ApJ](#), **773**, [L4](#)

Vera-Ciro C. A., Sales L. V., Helmi A., Frenk C. S., Navarro J. F.,
Springel V., Vogelsberger M., White S. D. M., 2011, [MNRAS](#),
416, [1377](#)