Supplemental Table 1. A descriptive summary table generated by TernTables.

Category Variable	Total (N = 929)
Patient Demographics	
Age (yr)	59.8 ± 11.9
Sex	
Female	445 (48%)
Male	484 (52%)
Surgical Findings	
Colonic Obstruction	180 (19%)
Bowel Perforation	27 (3%)
Tumor Characteristics	
Positive Lymph Nodes (n)	2 [1–5]
> 4 Positive Nodes	255 (27%)
Tumor Adherence	135 (15%)
Tumor Differentiation	
Well	93 (10%)
Moderate	663 (73%)
Poor	150 (17%)
Extent Of Local Spread	
Submucosa	21 (2%)
Muscle	106 (11%)
Serosa	759 (82%)
Contiguous Structures	43 (5%)
Outcomes	
Recurrence	
No Recurrence	461 (50%)
Recurrence	468 (50%)
Treatment Arm	
Levamisole + 5FU	304 (33%)
Levamisole	310 (33%)
Observation	315 (34%)

Supplemental Table 2. A two-group comparison table generated by TernTables.

Category Variable	No Recurrence (n = 461)	Recurrence (n = 468)	Total (N = 929)	OR	P value
Patient Demographics					
Age (yr)	60.5 ± 11.5	59 ± 12.4	59.8 ± 11.9	-	0.065
Sex					
Female	216 (47%)	229 (49%)	445 (48%)	1.00 (ref.)	0.570
Male	245 (53%)	239 (51%)	484 (52%)	0.92 [0.71–1.19]	-
Surgical Findings					
Colonic Obstruction	81 (18%)	99 (21%)	180 (19%)	1.26 [0.91–1.75]	0.194
Bowel Perforation	10 (2%)	17 (4%)	27 (3%)	1.70 [0.77–3.75]	0.258
Tumor Characteristics					
Positive Lymph Nodes (n)	2 [1–3]	3 [2–6]	2 [1–5]	-	**2E-14**
> 4 Positive Nodes	75 (16%)	180 (38%)	255 (27%)	**3.22 [2.36–4.38]**	**6E-14**
Tumor Adherence	53 (11%)	82 (18%)	135 (15%)	**1.64 [1.13–2.37]**	**0.012**
Tumor Differentiation					
Well	49 (11%)	44 (10%)	93 (10%)	-	0.089
Moderate	337 (75%)	326 (71%)	663 (73%)	-	-
Poor	62 (14%)	88 (19%)	150 (17%)	-	-
Extent Of Local Spread					
Submucosa	16 (3%)	5 (1%)	21 (2%)	-	**7E-6**
Muscle	72 (16%)	34 (7%)	106 (11%)	-	-
Serosa	359 (78%)	400 (85%)	759 (82%)	-	-
Contiguous Structures	14 (3%)	29 (6%)	43 (5%)	-	-
Treatment Details					
Treatment Arm					
Levamisole + 5FU	185 (40%)	119 (25%)	304 (33%)	-	**1E-5**
Levamisole	138 (30%)	172 (37%)	310 (33%)	-	-
Observation	138 (30%)	177 (38%)	315 (34%)	-	-

Supplemental Table 3. A three-group comparison table generated by TernTables.

Category Variable	Observation (n = 315)	Levamisole (n = 310)	Levamisole + 5FU (n = 304)	Total (N = 929)	P value
Patient Demographics					
Age (yr)	59.5 ± 12	60.1 ± 11.6	59.7 ± 12.3	59.8 ± 11.9	0.780
Sex					
Female	149 (47%)	133 (43%)	163 (54%)	445 (48%)	**0.028**
Male	166 (53%)	177 (57%)	141 (46%)	484 (52%)	-
Surgical Findings					
Colonic Obstruction	63 (20%)	63 (20%)	54 (18%)	180 (19%)	0.683
Bowel Perforation	9 (3%)	10 (3%)	8 (3%)	27 (3%)	0.907
Tumor Characteristics					
Positive Lymph Nodes (n)	2 [1–5]	2 [1–5]	2 [1–4]	2 [1–5]	0.550
> 4 Positive Nodes	87 (28%)	89 (29%)	79 (26%)	255 (27%)	0.749
Tumor Adherence	47 (15%)	49 (16%)	39 (13%)	135 (15%)	0.562
Tumor Differentiation					
Well	27 (9%)	37 (12%)	29 (10%)	93 (10%)	0.522
Moderate	229 (74%)	219 (73%)	215 (72%)	663 (73%)	-
Poor	52 (17%)	44 (15%)	54 (18%)	150 (17%)	-
Extent Of Local Spread					
Submucosa	8 (3%)	3 (1%)	10 (3%)	21 (2%)	0.264
Muscle	38 (12%)	36 (12%)	32 (11%)	106 (11%)	-
Serosa	249 (79%)	259 (84%)	251 (83%)	759 (82%)	-
Contiguous Structures	20 (6%)	12 (4%)	11 (4%)	43 (5%)	-
Outcomes					
Recurrence					
No Recurrence	138 (44%)	138 (45%)	185 (61%)	461 (50%)	**1E-5**
Recurrence	177 (56%)	172 (55%)	119 (39%)	468 (50%)	-