

OPEN ACCESS

EDITED BY

Neus Garcias-Bonet, King Abdullah University of Science and Technology, Saudi Arabia

REVIEWED BY

Julie L. Meyer, University of Florida, United States Alejandra Prieto-Davó, National Autonomous University of Mexico, Mexico Amanda Shore, Farmingdale State College, United States

*CORRESPONDENCE

Jason D. Selwyn

jason.selwyn@tamucc.edu

[†]PRESENT ADDRESS

Jason D. Selwyn, Genomics Core Laboratory, Texas A&M University - Corpus Christi, Corpus Christi, TX, United States

RECEIVED 04 September 2024 ACCEPTED 03 February 2025 PUBLISHED 20 February 2025

CITATION

Selwyn JD, Despard BA, Galvan-Dubois KA, Trytten EC and Vollmer SV (2025) Antibiotic pretreatment inhibits white band disease infection by suppressing the bacterial pathobiome. Front. Mar. Sci. 12:1491476. doi: 10.3389/fmars.2025.1491476

COPYRIGHT

© 2025 Selwyn, Despard, Galvan-Dubois, Trytten and Vollmer. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted

which does not comply with these terms.

Antibiotic pretreatment inhibits white band disease infection by suppressing the bacterial pathobiome

Jason D. Selwyn^{1,2*†}, Brecia A. Despard^{1,2}, Kai A. Galvan-Dubois^{1,2}, Emily C. Trytten^{1,2} and Steven V. Vollmer^{1,2}

¹Marine Science Center, Northeastern University, Nahant, MA, United States, ²Department of Marine and Environmental Sciences, Northeastern University, Boston, MA, United States

Diseases have caused unprecedent mortality in Caribbean coral communities. White band disease (WBD) has killed up to 95% of all endangered Caribbean Acroporids since it was first observed in 1979. Despite the devastating impacts of WBD, its etiology is currently unknown although recent research identified two bacterial strains - ASVs classified as a Cysteiniphilum litorale and a Vibrio sp., as the most likely pathogens. To better understand the disease etiology of WBD, we pretreated corals with antibiotics to determine how prophylactic use of antibiotics impacts the transmission of WBD in a replicated tank-based experiment. We found the prophylactic use of antibiotics led to significantly reduced infection rates in disease exposed corals with a 30-percentage point decrease in the infection rate. Analyses of 16S rRNA amplicon gene sequencing data in the disease exposed corals demonstrated that antibiotic pretreatment resulted in coral microbiomes which were less speciose and contained relatively fewer Vibrio spp. than untreated corals, indicating that the benefit of the antibiotic pretreatment was its ability to reduce the relative abundance of intrinsic secondary opportunists and/or opportunistic pathogens suggesting their likely importance to the etiology of WBD. We propose two distinct etiologies involving either an extrinsic keystone pathogen (Cysteiniphilum litorale) or overgrowth of intrinsic opportunistic pathogens (Vibrio spp.). Future research should isolate these strains to confirm the etiology of white band disease.

KEYWORDS

white band disease, *Acropora*, keystone pathogen, opportunistic pathogen, *Vibrio*, *Cysteiniphilum litorale*