

RESEARCH ARTICLE

Kin-Aggregations Explain Chaotic Genetic Patchiness, a Commonly Observed Genetic Pattern, in a Marine Fish

Jason D. Selwyn^{1*}, J. Derek Hogan¹, Alan M. Downey-Wall², Lauren M. Gurski¹, David S. Portnoy³, Daniel D. Heath⁴

1 HoBi Lab, Department of Life Sciences, Texas A&M University—Corpus Christi, Corpus Christi, Texas, United States of America, **2** Marine Science Center, Northeastern University, Nahant, Massachusetts, United States of America, **3** Marine Genomics Laboratory, Department of Life Sciences, Texas A&M University—Corpus Christi, Corpus Christi, Texas, United States of America, **4** Great Lakes Institute for Environmental Research, University of Windsor, Windsor, Ontario, Canada

* jason.selwyn@tamucc.edu



OPEN ACCESS

Citation: Selwyn JD, Hogan JD, Downey-Wall AM, Gurski LM, Portnoy DS, Heath DD (2016) Kin-Aggregations Explain Chaotic Genetic Patchiness, a Commonly Observed Genetic Pattern, in a Marine Fish. PLoS ONE 11(4): e0153381. doi:10.1371/journal.pone.0153381

Editor: Heather M. Patterson, Department of Agriculture and Water Resources, AUSTRALIA

Received: January 25, 2016

Accepted: March 29, 2016

Published: April 27, 2016

Copyright: © 2016 Selwyn et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by the Natural Sciences and Engineering Research Council of Canada CRO grant# 227965 awarded to P.F. Sale, a Global Environment Facility Coral Reef Targeted Research project and by a Natural Sciences and Engineering Research Council Tier I Canada Research Chair grant to DDH; a Great Lakes Institute for Environmental Research post-doctoral fellowship from the University of Windsor and Texas A&M University—Corpus Christi start-up funds to JDH.

Abstract

The phenomenon of chaotic genetic patchiness is a pattern commonly seen in marine organisms, particularly those with demersal adults and pelagic larvae. This pattern is usually associated with sweepstakes recruitment and variable reproductive success. Here we investigate the biological underpinnings of this pattern in a species of marine goby *Coryphopterus personatus*. We find that populations of this species show tell-tale signs of chaotic genetic patchiness including: small, but significant, differences in genetic structure over short distances; a non-equilibrium or “chaotic” pattern of differentiation among locations in space; and within locus, within population deviations from the expectations of Hardy-Weinberg equilibrium (HWE). We show that despite having a pelagic larval stage, and a wide distribution across Caribbean coral reefs, this species forms groups of highly related individuals at small spatial scales (<10 metres). These spatially clustered family groups cause the observed deviations from HWE and local population differentiation, a finding that is rarely demonstrated, but could be more common than previously thought.

Introduction

Marine organisms with dispersive pelagic larvae are expected to be characterized by little to no genetic differentiation among populations over large geographic areas due to high gene flow. However, many marine species exhibit slight, yet significant, levels of genetic heterogeneity over various local spatial scales [1,2] which may be temporally unstable. Typically in these situations, molecular markers deviate from the expectations of Hardy-Weinberg equilibrium (HWE) within some samples [3–5]. This complex pattern, termed chaotic genetic patchiness (CGP), has been attributed to non-equilibrium conditions caused by variation in reproductive success of breeding adults during larval recruitment [2,6]. By seeking to understand the biological mechanisms underlying these patterns we can deepen our understanding of the ecology and evolution of complex marine populations.