

GRADE Summary of Findings — pelvic (WPRT) vs prostate-only RT (PORT)

Outcome	No. of trials (pts)	Relative effect (HR, 95% CI)	Anticipated absolute effect at 7 yr (per 1,000)	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication bias	Certainty
Overall survival	5 (5,172)	HR 1.07 (0.94 to 1.21)	PORT: 211 per 1,000 WPRT: 224 (199–249) Δ +13 (-11 to +38)	✓ No concerns	✓ No concerns	↓ Serious	✓ No concerns	– Undetected	⊕⊕⊕○ Moderate
Biochemical / progression (mixed defn)	5 (5,172)	HR 0.84 (0.54 to 1.29)	PORT: 399 per 1,000 WPRT: 348 (241–482) Δ -51 (-159 to +83)	↓ Serious	↓↓ Very serious	↓ Serious	↓ Serious	– Undetected	⊕○○○ Very low
Distant metastasis / MFS (mixed defn)	4 (4,728)	HR 0.92 (0.54 to 1.57)	PORT: 289 per 1,000 WPRT: 270 (168–415) Δ -20 (-121 to +126)	✓ No concerns	↓ Serious	↓ Serious	↓ Serious	– Undetected	⊕○○○ Very low

Anticipated absolute effects per 1,000 at 7 years: baseline = reconstructed-IPD PORT arm; WPRT = baseline transformed by the pooled HR under proportional hazards. Domains: ✓ no concerns · ↓ serious · ↓↓ very serious · – undetected. Publication bias not formally testable at k<10.