

# Weak Supervision for the Strong Force


Jesse Thaler



QUC Summer School, KIAS — July 12, 2022




# The NSF AI Institute for Artificial Intelligence and Fundamental Interactions (IAIFI /ai-fai/ iaifi.org)



*Advance physics knowledge — from the smallest building blocks of nature  
to the largest structures in the universe — and galvanize AI research innovation*




# The NSF AI Institute for Artificial Intelligence and Fundamental Interactions (IAIFI /ai-fai/ iaifi.org)



*Infuse physics intelligence  
into artificial intelligence*

*Machine learning that incorporates  
first principles, best practices, and domain knowledge  
from fundamental physics*


*Advance physics knowledge — from the smallest building blocks of nature  
to the largest structures in the universe — and galvanize AI research innovation*



# IAIFI


**Summer School & Workshop**

August 1–9, 2022



# Weak Supervision for the Strong Force


An IAI FI origin story: **Patrick and Eric** come to my office at MIT...




*“What do you mean by **quarks** and **gluons**? ”*

[Komiske, Metodiev, Schwartz, [JHEP 2017](#)]





# Weak Supervision for the Strong Force



Five years later,  
I finally know an answer!

*Required weak supervision, topic modeling,  
permutation-invariant networks,  
simulation-based inference, optimal transport, ...*

[Komiske, Kryhin, JDT, arXiv 2022]




“What do you mean by *quarks* and *gluons*? ”

[Komiske, Metodiev, Schwartz, JHEP 2017]




# Outline




2017: A Quark/Gluon Conundrum



2018: Leveraging Weak Supervision




2019-2022: The Strong Force Revisited



## 2017: A Quark/Gluon Conundrum




## 2018: Leveraging Weak Supervision





## 2019-2022: The Strong Force Revisited



# Particle Physics 101



# Particle Physics 101




# Particle Physics 101




# Collider Event

Every 25 nanoseconds at the LHC









T E H M





Run Number: 159224, Event Number: 3533152

Date: 2010-07-18 11:05:54 CEST




ATLAS  
EXPERIMENT  
<http://atlas.ch>




# What is a “Quark Jet” vs. “Gluon Jet”?

**Quark (color triplet) vs. Gluon (color octet)**  
But jet constituents are **color-singlet hadrons!**




[Gras, Hoeche, Kar, Larkoski, Lönnblad, Plätzer, Siódmok, Skands, Soyez, JDT, [JHEP 2017](#); citing long history going back at least to Nilles, Streng, [PRD 1981](#)]

# Emphasizing the Challenge



## Altarelli-Parisi Splitting


Core prediction of QCD



$$dP_{i \rightarrow ig} \simeq \frac{2\alpha_s}{\pi} C_i \frac{d\theta}{\theta} \frac{dz}{z}$$


Collinear   Soft

# Emphasizing the Challenge



# Emphasizing the Challenge

Theory



## Energy Flow:

Robust to hadronization and detector effects...  
...but blind to direct **quark/gluon** information



$$\hat{\mathcal{E}} \simeq \lim_{t \rightarrow \infty} \hat{n}_i T^{0i}(t, vt\hat{n})$$

(more about this in backup)

[see e.g. Sveshnikov, Tkachov, [PLB 1996](#); Hofman, Maldacena, [JHEP 2008](#); Mateu, Stewart, JDT, [PRD 2013](#); Belitsky, Hohenegger, Korchemsky, Sokatchev, Zhiboedov, [PRL 2014](#); Chen, Moult, Zhang, Zhu, [PRD 2020](#)]

# Color-Blind Tests of QCD

*First ever analysis using public LHC data!*




$$dP_{i \rightarrow ig} \simeq \frac{2\alpha_s}{\pi} C_i \frac{d\theta}{\theta} \frac{dz}{z}$$


*Insensitive to quark/gluon composition. Can we do the opposite?!*

[Tripathee, Xue, Larkoski, Marzani, JDT, [PRL 2017](#), [PRD 2017](#);  
based on Larkoski, Marzani, JDT, [PRD 2015](#)]






*Jets are manifestation of quarks and gluons,  
but there is **no unambiguous way**  
to tell a quark jet from a gluon jet*



2017: A Quark/Gluon Conundrum







2018: Leveraging Weak Supervision



2019-2022: The Strong Force Revisited


# Blind Source Separation for Cosmology



[Planck Outreach]

# Blind Source Separation for Quark/Gluon Jets?

While you can't unambiguously label individual jets, you can extract **quark** and **gluon** distributions from **hadron-level measurements**



In natural language processing, this is known as “**topic modeling**”

[Komiske, Metodiev, JDT, [JHEP 2018](#); cf. ATLAS, [PRD 2019](#); using Metodiev, Nachman, JDT, [JHEP 2017](#); Metodiev, JDT, [PRL 2018](#)  
[see also Blanchard, Flaska, Handy, Pozzi, Scott, [PLMR 2013](#); Katz-Samuels, Blanchard, Scott, [JMLR 2016](#)]  
[Yes, for those of you obsessed with these things, I flipped the color of quark and gluon on this slide.]





# Fundamental Assumption

One can make jet samples that are mixtures of “quarks” and “gluons”

$$p_{\text{mixed}}(\vec{x}) = f_q \, p_{\text{quark}}(\vec{x}) + (1 - f_q) \, p_{\text{gluon}}(\vec{x})$$

↳ Jet Features





Non-trivial! Assumes A/B mixtures have unbiased jet properties

# Fundamental Trick

From A/B mixtures you can define “quarks” and “gluons”

$$p_{\text{quark}}(\vec{x}) = \frac{p_A(\vec{x}) - \kappa_{AB} p_B(\vec{x})}{1 - \kappa_{AB}}$$


$$p_{\text{gluon}}(\vec{x}) = \frac{p_B(\vec{x}) - \kappa_{BA} p_A(\vec{x})}{1 - \kappa_{BA}}$$




Choosing  $\kappa$  as big as possible yields “mutually irreducible” distributions

[Katz-Samuels, Blanchard, Scott, [JLMR 2016](#); advocated for in Komiske, Metodiev, [JDT, JHEP 2018](#)]

# Jet Topics Result from ATLAS



Track multiplicity for  
“Topic 1” and “Topic 2”



[ATLAS, PRD 2019]


*“Ok, but don’t jet topics  
depend on choice of jet features?”*



Yes! But to find maximal separability,  
we can **leverage weak supervision!**  
**CWoLa = Classification Without Labels**  
(More accurately, classification with noisy labels)

# Reminder: Fully-Supervised Binary Classification

*Training on pure, perfectly labeled examples*



**Minimize Loss Function**

(asymptotically)

$$h(\vec{x}) = \frac{p_{\text{sig}}(\vec{x})}{p_{\text{sig}}(\vec{x}) + p_{\text{bkgd}}(\vec{x})}$$

**Optimal Classifier (Neyman–Pearson)**


$$\ell_{\text{MSE}} = \left\langle (h(\vec{x}) - 1)^2 \right\rangle_{\text{signal}} + \left\langle (h(\vec{x}) - 0)^2 \right\rangle_{\text{background}}$$

↑      ↑  
Classifier   Inputs

(See backup for Lagrangian formulation of what this is doing)

# Classification Without Labels (CWoLa)

*Train directly on *mixed data*!*




$$h_{\text{mixed}}(\vec{x}) = \frac{p_A(\vec{x})}{p_A(\vec{x}) + p_B(\vec{x})}$$
$$\neq$$
$$h_{\text{pure}}(\vec{x}) = \frac{p_q(\vec{x})}{p_q(\vec{x}) + p_g(\vec{x})}$$

but...

$$\frac{\partial h_{\text{mixed}}(\vec{x})}{\partial h_{\text{pure}}(\vec{x})} > 0$$



*Weak supervision yields *same decision boundaries* as strong supervision!*

[Metodiev, Nachman, [JDT, JHEP 2017](#); see also Blanchard, Flaska, Handy, Pozzi, Scott, [PLMR 2013](#); Cranmer, Pavez, Louppe, [arXiv 2015](#); Dery, Nachman, Rubbo, Schwartzman, [JHEP 2017](#); Cohen, Freytsis, Ostdiek, [JHEP 2018](#); Komiske, Metodiev, Nachman, Schwartz, [PRD 2018](#); Collins, Howe, Nachman, [PRL 2018, PRD 2019](#)]



# The Well-Read CWoLa

*Weak supervision meets topic modeling*




$$h_{\text{pure}} \in [0, 1]$$

$$h_{\text{mixed}} \in \left[ \frac{f_g^A}{f_g^A + f_g^B}, \frac{f_q^A}{f_q^A + f_q^B} \right]$$


$$h_{\text{mixed}} \in \left[ \frac{f_g^A}{f_g^A + f_g^B}, \frac{f_q^A}{f_q^A + f_q^B} \right]$$

[using Katz-Samuels, Blanchard, Scott, [JLMR 2016](#); Metodiev, Nachman, [JDT, JHEP 2017](#); Metodiev, [JDT, PRL 2018](#)]  
[Komiske, Metodiev, [JDT, JHEP 2018](#)]






*By assuming jet samples are mixtures of “quarks” and “gluons”,  
one can **operationally define** jet categories*



2017: A Quark/Gluon Conundrum




2018: Leveraging Weak Supervision



2019-2022: The Strong Force Revisited


# “Collision Course”

“Theoretical Physics for Machine Learning”  
Aspen Center for Physics, January 2019



# “Collision Course”



“Theoretical Physics for Machine Learning”  
Aspen Center for Physics, January 2019



*Cue the Montage!*

# 2019: Point Cloud Learning

## Energy Flow Networks



*Set-based architecture with  
interpretable latent space:*

$$S(\mathcal{J}) = F(V_1, V_2, \dots, V_\ell)$$

$$V_a(\mathcal{J}) = \sum_{i \in \mathcal{J}} E_i \Phi_a(\hat{n}_i)$$


*Collinear singularity of QCD!*

$$dP_{i \rightarrow ig} \simeq \frac{2\alpha_s}{\pi} C_i \frac{d\theta}{\theta} \frac{dz}{z}$$



[Komiske, Metodiev, JDT, [JHEP 2019](#); see also Komiske, Metodiev, JDT, [JHEP 2018](#); code at [energyflow.network](#);  
special case of Zaheer, Kottur, Ravanbakhsh, Poczos, Salakhutdinov, Smola, [NIPS 2017](#);  
other set-based architecture in Qu, Gouskos, [PRD 2020](#); Mikuni, Canelli, [EPJP 2020](#); Dolan, Ore, [arXiv 2020](#)]

# 2019: Metric Space for Colliders

*Energy Mover's Distance*




## Optimal Transport for Collider Geometry



[Komiske, Metodiev, JDT, [PRL 2019](#); code at Komiske, Metodiev, JDT, [energyflow.network](#); open data study in Komiske, Mastandrea, Metodiev, Naik, JDT, [PRD 2020](#)]  
[based on Peleg, Werman, Rom, [IEEE 1989](#); Rubner, Tomasi, Guibas, [ICCV 1998](#), [ICJV 2000](#); Pele, Werman, [ECCV 2008](#); Pele Taskar, [GSI 2013](#)]  
[flavored variant in Crispim Romão, Castro, Milhano, Pedro, Vale, [EPJC 2021](#); linearized and unbalanced transport in Cai, Cheng, Craig, Craig, [PRD 2020](#), [PRD 2022](#)]

# 2020: Metric Space in Open Data

*t-SNE Projection*




[Komiske, Mastandrea, Metodiev, Naik, JDT, [PRD 2020](#);  
using van der Maaten, Hinton, [JMLR 2008](#); using [CMS Open Data](#)]



# 2020: Metric Space in Open Data



*t-SNE Projection*



*Coordinate system of QCD!*

A diagram illustrating the coordinate system of QCD. It shows a point on a horizontal axis with a vertical arrow pointing downwards. A curved arrow labeled  $\theta$  indicates a rotation around the vertical axis. A vertical arrow labeled  $z$  indicates a vertical displacement. Below the diagram is the equation:



$$dP_{i \rightarrow ig} \sim \frac{2\alpha_s}{\pi} C_i \frac{d\theta}{\theta} \frac{dz}{z}$$

[Komiske, Mastandrea, Metodiev, Naik, JDT, PRD 2020;  
using van der Maaten, Hinton, JMLR 2008; using CMS Open Data]



# 2020: Metric Space in Open Data

## Correlation Dimension



[Komiske, Mastandrea, Metodiev, Naik, JDT, [PRD 2020](#);  
 using Grassberger, Procaccia, [PRL 1983](#); Kégl, [NIPS 2002](#); using [CMS Open Data](#)]


*Scaling behavior of QCD!*

$$dP_{i \rightarrow ig} \simeq \frac{2\alpha_s}{\pi} C_i \frac{d\theta}{\theta} \frac{dz}{z}$$

*Hmm, detector effects  
 distort underlying physics...*




# 2020: Detector Unfolding



OmniFold




*Multi-dimensional unbinned detector corrections  
via iterated application of machine-learned reweighting*



[Andreassen, Komiske, Metodiev, Nachman, JDT, [PRL 2020](#); + Suresh, [ICLR SimDL 2021](#);  
Komiske, McCormack, Nachman, [PRD 2021](#); see unfolding comparison in Petr Baron, [APPB 2021](#)]  
[see alternative in Bellagente, Butter, Kasieczka, Plehn, Rousselot, Winterhalder, Ardizzone, Köthe, [SciPost 2020](#)]



**Maintain a physical  
distance of 1.5 m**



**1.5 Metres  
(or about 3 koalas)**




For updates, visit  
[covid19.act.gov.au](https://covid19.act.gov.au)





*Oh, that...*

# 2021: Launch of IAIFI





*Can we finally put the “AI” in Altarelli-Parisi?*



$$dP_{i \rightarrow ig} \sim \frac{2\alpha_s}{\pi} C_i \frac{d\theta}{\theta} \frac{dz}{z}$$


Must separate **quarks** from **gluons**!



# 2022: Putting All the Pieces Together



Quark and Gluon Jets  
from the Strong Force





...Confronted with  
Public Collider Data...





...Unfolded for  
Detector Effects...



...Disentangled using Weak Supervision on  
Set-Based Classifiers with Topic Modeling...



...Triangulated with  
Optimal Transport!






[Komiske, Kryhin, JDT, [arXiv 2022](#); using Komiske, Mastandrea, Metodiev, Naik, JDT, [PRD 2020](#); Andreassen, Komiske, Metodiev, Nachman, JDT, [PRL 2020](#); + Suresh, [ICLR SimDL 2021](#); Metodiev, Nachman, JDT, [JHEP 2017](#); Komiske, Metodiev, JDT, [JHEP 2019](#); Metodiev, JDT, [PRL 2018](#); Komiske, Metodiev, JDT, [JHEP 2018](#); Komiske, Metodiev, JDT, [PRL 2019](#)]



# 2022: Putting All the Pieces Together



Plus some technical details as to how we extract reducibility factors, determine correlated uncertainties, and validate the method...





Once you know *quark fractions*, the rest is *linear algebra*

[Komiske, Kryhin, JDT, [arXiv 2022](#); using Komiske, Mastandrea, Metodiev, Naik, JDT, [PRD 2020](#); Andreassen, Komiske, Metodiev, Nachman, JDT, [PRL 2020](#); + Suresh, [ICLR SimDL 2021](#); Metodiev, Nachman, JDT, [JHEP 2017](#); Komiske, Metodiev, JDT, [JHEP 2019](#); Metodiev, JDT, [PRL 2018](#); Komiske, Metodiev, JDT, [JHEP 2018](#); Komiske, Metodiev, JDT, [PRL 2019](#)]



# Dimensionality of Quark/Gluon Jets



First Principles  
QCD Calculation:

$$\dim_{IJ}(Q) \simeq \frac{2\alpha_s}{\pi} (C_I + C_J) \ln \frac{p_T/2}{Q}$$



*Color Factors of QCD!*

$$dP_{i \rightarrow ig} \simeq \frac{2\alpha_s}{\pi} C_i \frac{d\theta}{\theta} \frac{dz}{z}$$

[Komiske, Kryhin, JDT, arXiv 2022]




# Dimensionality of Quark/Gluon Jets




[Komiske, Kryhin, JDT, arXiv 2022]





*We gained new insights into strong force by fusing advances in machine learning with insights from quantum field theory*

# Summary




## 2017: A Quark/Gluon Conundrum

*Quark/gluon jets offer an extreme example where fully supervised learning is fundamentally ambiguous*



## 2018: Leveraging Weak Supervision


*Assuming that “quark” and “gluon” jet categories exist, one can use machine learning to operationally define them*




## 2019-2022: The Strong Force Revisited

*Jet physics has crossed a important threshold where machine learning is now yielding insights that go beyond traditional analysis techniques*


# *Backup Slides*










# QCD through the Lens of Condensed Matter




[Komiske, Moult, JDT, Zhu, arXiv 2022; see talks by Moult, BOOST 2019, BOOST 2020]



# QCD through the Lens of Cosmology



CMS Open Data,  $R_L \in (0.3, 0.4)$



[Chen, Moult, JDT, Zhu, arXiv 2022]



# Principles of Fundamental Physics

*Robustness of Energy Flow*

[Komiske, Metodiev, JDT, JHEP 2018]



Patrick Komiske



Eric Metodiev



SF




**Power of Artificial Intelligence**

*Point Cloud Learning*

[Zaheer, Kottur, Ravanbakhsh, Poczos, Salakhutdinov, Smola, NIPS 2017]

# Energy Flow Networks


<https://energyflow.network/>  
[Komiske, Metodiev, JDT, JHEP 2019]



# Triangulating the Space of Jets





[Chu, MIT News July 2019]



[Komiske, Metodiev, JDT, [PRL 2019](#); code at Komiske, Metodiev, JDT, [energyflow.network](#);  
see alternative graph network approach in Mullin, Pacey, Parker, White, Williams, [JHEP 2021](#)]

# Rapidity-Dependent Quark/Gluon Fraction




[Komiske, Kryhin, JDT, arXiv 2022]




# Forward/Central Differences


## Multiplicity



## Weak Classification




## Weak Regression




[Komiske, Kryhin, JDT, arXiv 2022]

# Extracted Fractions with Different Observables


## Traditional Observables



## Weak Supervision




## Quark Fractions




[Komiske, Kryhin, JDT, arXiv 2022]

# Three Different Strategies for K




**Anchor Bin**



**Likelihood Ratio Fit**




**ROC Curve Fit**









[Komiske, Kryhin, JDT, arXiv 2022]

# Six Traditional Observables




[Komiske, Kryhin, JDT, arXiv 2022]

# Testing Sample Independence



[Komiske, Kryhin, JDT, arXiv 2022]

# Unfolding Checks



# Likelihood Ratio Trick

Many HEP problems can be expressed in this form!

Key example of *simulation-based inference*

Goal: Estimate  $p(x)$  /  $q(x)$

Training Data: Finite samples  $P$  and  $Q$

Learnable Function:  $f(x)$  parametrized by, e.g., neural networks

Loss Function(al):  $L = -\langle \log f(x) \rangle_P + \langle f(x) - 1 \rangle_Q$

Asymptotically:  $\arg \min_{f(x)} L = \frac{p(x)}{q(x)}$  *Likelihood ratio*

$-\min_{f(x)} L = \int dx p(x) \log \frac{p(x)}{q(x)}$  *Kullback–Leibler divergence*

[see e.g. Cranmer, Pavez, Loupe, [arXiv 2015](#); D’Agnolo, Wulzer, [PRD 2019](#); simulation-based inference in Cranmer, Brehmer, Loupe, [PNAS 2020](#); relation to f-divergences in Nguyen, Wainwright, Jordan, [AoS 2009](#); Nachman, Thaler, [PRD 2021](#)]

# Likelihood Ratio Trick

Many HEP problems can be expressed in this form!

Key example of *simulation-based inference*

Asymptotically, same structure as *Lagrangian mechanics!*

Action:  $L = \int dx \mathcal{L}(x)$

Lagrangian:  $\mathcal{L}(x) = -p(x) \log f(x) + q(x)(f(x) - 1)$

Euler-Lagrange:  $\frac{\partial \mathcal{L}}{\partial f} = 0$

Solution:  $f(x) = \frac{p(x)}{q(x)}$

Requires shift in focus from solving problems to specifying problems

[see e.g. Cranmer, Pavez, Louppe, [arXiv 2015](#); D'Agnolo, Wulzer, [PRD 2019](#); simulation-based inference in Cranmer, Brehmer, Louppe, [PNAS 2020](#); relation to f-divergences in Nguyen, Wainwright, Jordan, [AoS 2009](#); Nachman, Thaler, [PRD 2021](#)]

# Machine Learning Requirements

*If you have in hand...*

Well-specified loss  
Reliable training data  
Learnable function

*...then you can leverage ML!*

Many HEP tasks can be phrased in this language

Physics input essential for robust usage of these tools

[see [HEPML-LivingReview](#) for extensive bibliography]

