

Caching in or falling back at the Sevilleta

Sora Kim¹ and Justin Yeakel¹

¹*School of Natural Sciences, University of California, Merced, Merced, CA 95340, USA*

I. MODEL DESCRIPTION

We constructed a two-site stage-class model that tracked shark population dynamics over time, where one site was designated the nursery. We considered four key dynamics to characterize changes in population size for both sites: reproduction, somatic growth, mortality, and migration. Vital rates controlling these dynamics were site- and temperature-specific, where temperatures oscillated annually.

Reproduction: The per-capita reproductive rates $r_{n,a}$ are assumed to differ between the nursery (n subscript) and adult (a subscript) site. We set the reproductive rate at the adult site $r_a = 0$, whereas the reproductive rate at the nursery was scaled to seasonal ocean temperature, such that $r_n = 0.47 \times 10^{-7}$ female pups per second during the warmest part of the year, and 1/2 that during the coldest.

Somatic growth: Partitioning of metabolism B between growth and maintenance purposes can be used to derive a general equation for both the growth trajectories and growth rates of organisms ranging from bacteria to metazoans [1–5]. This relation is derived from the balance condition $B_0 m^\eta = E_m \dot{m} + B_m m$, [1–5] where E_m

is the energy needed to synthesize a unit of mass, B_m is the metabolic rate to support an existing unit of mass, and m is the mass of the organism at any point in its development. This balance has the general solution [5?]

$$\left(\frac{m(t)}{M}\right)^{1-\eta} = 1 - \left[1 - \left(\frac{m_0}{M}\right)^{1-\eta}\right] e^{-a(1-\eta)t/M^{1-\eta}}, \quad (1)$$

where, for $\eta < 1$, $M = (B_0/B_m)^{1/(1-\eta)}$ is the asymptotic mass, $a = B_0/E_m$, and m_0 is mass at birth. We now use this solution to define the timescale for reproduction and recovery from starvation (Fig. ??; see [2] for a detailed presentation of these timescales). The time that it takes to reach a particular mass ϵM is given by the timescale

$$\tau(\epsilon) = \ln \left[\frac{1 - (m_0/M)^{1-\eta}}{1 - \epsilon^{1-\eta}} \right] \frac{M^{1-\eta}}{a(1-\eta)}, \quad (2)$$

where ϵ is used to partition growth into discrete size classes.

Mortality:

Migration:

-
- [1] G. B. West, J. H. Brown, and B. J. Enquist, *Nature* **413**, 628 (2001).
 [2] M. E. Moses, C. Hou, W. H. Woodruff, G. B. West, J. C. Nekola, W. Zuo, and J. H. Brown, <http://dx.doi.org.proxy.lib.sfu.ca/10.1086/679735> **171**, 632 (2008).
 [3] J. F. Gillooly, E. L. Charnov, G. B. West, V. M. Savage,

- and J. H. Brown, *Nature* **417**, 70 (2002).
 [4] C. Hou, W. Zuo, M. E. Moses, W. H. Woodruff, J. H. Brown, and G. B. West, *Science* **322**, 736 (2008).
 [5] C. P. Kempes, S. Dutkiewicz, and M. J. Follows, *PNAS* **109**, 495 (2012).