

Homework 7

Jonathan Zhang EID: jdz357

Exercise (8.1). *If A is positive definite, show that its eigenvalues are positive. If A is symmetric and has positive eigenvalues, then A is positive definite.*

Proof. Suppose that A is positive definite. Let (λ, v) be an eigenpair for A . Then by the positive definiteness,

$$0 < (v, Av) = (v, \lambda v) = \lambda (v, v) = \lambda \|v\|^2.$$

Since this quantity is strictly positive, we must have $\lambda > 0$.

On the other side, assume now that A has all positive eigenvalues. Then, A admits a diagonalization $A = U\Lambda U^*$ for some $U^* = U^{-1}$ unitary matrix. Pick $x \neq 0$, and set $y = U^*x \neq 0$. Then

$$(x, Ax) = (x, U\Lambda U^*x) = (y, \Lambda y).$$

Expand $y = \sum y_i e_i$, then

$$(y, \Lambda y) = \sum_i \bar{y}_i y_i \lambda_i > 0$$

since $\lambda_i > 0$ and $\bar{y}_i y_i = |y_i|^2 \geq 0$ since $y \neq 0$ by assumption. This shows that $(x, Ax) = (y, \Lambda y) > 0$ and A is positive definite.

□

Exercise (8.3). Suppose we wish to find $u \in H^2(\Omega)$ satisfying $-\Delta u + cu^2 = f \in L^2(\Omega)$ for $\Omega \subset \mathbb{R}^d$ is a bounded Lipschitz domain. For consistency, we would require that $cu^2 \in L^2(\Omega)$. Determine the smallest p such that if $c \in L^p(\Omega)$, then this holds. Depends on d .

Proof. We want $cu^2 \in L^2(\Omega)$. Suppose that $u \in L^r(\Omega)$ and $c \in L^p(\Omega)$. Then by Generalized Hölder's, we have the estimate

$$\|cu^2\|_2^2 = \int_{\Omega} c^2 u^4 \leq \|c\|_p^2 \|u\|_r^4$$

provided that $2/p + 4/r = 1$. That is, given r , we determine $p = \frac{2}{1-4/r}$, and if $c \in L^p$, then we have $cu^2 \in L^2(\Omega)$.

We make use now of the Sobolev embedding theorem. We know $u \in H^2(\Omega) = W^{2,2}(\Omega)$. Let $j = 0, m = 2, p = 2$. Then we have

1. If $4 < d$, then $W^{2,2}(\Omega) \hookrightarrow W^{0,r}(\Omega)$ for every $r \leq \frac{2d}{d-4}$. Let us first investigate when $r = \frac{2d}{d-4}$. Substituting in, we find $p = \frac{2d}{8-d}$. We immediately find that we have problems when $d > 8$, which is admissible in this case. Restricting to $p \geq 1$, we obtain $r \geq 4$, so we must have $r \in [4, \frac{2d}{d-4}]$.
2. If $d < 4$, then $W^{2,2}(\Omega) \hookrightarrow C_B^0(\Omega) = C^0(\Omega) \cap L^\infty(\Omega)$. In this case, we obtain $r = \infty$, and calculate $p = 2$, so c can be in L^2 .

□

Exercise (8.5). Suppose that $\Omega \subset \mathbb{R}^d$ is a smooth, bounded, connected domain. Let

$$H = \left\{ u \in H^2(\Omega) : \int_{\Omega} u = 0, \nabla u \cdot n = 0 \text{ on } \partial\Omega \right\}.$$

Show that H is a Hilbert space, and prove there exists $C > 0$ such that

$$\|u\|_{H^1(\Omega)} \leq C \sum_{|\alpha|=2} \|D^{\alpha}u\|_{L^2(\Omega)}$$

for every $u \in H$.

Proof. 1. Hilbert Space.
 2. Proof by contradiction.

□

Exercise (8.8). $\Omega \subset \mathbb{R}^d$ bounded, connected, Lipschitz domain. Let $V \subset \Omega$ have positive measure. Let $H = \{u \in H^1(\Omega) : u|_V = 0\}$.

1. Why is H a Hilbert space?
2. Prove that there exists $C > 0$ such that

$$\|u\| \leq C \|\nabla u\|$$

for every $u \in H$.

1. H is the null space of the restriction to V operator. Since restriction is continuous, then H is closed. As a closed subspace of Hilbert space $H^1(\Omega)$, then H is itself a Hilbert space.
2. Go by contradiction. Assume to the contrary that there exists a sequence $u_n \in H$ such that

$$\|u_n\| = 1 \quad \text{and} \quad \|\nabla u_n\| \rightarrow 0.$$

Now, from every bounded sequence in a Hilbert space, we can extract a weakly convergent subsequence $u_{n_k} \rightharpoonup u \in H$. Weak convergence of u_{n_k} implies weak convergence of $\nabla u_{n_k} \rightharpoonup \nabla u$ in L^2 . Since the norm is continuous we pass to the limit and we have that $\nabla u = 0$, i.e. u is a constant. But if u vanishes on V , then the only possibility is that $u \equiv 0$.

On the other side, we know from the Rellich-Kondrachov Theorem that we have $u_{n_k} \rightarrow u$ in L^2 . Convergence in L^2 also implies convergence in the norm, but since $\|u_n\| = 1$ that means $\|u\| = 1$, a contradiction.