

Homework 2

Jonathan Zhang EID: jdz357

Problem 6.9:

Suppose that $f \in L^p(\mathbb{R}^d)$ for $p \in (1, 2)$. Show that there exist $f_1 \in L^1(\mathbb{R}^d)$ and $f_2 \in L^2(\mathbb{R}^d)$ such that $f = f_1 + f_2$. Define $\hat{f} = \hat{f}_1 + \hat{f}_2$. Show that this definition is well defined, that is, independent of the choices f_1 and f_2 .

Problem 6.11:

Suppose that f and g are in $L^2(\mathbb{R}^d)$. The convolution $f * g \in L^\infty(\mathbb{R}^d)$, so it may not have a Fourier Transform. Prove that $f * g = (2\pi)^{d/2} \mathcal{F}^{-1}(\hat{f}\hat{g})$ is well defined, using the natural Fourier inverse integration formula.

Problem 6.15:

Is it possible for there to be a continuous function f defined on \mathbb{R}^d with the following properties?

1. There is no polynomial P in d variables such that $|f(x)| \leq P(x)$ for all $x \in \mathbb{R}^d$.
2. The distribution $\phi \mapsto \int \phi f$ is tempered.

Problem 6.18:

The gamma function is $\Gamma(s) = \int_0^\infty t^{s-1} e^{-t} dt$. Let $\phi \in \mathcal{S}(\mathbb{R}^d)$ and $0 < \alpha < d$.

1. Show that

Problem 6.20:

Argue that $\mathcal{D}(\mathbb{R}^d)$ is dense in \mathcal{S} . Show also that \mathcal{S}' is dense in \mathcal{D}' and that distributions with compact support are dense in \mathcal{S}' .